

38. Family SCENOPINIDAE*

By Lewis P. Kelsey

The family Scenopinidae is distributed worldwide and currently includes nearly 400 described species. The Australasian/Oceanian fauna contains 86 species, mostly from Australia. Six of the 17 known genera are found in the Australasian/Oceanian Regions, including 2 worldwide genera (*Scenopinus*, *Metatrachia*) and 4 genera endemic to Australia (*Neopseudotrichia*, *Paramonova*, *Paratrachia*, *Riekiella*).

Most of the species represented have a rather limited distributional pattern restricted at or near the type locality. However, there are 4 species with wide distributions undoubtedly associated with the movement of trade goods (*Scenopinus fenestralis* and *S. glabifrons* are cosmopolitan, *S. lucidus* is from Egypt, Nigeria, and the Hawaiian Islands, and *S. papuanus* is from Hawaii, Taiwan, New Guinea, and Sri Lanka).

Very little is known of the larval biology except that they seem to be predators of other insects. Larvae have been found associated with stored-grain pests, in nests of birds and rodents, in burrows of beetle larvae attacking a rather wide range of trees and shrubs, and in soil in association with theriid larvae, which they closely resemble. Adults are taken in the field by sweeping flowers with open nectaries that are accessible to the sponging mouthparts of the adult. In buildings they are taken at windows, giving rise to the common name "window flies" for species of this family.

Refs.: Kelsey (1969, 1970a,b, 1987, rev., keys), Paramonov (1955g, syst., biol.), Krivosheina (1980, biol.).

Genus METATRICHIA Coquillett

METATRICHIA Coquillett, 1900d: 500. Type species: *Scenopinus bulbosa* Osten Sacken, 1877, orig. des.
papuana Kelsey, 1970a: 44. PNG (PNG).
waterhousei Paramonov, 1955g: 640 (*Pseudomphrale*). Australia (NSW).

Genus NEOPSEUDOTRICHIA Kelsey

NEOPSEUDOTRICHIA Kelsey, 1969: 299. Type species: *Neopseudotrichia kewi* Kelsey, 1969, orig. des.
canei Kelsey, 1969: 200. Australia (SA).
mariaensis, Paramonov, 1955, not Hardy, 1933, misid.
inglewoodi Kelsey, 1970b: 141. Australia (Qld).
kewi Kelsey, 1969: 300. Australia (ACT); Australia (NSW).
mariaensis Hardy, 1933b: 419 (*Pseudotrichia*). Australia (Tas). N. comb.
monteithi Kelsey, 1970b: 144. Australia (Qld); Australia (NSW).

Genus PARAMONOVA Kelsey

PARAMONOVA Kelsey, 1970b: 125. Type species: *Riekiella quadrifida* Kelsey, 1969, orig. des.
arila Kelsey, 1987: 10. Australia (Qld).
brevipalpa Kelsey, 1970b: 127. Australia (NSW).
bungabandii Kelsey, 1975: 205. Australia (WA).
burrijunana Kelsey, 1987: 11. Australia (WA).
bushi Kelsey, 1971: 203. Australia (WA).
collessi Kelsey, 1969: 307 (*Riekiella*). Australia (NSW). N. comb.

*Published as Miscellaneous Paper No. 1146 of the Delaware Agricultural Experiment Station, Contribution No. 573 of the Department of Entomology and Applied Ecology, University of Delaware, Newark, Delaware.

complexa Kelsey, 1969: 309 (*Riekiella*). Australia (ACT). **N. comb.**
dycei Kelsey, 1969: 310 (*Riekiella*). Australia (NSW). **N. comb.**
emarginata Kelsey, 1970b: 129 (*Riekiella*). Australia (WA).
fonsamnicula Kelsey, 1980: 9. Australia (NSW).
hamata Kelsey, 1980: 10. Australia (NSW).
mangara Kelsey, 1987: 12. Australia (WA).
poocherania Kelsey, 1987: 13. Australia (SA).
quadrifida Kelsey, 1969: 313 (*Riekiella*). Australia (Qld). **N. comb.**
taylori Kelsey, 1969: 315 (*Riekiella*). Australia (Vic). **N. comb.**
uncata Kelsey, 1969: 315 (*Riekiella*). Australia (NSW). **N. comb.**
zentae Kelsey, 1975: 207. Australia (SA).

Genus PARATRICHIA Kelsey

PARATRICHIA Kelsey, 1969: 320. Type species: *Paratrichia lobosa* Kelsey, 1969, orig. des.
lobosa Kelsey, 1969: 320. Australia (ACT).
spicata Kelsey, 1975: 208. Australia (WA).
westralica Paramonov, 1955g: 419 (*Pseudomphrale*). Australia (WA). **N. comb.**

Genus RIEKIELLA Paramonov

RIEKIELLA Paramonov, 1955g: 652. Type species: *Riekiella bicornis* Paramonov, 1955, orig. des.
arilaringa Kelsey, 1987: 16. Australia (WA).
arnheimi Kelsey, 1975: 210. Australia (NT).
bicornis Paramonov, 1955g: 653. Australia (Qld).
binnaliweanda Kelsey, 1987: 18. Australia (WA).
bolgarti Kelsey, 1969: 55 (*Scenopinus*). Australia (WA). **N. comb.**
burririmbaja Kelsey, 1987: 19. Australia (NT).
cetratra Kelsey, 1971: 196. Australia (NT).
cornuta Kelsey, 1971: 197. Australia (NT).
culbara Kelsey, 1987: 20. Australia (NT).
cuspidata Kelsey, 1971: 198. Australia (Qld).
denticulata Kelsey, 1971: 200. Australia (NT).
kilyalpa Kelsey, 1987: 21. Australia (WA).
longitermina Kelsey, 1975: 211. Australia (WA).
longiventris Kelsey, 1969: 312. Australia (WA).
malkana Kelsey, 1987: 22. Australia (WA).
maloga Kelsey, 1987: 23. Australia (WA).
murilla Kelsey, 1987: 24. Australia (WA).
nancyae Kelsey, 1987: 25. Australia (WA).
nicholsi Kelsey, 1971: 200. Australia (NT).
noona Kelsey, 1987: 27. Australia (WA).
rossi Kelsey, 1969: 123. Australia (WA).
rupari Kelsey, 1987: 27. Australia (WA).
tamala Kelsey, 1987: 29. Australia (WA).
woodwardi Kelsey, 1971: 201. Australia (NT).
yoota Kelsey, 1987: 30. Australia (WA).

Genus SCENOPINUS Latreille

OMPHRALE Meigen, 1800: 29. Suppressed by I.C.Z.N., 1963b: 339.
SCENOPINUS Latreille, 1802: 463. Type species: *Musca fenestralis* Linnaeus, 1758, mon.
SCENOPOEUS Agassiz, 1847: 333 (unjust. emend. of *Scenopinus*).
LUCIDOMPHRALE Kröber, 1937: 219. Type species: *Scenopinus lucida* Becker, 1902, orig. des.

- PAROMPHRALE* Kröber, 1937: 222. Type species: *Scenopinus glabrifrons* Meigen, 1824, orig. des.
- SCENOPINULA* Paramonov, 1955g: 651. Type species: *Scenopinula pallidipennis* Paramonov, 1955, mon.
- SCAENOPIUS*, error for *Scenopinus*.
- adventicius* Hardy, 1960: 329. Hawaiian Is.
- adyeri* Kelsey, 1987: 5. Australia (WA).
- arrectus* Kelsey, 1975: 198. Australia (WA)
- ascensus* Kelsey, 1987: 6. Australia (Qld).
- bicuspis* Kelsey, 1971: 185. Australia (NT).
- biroi* Kertész, 1899b: 173. PNG (PNG); Australia (NSW, Qld).
- brittoni* Kelsey, 1970: 111. Australia (WA).
- casurarinus* Kelsey, 1975: 200. Australia (NT).
- cristafrons* Kelsey, 1971: 186. Australia (NT).
- darwini* Kelsey, 1969: 64. Australia (NT).
- dilkara* Kelsey, 1987: 7. Australia (WA).
- erugatus* Kelsey, 1975: 201. Australia (WA).
- fenestralis* Linnaeus, 1758: 597 (*Musca*). "Europa"; Australia (Qld); cosmopol.
- perkinsi*, Paramonov, 1955, not Hardy, 1942, misid.
- fijianus* Kröber, 1939a: 399 (*Omphrale*). Fiji.
- fragosifrons* Kelsey, 1969: 68. Australia (WA).
- glabrifrons* Meigen, 1824: 114. Austria; Australia (NSW, Qld, WA); cosmopol.
- civiculus* Hardy, 1933b: 418. Australia (NSW & Qld & SA).
- harleyi* Kelsey, 1971: 187. Australia (Qld).
- limbunyai* Kelsey, 1971: 190. Australia (NT).
- lucidus* Becker, 1902: 39. Egypt; Hawaiian Is; Algeria, Christmas I (Indian Ocean), Afrotrop. Reg.
- lunulanotafrons* Kelsey, 1971: 191. Australia (NT).
- mendumae* Kelsey, 1971: 188. Australia (NT).
- millarae* Kelsey, 1971: 189. Australia (Qld).
- minutus* Kelsey, 1969: 78. Australia (NSW).
- nakkari* Kelsey, 1987: 8. Australia (WA).
- norrisi* Kelsey, 1969: 82. Australia (Qld).
- pallidipennis* Paramonov, 1955g: 651 (*Scenopinula*). Australia (NSW); Australia (SA). N. comb.
- papuanus* Kröber, 1912e: 25 (*Omphrale*). PNG (PNG) & Taiwan; Guam, Hawaiian Is, Northern Marianas; Christmas I (Indian Ocean), Cocos (Keeling) Is, Sri Lanka.
- zeylanicus* Senior-White, 1922a: 205. Sri Lanka.
- niger*, Grimshaw, 1901, not De Geer, 1776, misid.
- perkinsi* Hardy, 1942: 202. Australia (Qld); Australia (NSW).
- renneri* Kelsey, 1975: 204. Australia (NT).
- sanfordi* Kelsey, 1971: 192. Australia (NT).
- spurrelli* Kelsey, 1969: 90. Australia (Vic).
- torrensianus* Kelsey, 1980: 7. Australia (SA).
- trigelasinus* Kelsey, 1971: 194. Australia (NT).
- triquetrus* Kelsey, 1971: 192. Australia (NT).