

SPOTTED REDSHANK

Tringa erythropus

non-breeding visitor, vagrant

monotypic

The Spotted Redshank breeds in boreal regions of Eurasia from Scandinavia to the Russian Far east, and winters in S Europe through c. Africa and SE Asia (Dement'ev and Gladkov 1951c, Cramp and Simmons 1983, AOU 1998). Vagrants have been recorded in the Pacific Basin to Australia, Guam, and Chuuk (Pratt et al. 1987, Higgins and Davies 1996, Wiles 2005) and at scattered locations throughout North America (AOU 1998). In the Hawaiian Islands there is a single record from the *Southeastern Hawaiian Islands*, a first-year bird that over-wintered at the at Kona (Kealakehe) STP, **Hawai'i I** 3 Nov 2014-15 Feb 2015 ([HRBP](#) 6424-6427, published *NAB* 69:514). Based on this record, Spotted Redshank was accepted by the [HBRC](#) to the Hawaiian Island Checklist in Jul 2015.

[Acronyms and Abbreviations](#)

[Literature cited](#)

Citation: Pyle, R.L., and P. Pyle. 2017. The Birds of the Hawaiian Islands: Occurrence, History, Distribution, and Status. B.P. Bishop Museum, Honolulu, HI, U.S.A. Version 2 (1 January 2017) <http://hbs.bishopmuseum.org/birds/rlp-monograph/>