

## TEREK SANDPIPER

*Xenus cinereus*

**non-breeding visitor, vagrant**

monotypic

The Terek Sandpiper breeds across c. Russia and winters in the S Hemisphere from S Africa to Australia (Dement'ev and Gladkov 1951c, Cramp and Simmons 1983, Higgins and Davies 1996, AOU 1998). Migrants and vagrants have been recorded E in the Pacific Basin to Yap (Pratt et al. 1987, Wiles 2005) and in Alaska, British Columbia, and California (AOU 1998, CBRC 2007). In the Hawaiian Islands there is a single record from the *Southeastern Hawaiian Islands*, a bird present in a flooded field near Kahuku, *O'ahu* 9 Jan-6 Mar 2014 (Pohlman and VanderWerf 2014; [HRBP](#) 6399-6402 published *NAB* 68:444). Terek Sandpiper was accepted by the [HBRC](#) to the Hawaiian Island Checklist in Sep 2014.

[Acronyms and Abbreviations](#)

[Literature cited](#)

Citation: Pyle, R.L., and P. Pyle. 2017. The Birds of the Hawaiian Islands: Occurrence, History, Distribution, and Status. B.P. Bishop Museum, Honolulu, HI, U.S.A. Version 2 (1 January 2017) <http://hbs.bishopmuseum.org/birds/rlp-monograph/>