

Numbers of Hawaiian Species: Supplement 1¹

SCOTT E. MILLER and LUCIUS G. ELDREDGE² (Hawaii Biological Survey, Bishop Museum, P.O. Box 19000, 1525 Bernice Street, Honolulu, Hawaii 96817, USA)

This is a supplement to our earlier tabulation of species known from Hawaii (Eldredge & Miller 1995; see also Mlot 1995). The Hawaii Biological Survey is currently working on species checklists and bibliographies for many of these taxa, so this should be viewed as an interim report subject to further change. See Eldredge & Miller (1995) for definitions and scope. We have not included bacteria and viruses, but have now included human parasitic protists and helminths. Allison *et al.* (1995, table 21.3) summarized the numbers of marine species based on Eldredge & Miller (1995).

From literature and unpublished sources, approximately 22,056 species have been recorded from the Hawaiian Islands and surrounding waters. Of these, 8,850 are endemic to the Hawaiian Islands, and 4,465 are nonindigenous species of protists, fungi, plants and animals. These numbers include significant refinements from the previous compilation by Eldredge & Miller (1995), based on continued literature survey and review by specialists, especially among the parasitic Protozoa and helminths. Many gaps in knowledge of Hawaii's biota remain and many species of protists, algae, fungi, worms and arthropods remain to be described.

Table 1. Estimates of numbers of species of the Hawaiian biota (based on Eldredge & Miller 1995, as updated by this paper and other papers in this issue of *Records of the Hawaii Biological Survey*).

Taxon	Total	Endemic	NIS
Algae and other protists	2228	4	5?
Fungi and lichens	2023	240	0?
Flowering plants	1910	850	877
Other plants	703	243	50
Mollusks	1656	956	75+
Insects	7862	5237	2527
Other arthropods	1795	335	519
Other invertebrates	2338	785	250
Fish	1195	139	73
Amphibians	4	0	4
Reptiles	24	0	20
Birds	274	60	46
Mammals	44	1	19
Total	22056	8850	4465

Endemic species are restricted to Hawaii; nonindigenous (= alien, = exotic, includes introduced) species do not naturally occur in Hawaii; indigenous species occur naturally in Hawaii but are not endemic. See Eldredge & Miller 1995 for additional definitions and qualifications.

1. Contribution No. 1996-001 to the Hawaii Biological Survey.

2. Executive Secretary, Pacific Science Association, P.O. Box 17801, Honolulu, Hawaii, 96817, USA.

Freshwater and terrestrial Cyanophyta (blue-green algae)

We have not yet reconciled the many old records (e.g., MacCaughey 1918, Tilden 1901, 1910) with current nomenclature. Some recent records include Doty & Watson 1981, MacEntee & Bold 1978, and Zimmerman & Bergman 1990. Patterson *et al.* (1991 and included references) included many Hawaiian species in a phytochemical screening program. The recorded flora of freshwater and terrestrial forms likely approaches 100 species.

Freshwater and terrestrial algae

We have not yet reconciled the many old records (e.g., MacCaughey 1918, Tilden 1901) with current nomenclature, but our original estimate of over 120 species was too low. Recent authoritative records include over 172 diatoms (Fungladda *et al.* 1983), 26 desmids (Prescott *et al.* 1975–1983), and 34 macroalgae in streams (Vis *et al.* 1994). The total likely exceeds 250 species.

Protozoa

The protists of the Hawaiian islands have never been systematically surveyed, but we have accumulated records of 207 species (excluding Foraminifera) from a variety of sources. Only authoritative identifications have been included. We expect that the ultimate number of species will be vastly greater.

Free-living marine protozoa: 39 species (Phillips 1977, Larsen & Patterson 1990), in addition to over 1000 species of Foraminifera (Chave 1987).

Free-living freshwater protozoa: 61 species, including 59 testaceous rhizopods (Van Winkle 1927), *Paramecium primaurelia* Sonneborn 1975, and *Tetrahymena tropicalis* Nanney & McCoy 1976 (= *T. pyriformis* variety 9 of Elliott *et al.* 1964).

Free-living soil protozoa: At least 31 ciliates, 9 flagellates, 6 Sarcodina, and 10 Testacea recorded from various habitats on Hawaii Island (Bamforth & Eggler 1973, Foissner 1993, 1994).

Parasitic protozoa of terrestrial vertebrates including humans: 60 species. Alicata (1969) detailed the species known from humans and domestic animals. Other key references include: Colgrove (1976 — dog), Metcalf (1923 — amphibian), Upton *et al.* (1994 — reptiles), Wallace & Frenkel (1975 — cat), Van Hoven *et al.* (1992 — deer), and Van Riper (1991 — birds).

Parasitic protozoa of invertebrates: 11 species symbionts in 3 species of termites (Kirby 1941–1946, Lai *et al.* 1983), at least 9 species from insects (Fujii & Tamashiro 1967, Tamashiro 1968, Tanabe & Tamashiro 1967, Tanada 1957), 2 species from marine crabs (Ball 1963).

Parasitic protozoa of marine fishes: 8 species (Basson *et al.* 1990, Lom 1970, Moser & Noble 1977a, b).

Fungi

We have not yet tabulated the number of fungi beyond our original estimates, but it is worth pointing to the recent work on yeasts associated with flowering plants (Starmer 1981), especially *Opuntia* cactus (Starmer *et al.* 1990); the recent reviews of rust fungi (over 74 species — Gardner & Hodges 1989, Gardner 1994); and Agaricales (101 taxa — Desjardin *et al.* 1992).

Platyhelminthes (zooparasitic)

Of 583 species, 56 are cestodes (tapeworms) including 20 from marine fishes (Cornford 1974, Carvajal *et al.* 1976, Beveridge & Campbell 1993, Brill *et al.* 1987, Deardoff *et al.* 1984, Gulyaev & Korotaeva 1989, Yamaguti 1968a), 1 from freshwater fishes (Font & Tate 1994), 21 from terrestrial vertebrates (including human) (Alicata 1969, Olsen & Haas 1976, Lewis & Holmes 1971), of 2 species from intermingling mammals 1 is new record (McKenzie & Davidson), 7 from green turtles (Dailey *et al.* 1992), 1 from porpoise (Testa & Dailey 1977), 4 from Hawaiian monk seal (Anderson 1987), and 1 from “megamouth” shark (Dailey & Vogelhein 1982); the remaining 527 species are trematodes (flukes), 507 from marine fishes (Lebedev 1984, Mamaev 1989, Oliver 1983, Yamaguti 1965, 1968b, 1970), 16 from terrestrial vertebrates (including human) (Alicata 1969, O’Leary *et al.* 1985), 1 unidentified species (larva) from freshwater fish (Font & Tate 1994), 1 from nonindigenous, marine, cage-cultured tilapia (Kaneko *et al.* 1988), 1 species from the cane toad (Speare 1990), 1 species with secondary intermediate stage in coral *Porites compressa* (Aeby 1991).

Acanthocephala

Three species not reported in 1995: *Moniliformis moniliformis* from rats (Alicata 1969) and cockroach (Schaefer 1970); *Corynosoma rauschi* from Hawaiian monk seal (Dailey *et al.* 1988); and *Mediorhynchus orientalis* from birds (Schmidt & Kuntz 1977); also unidentified species from freshwater fish (Font & Tate 1994).

Nematoda (zooparasitic)

Of 131 zooparasitic species, 93 reported from terrestrial vertebrates (including man) (Alicata 1969), of 22 species reported from intermingling mammals 12 are new records (McKenzie & Davidson 1989), 1 species from Hawaiian monk seal (Dailey 1988), 1 from albatross (Langston & Hillgarth 1995), 3 from endemic birds (Cid del Prado *et al.* 1985), 3 from gamebirds (Lewin & Holmes 1971, Smith 1973), 18 from marine fishes (Bruce *et al.* 1994, Deardoff & Overstreet 1982, Deardoff *et al.* 1982, 1983, Deardoff 1987, Noble 1966, Solovjeva & Pozdnyakov 1984), and 1 from freshwater fish (Font & Tate 1994).

Tardigrada

Pollock (1995) published the first record of 2 species of marine tardigrades—*Dipodartus borrori* and *D. anaholiensis*—from Anahola Bay, Kauai.

Bryozoa

Two Hawaiian freshwater bryozoan species were overlooked in our earlier report (Bailey-Brock & Hayward 1984); one new deepwater, marine species described (Chimonides & Cook 1994).

Mollusca

Cowie *et al.* (1995) previously cited as “in press” has been published; the 75 non-indigenous species in the table represents non-marine mollusks only— numbers of non-indigenous marine mollusks have not yet been ascertained (R.H. Cowie, pers. comm.).

Annelida: Hirudinea

Two leeches collected in association with freshwater fishes to be added to species list: *Myzobdella lugubris* and *Cystobranchus* sp. (Font & Tate 1994).

Arthropoda: Crustacea

Recent additions to the crustacean fauna of the Hawaiian Islands include: cyclopoid copepod *Haplostomides hawaiiensis* [n.sp.](Ooishi 1994); harpacticoid copepod *Psammopsyllus stri* (Kunz 1993); ascothoracid cirriped *Laura bicornuta* [n.sp.] and *Zoanthoeus cerebroides* [n.sp.](Grygier 1985); mysids *Anisomysis hawaiiensis* [n.sp.], *A. extranea* [n.sp.], *A. xenops* (Murano 1995), *Holmesimysis costata* (Holmquist 1979), and *Gnathophausia longispina* (Wilson & Boehlert 1993); marine isopods *Neonaesa rugosa* and *Cymodoella hawaiiensis* [n.sp.](Bruce 1994); deep-water caridean shrimp *Opaepele loihi* [n.gen. & n.sp.](Williams & Dobbs 1995).

Terrestrial and freshwater Arthropoda

The numbers for some groups have changed because of continued refinement of synonymies and distribution status in the database (G.M. Nishida, pers. comm.), as well as recent publications (Gillespie 1994—3 new endemic spiders; Asquith 1994—39 new endemic bugs). Many additional endemic species of arthropods remain to be described (e.g. Howarth 1990). Also, for the original figure for nonindigenous species of the insect order Diptera, there was a typographical error in the table. The number for NIS should read 361 rather than 631 (this did not impact the total numbers).

Insecta: 7862 total species, including 5237 endemic and 2527 nonindigenous;

Araneae: 200 total species, including 126 endemic and 70 nonindigenous;

Acari: 529 total species, including 103 endemic and 365 nonindigenous.

Symphyla: Has 5 nonindigenous species (Nishida 1994). It was overlooked in the original list.

Chordata: Reptilia

Our original list omitted 3 indigenous species of sea turtles and 3 nonindigenous species of freshwater turtles (C. Kishinami, pers. comm.).

Acknowledgements

Our literature survey was improved by access to the CAB Abstracts and AGRICOLA databases (CAB International and National Agriculture Library, U.S. Dept. of Agriculture, respectively). We thank the libraries of Bishop Museum, Santa Barbara Museum of Natural History, University of California at Santa Barbara, and University of Hawaii at Manoa for access to literature. Various specialists have assisted in literature search and analysis, or submitted reprints. The specimen database for the U.S. National Animal Parasite Collection, U.S. Department of Agriculture, Beltsville, Maryland (E.P. Hoberg & P.A. Pilitt) also helped our literature search. This is an interim product from a project supported by the John D. and Catherine T. MacArthur Foundation and National Biological Service.

Literature Cited

- Aeby, G.S.** 1991. Behavioral and ecological relationships of a parasite and its hosts within a coral reef system. *Pac. Sci.* **45**: 263–69.
- Alicata, J.E.** 1969. *Parasites of man and animals in Hawaii*. S. Karger, Basel. 190 p.
- Allison, A., S.E. Miller & G.M. Nishida.** 1995. Hawaii Biological Survey: a model for the Pacific Region, p. 349–55. In: Maragos, J.E., M.N.A. Peterson, L.G. Eldredge, J.E. Bardach & H.F. Takeuchi, eds., *Marine and coastal biodiversity in the tropical island Pacific region*. Volume 1. Species systematics and information management priorities. East-West Center, Honolulu. xxii + 424 p.
- Anderson, K.I.** 1987. A redescription of *Diphyllbothrium stemmacephalum* Cobbold, 1858 with comments on other marine species of *Diphyllbothrium* Cobbold, 1958. *J. Nat. Hist.* **21**: 411–27.
- Asquith, A.** 1994. Revision of the endemic Hawaiian genus *Sarona* Kirkaldy (Heteroptera: Miridae: Orthotylinae). *Bishop Mus. Occas. Pap.* **40**: 1–81.
- Bailey-Brock, J.H. & P.J. Hayward.** 1984. A freshwater bryozoan, *Hyalinella vaihiriae* Hastings (1929), from Hawaiian prawn farms. *Pac. Sci.* **38**: 199–204.
- Ball, G.H.** 1963. *Cephaloidophora carpilodei* n. sp. and *C. pinguis* n. sp., gregarines of a xanthid crab in Hawaii. *J. Protozool.* **10**: 321–27.
- Bamforth, S.S. & W.A. Egger.** 1973. Protozoa from pumice soils of Kilauea Iki Volcano, Hawaii. *J. Protozool.* **20**: 496. [abstract]
- Basson, L., J.G. Van As & L. Fishelson.** 1990. A new species of *Trichodina* (Ciliophora: Peritrichia) from the intestine of the surgeonfish *Acanthurus xanthopterus*. *Int. J. Parasitol.* **20**: 785–87.
- Beveridge, I. & R. A. Campbell.** 1993. A revision of *Dasyrhynchus* Pinter (Cestoda: Trypanorhyncha), parasitic in elasmobranch and teleost fishes. *Syst. Parasitol.* **24**: 129–57.
- Brill, R.W., R. Bourke, J. A. Brock & M. D. Dailey.** 1987. Prevalence and effects of infection of the dorsal aorta in yellowfin tuna, *Thunnus albacares*, by the larval cestode, *Dasyrhynchus talismani*. *Fish. Bull.* **85**: 767–76.
- Bruce, N.L.** 1994. New records of marine isopod crustaceans (Sphaeromatidae) from Hawaiian waters. *Crustaceana* **67**: 381–86.
- , **R.D. Adlard & L.R.G. Cannon.** 1994. Synoptic checklist of ascaridoid parasites (Nematoda) from fish hosts. *Invertebr. Taxon.* **8**: 583–674.
- Carvajal, J., R.A. Campbell & E.M. Cornford.** 1976. Some tyrypanorhynch cestodes from Hawaiian fishes with descriptions fo new species. *J. Parasitol.* **62**: 70–77.
- Chimonides, P.J. & P.L. Cook.** 1994. Notes on the genus *Cranosina* (Bryozoa, Cheilostomida). *Zool. Scripta* **23**: 43–49.
- Cid del Prado, Vera, I., A.R. Maggenti & C. van Riper, III.** 1985. New species of Spiruridae (Nemata: Spirurida) from endemic Hawaiian honeycreepers (Passeriformes: Drepanididae), the Japanese white-eye (Passeriformes: Zosteropidae) and a new species of Acuariidae (Nemata: Spirurida) from the Japanese white-eye collected on the island of Hawaii. *Proc. Helminthol. Soc. Wash.* **52**: 247–59.
- Colgrove, D.J.** 1976. Canine babesiosis in Hawaii. *J. Amer. Vet. Med. Assoc.* **168**: 591–92.
- Cornford, E.M.** 1974. Two tetraphyllidean cestodes from Hawaiian stingrays. *J. Parasitol.* **60**: 942–48.

- Cowie, R.H., N.L. Evenhuis & C.C. Christensen.** 1995. *Catalog of the native land and freshwater molluscs of the Hawaiian Islands*. Backhuys Publ., Leiden. vi + 248 p.
- Dailey, M.D. & W. Vogelbein.** 1982. Mixodigmatidae, a new family of cestode (Trypanorhyncha) from a deep sea, planktivorous shark. *J. Parasitol.* **68**: 145–49.
- , **M.L. Fast & G.H. Balazs.** 1992. A survey of the Trematoda (Platyhelminthes: Digenea) parasitic in green turtles, *Chelonia mydas* (L.) from Hawaii. *Bull. South. Calif. Acad. Sci.* **91**: 84–91.
- , **R.V. Santangelo & W. G. Gilmartin.** 1988. A coprological survey of helminth parasites of the Hawaiian monk seal from the northwestern Hawaiian Islands. *Mar. Mamm. Sci.* **4**: 125–31.
- Deardoff, T.L.** 1987. Redescription of *Pulchrascaris chiloscylli* (Johnston and Mawson, 1951) (Nematoda: Anisakidae), with comments on species in *Pulchrascaris* and *Terranova*. *Proc. Helminthol. Soc. Wash.* **54**: 28–39.
- & **R.M. Overstreet.** 1982. *Hysterothylacium pelagicum* sp. n. and *H. cornutum* (Stossich, 1904) (Nematoda: Anisakidae) from marine fishes. *Proc. Helminthol. Soc. Wash.* **49**: 246–51.
- , **M.M. Kliks, M.E. Rosenfeld, R.A. Rychlinski & R.S. Desowitz.** 1982. Larval ascaridoid nematodes from fishes near the Hawaiian Islands, with comments on pathogenicity experiments. *Pac. Sci.* **36**: 187–201.
- , **R.B. Raybourne & T.E. Mattis.** 1984. Infections with trypanorhynch plerocerci (Cestoda) in Hawaiian fishes of commercial importance. *Sea Grant Q.* **6**: 1–6.
- Desjardin, D.E., G.J. Wong & D.E. Hemmes.** 1992. Agaricales of the Hawaiian Islands. I. Marasmioid fungi: new species, new distributional records, and poorly known taxa. *Can. J. Bot.* **70**: 530–42.
- Doty, M.S. & L.-L. Watson.** 1981. Terrestrial algae, p. 165–71. In: Mueller-Dombois, D., K.W. Bridges & H.L. Carson, eds., *Island ecosystems: biological organization in selected Hawaiian communities*. Hutchinson Ross Publ. Co., Stroudsburg, Penn.
- Eldredge, L.G. & S.E. Miller.** 1995. How many species are there in Hawaii? *Bishop Mus. Occas. Pap.* **41**: 3–18.
- Elliott, A.M., M.A. Studier & J.A. Work.** 1964. *Tetrahymena pyriformis* from several Pacific islands and Australia. *J. Protozool.* **11**: 370–78.
- Foissner, W.** 1993. *Corticocolpoda kaneshiroae* n. g., n. sp., a new colpodid ciliate (Protozoa, Ciliophora) from the bark of ohia trees in Hawaii. *J. Euk. Microbiol.* **40**: 764–75.
- . 1994. *Bryometopus hawaiiensis* sp. n., a new colpodid ciliate from a terrestrial biotope of the Hawaiian archipelago (Protozoa: Ciliophora). *Ann. Naturhist. Mus. Wien (B)* **96**: 19–27.
- Font, W.F. & D.C. Tate.** 1994. Helminth parasites of native Hawaiian freshwater fishes: an example of extreme ecological isolation. *J. Parasitol.* **80**: 682–88.
- Fujii, J.K. & M. Tamashiro.** 1972. *Nosema tephrititae* sp. n., a microsporidian pathogen of the Oriental Fruit Fly, *Dacus dorsalis* Hendel. *Proc. Hawaii. Entomol. Soc.* **21**: 191–203.
- Fungladda, N., I. Kaczmarek & S.R. Rushforth.** 1983. A contribution to the freshwater diatom flora of the Hawaiian islands. *Biblioth. Diatomol.* **2**: 1–103.
- Gardner, D.E.** 1994. The native rust fungi of Hawaii. *Can. J. Bot.* **72**: 976–89.
- & **C.S. Hodges, Jr.** 1989. The rust fungi (Uredinales) of Hawaii. *Pac. Sci.* **43**: 41–55.

- Gillespie, R.G.** 1994. Hawaiian spiders of the genus *Tetragnatha*: III. *Tetragnatha acuta* clade. *J. Arachnol.* **22**: 161–68.
- Grygier, M.J.** 1985. Lauridae: taxonomy and morphology of ascothoracid crustacean parasites of zoanths. *Bull. Mar. Sci.* **36**: 278–303.
- Gulyaev, V.D. & V.D. Korotaeva.** 1989. [Redescription of the type species of *Glossobothrium Yamaguti*, 1952 (Cestoda, Pseudophyllidea.) *Vestn. Zool.* **1989**(6): 8–11. [in Russian]
- Holmquist, C.** 1979. *Mysis costata* Holmes, 1900, and its relations (Crustacea, Mysidacea). *Zool. Jahrb. (Syst.)* **106**: 471–99.
- Howarth, F.G.** 1990. Hawaiian terrestrial arthropods: An overview. *Bishop Mus. Occas. Pap.* **30**: 4–26.
- Kaneko, J.J., II, R. Yamada, J.A. Brock & R.M. Nakamura.** 1988. Infection of tilapia, *Oreochromis mossambicus* (Trewavas), by a marine monogenean, *Neobenedenia melleni* (MacCallum, 1927), 1963 in Kaneohe Bay, Hawaii, USA, and its treatment. *J. Fish Dis.* **11**: 295–300.
- Kirby, H.** 1941–1946. Devescovinid flagellates of termites. *Univ. Calif. Publ. Zool.* **45**: 1–318.
- Kunz, H.** 1993. Beitrag zur kenntnis von zwei *Psammopsyllus*-arten (Copepoda, Harpacticoida, Cyliindropsyllidae). *Crustaceana* **64**: 143–54.
- Lai, P.Y., M. Tamashiro & J.K. Fujii.** 1983. Abundance and distribution of the three species of symbiotic Protozoa in the hindgut of *Coptotermes formosanus* (Isoptera: Rhinotermitidae). *Proc. Hawaii. Entomol. Soc.* **24**: 271–76.
- Langston, N. & N. Hilgarth.** 1995. Moults varies with parasites in Laysan Albatross. *Proc. R. Soc. Lond. (B)* **261**: 239–43.
- Lebedev, B.I.** 1984. [New taxa of gastrocotylid-like monogenans.] *Parazity Zhivotnykh i Rastenii* [*Parasites of animals and plants*], p. 17–24. [In Russian with English summary.]
- Lewin, V. & J.C. Holmes.** 1971. Helminths from exotic game birds of the Puuwaawaa Ranch, Hawaii. *Pac. Sci.* **25**: 372–81.
- Lom, J.** 1970. Trichodinid ciliates (Peritrichida: Urceolariidae) from some marine fishes. *Fol. Parasitol.* **17**: 113–25.
- MacCaughy, V.** 1918. Algae of the Hawaiian Archipelago. *Bot. Gaz.* **65**: 42–57, 121–49.
- MacEntee, F.J. & H.C. Bold.** 1978. Some microalgae from sand. *Texas J. Sci.* **30**: 167–73.
- Mamaev, Y.** 1989. [The species composition and morphological characteristics of the genus *Microcotyle* (Microcotylidae, Monogenoidea)], p. 32–38 In: Lebedev, B.I., ed., *Parazitologicheskie Issledovaniya* [*Investigations in parasitology*]. [In Russian with English summary.]
- Margulis, L., J.O. Corliss, M. Melkonian & D.J. Chapman, eds.** 1990. *Handbook of Protoctista*. Jones and Bartlett, Boston. xli + 914 p.
- McKenzie, M.E. & W.R. Davidson.** 1989. Helminth parasites of intermingling axis deer, wild swine and domestic cattle from the island of Molokai, Hawaii. *J. Wildl. Dis.* **25**: 252–57.
- Metcalf, M.M.** 1923. The opalinid ciliate infusorians. *U.S. Natl. Mus. Bull.* **120**: 1–484.
- Mlot, C.** 1995. Biological surveys in Hawaii, taking inventory of a biological hot spot. *Science* **269**: 322–23.

- Moser, M. & E.R. Noble. 1977a. Three genera of Myxosporida (Protozoa) in macrourid fishes. *Int. J. Parasitol.* **7**: 93–96.
- & E.R. Noble. 1977b. *Zschokkella* (Protozoa: Myxosporida) in macrourid fishes. *Int. J. Parasitol.* **7**: 97–100.
- Murano, M. 1995. New and already known species of the genus *Anisomysis* (Mysidacea) from Hawaii and the Society Islands. *J. Crust. Biol.* **15**: 355–64.
- Nanney, D.L. & J.W. McCoy. 1976. Characterization of the species of the *Tetrahymena pyriformis* complex. *Trans. Am. Microscop. Soc.* **95**: 664–82.
- Nishida, G.M., ed. 1994. Hawaiian terrestrial arthropod checklist. Second edition. *Bishop Mus. Tech. Rep.* **4**: 1–287.
- Noble, E. R. 1966. A new camallanid nematode from Hawaii. *Pac. Sci.* **20**: 360–66.
- O’Leary, M.J., J.T. Berthiaume & V. Sakbun. 1985. Treatment of *Clonorchis sinensis* in Hawaii’s Laotian population—experience with praziquantel. *Hawaii Med. J.* **44**: 63–64.
- Oliver, G. 1983. *Diplectanum yamaguti* sp. n. (Monogenea, Monopisthocotylea, Diplectanidae), parasite de *Kyphosus cinerascens* (Forsskål) à Hawaii. *Zool. Scripta* **12**: 91–93.
- Olsen, O.W. & W.R. Haas. 1976. A new record of *Spirometra mansoni*, a zoonotic tapeworm, from naturally infected cats and dogs in Hawaii. *Hawaii Med. J.* **35**: 261–63.
- Ooishi, S. 1994. *Haplostomides hawaiiensis*, new species (Copepoda: Cyclopoida: Ascidiocolidae), associated with the ascidian *Polyclinum constellatum* at Honolulu, Hawaii. *Hydrobiologia* **292/293**: 89–96.
- Patterson, G.M.L., C.L. Baldwin, C.M. Bolis, F.R. Caplan, H. Karuso, L.K. Larsen, I.A. Levine, R.E. Moore, C.S. Nelson, K.D. Tschappat, G.D. Tuang, E. Furusawa, S. Furusawa, T.R. Norton & R.B. Raybourne. 1991. Antineoplastic activity of cultured blue-green algae (Cyanophyta). *J. Phycol.* **27**: 530–36.
- Pollock, L.W. 1995. New marine tardigrades from Hawaiian beach sand and phylogeny of the family Halechiniscidae. *Invert. Biol.* **114**: 220–35.
- Prescott, G.W. et al. 1975–1983. *A synopsis of North American desmids*. Part II. Desmidiaceae: Placodermae. Sections 1–5. University of Nebraska Press, Lincoln. [coauthors vary]
- Schaefer, P.W. 1970. *Periplanta americana* (L.) as intermediate host of *Moniliformis moniliformis* (Bremser) in Honolulu, Hawaii. *Proc. Helminthol. Soc. Wash.* **37**: 204–06.
- Schmidt, G.D. & R.E. Kuntz. 1977. Revision of *Mediorhynchus* Van Cleave 1916 (Acanthocephala) with a key to species. *J. Parasitol.* **63**: 500–07.
- Schottelius, J. 1989. Contribution to the characterization of South American Trypanosomatidae: I. The importance of lectins, neuraminic acid and neuraminidase for the differentiation of trypanosomes and leishmania from the New World. *Zool. Anz.* **223**: 67–81.
- Smith, H.E. 1973. Syngamus trachea: a first record in Hawaii. *Elepaio* **33**(9): 95–96.
- Solovjeva, G.F. & S.E. Pozdnyakov. 1984. [A new nematode species, *Hysterothylacium hospitum* n. sp. (Nematoda, Anisakidae), found in the spiral valve of a blue shark.] *Parazitologia* **18**(1): 66–68. [In Russian with English summary.]
- Sonneborn, T.M. 1975. The *Paramecium aurelia* complex of fourteen sibling species. *Trans. Am. Microscop. Soc.* **94**: 155–78.
- Speare, R. 1990. A review of the diseases of the cane toad, *Bufo marinus*, with comments

- on biological control. *Aust. Wildl. Res.* **17**: 387–410.
- Starmer, W.T.** 1981. A comparison of *Drosophila* habitats according to the physiological attributes of the associated yeast communities. *Evolution* **35**: 38–52.
- , **M.-A. Lachance, H.J. Phaff & W.B. Heed.** 1990. The biogeography of yeasts associated with decaying cactus tissue in North America, the Caribbean, and northern Venezuela. *Evol. Biol.* **24**: 253–96.
- Tamashiro, M.** 1968. Microbial control in Hawaii. *Proc. Hawaii. Entomol. Soc.* **20**: 157–67.
- Tanabe, A.M. & M. Tamashiro.** 1967. The biology and pathogenicity of a microsporidian (*Nosema trichoplusiae* sp. n.) of the cabbage looper, *Trichoplusia ni* (Hübner) (Lepidoptera: Noctuidae). *J. Invert. Pathol.* **9**: 188–95.
- Tanada, Y.** 1957. An annotated list of infectious diseases of insects in Hawaii. *Proc. Eighth Pac. Sci. Cong.* **3A**: 1235–1250.
- Testa, J. & M.D. Dailey.** 1977. Five new morphotypes of *Phyllobothrium delphini* (Cestoda: Tetraphyllidea), their relationship to existing morphotypes, and their zoogeography. *Bull. South. Calif. Acad. Sci.* **76**: 99–110.
- Tilden, J.E.** 1901. Collection of algae from the Hawaiian Islands. *Hawaii. Ann.* **1902**: 106–13.
- . 1910. *Minnesota algae*. Volume I. The Myxophyceae of North America and adjacent regions including Central America, Greenland, Bermuda, the West Indies and Hawaii. University of Minnesota, Minneapolis. iv + 328 p.
- Upton, S.J., A.A. Marchiondo & R.N. Williams.** 1988. New species of *Isospora* (Apicomplexa: Eimeriidae) from passeriform birds of Hawaii. *Syst. Parasitol.* **12**: 81–85.
- , **K. Hanley & T.J. Case.** 1994. *Eimeria frenatus* n.sp. and *Eimeria rochalimai* (Apicomplexa: Eimeriidae) from *Hemidactylus frenatus* (Sauria: Gekkonidae) in Hawaii, U.S.A. *Trans. Am. Microscop. Soc.* **113**: 390–94.
- Van Hoven, W., F.M.C. Gilchrist & T.C. Telfer.** 1992. A comparison of rumen ciliates in black-tailed deer from California and Hawaii. *Calif. Fish Game* **78**: 124–27.
- Van Riper, C., III.** 1991. The impact of introduced vectors and avian malaria on insular passeriform bird populations in Hawaii. *Bull. Soc. Vector Ecol.* **16**: 59–83.
- Van Winkle, M.E.** 1927. Freshwater testaceous rhizopods of Hawaii. Unpubl. M.S. thesis, University of Hawaii, Honolulu. 41 p.
- Vis, M.L., R.G. Sheath, J.A. Hambrook & K.M. Cole.** 1994. Stream macroalgae of the Hawaiian Islands: a preliminary survey. *Pac. Sci.* **48**: 175–87.
- Wallace, G.D. & J.K. Frenkel.** 1975. *Besnoitia* species (Protozoa, Sporozoa, Toxoplasmatidae): recognition of cyclic transmission by cats. *Science* **188**: 369–71.
- Williams A.B. & F.C. Dobbs.** 1995. A new genus and species of caridean shrimp (Crustacea: Decapods: Bresiliidae) from hydrothermal vents on Loihi Seamount, Hawaii. *Proc. Biol. Soc. Wash.* **108**: 228–37.
- Wilson, C.D. & G.W. Boehlert.** 1993. Population biology of *Gnathophausia longispina* (Mysidacea: Lophogastrida) from a central North Pacific seamount. *Mar. Biol.* **115**: 537–43.
- Yamaguti, S.** 1965. New digenetic trematodes from Hawaiian fishes, I. *Pac. Sci.* **19**: 458–81.
- . 1968a. Cestode parasites of Hawaiian fishes. *Pac. Sci.* **22**: 21–36.
- . 1968b. *Monogenetic trematodes of Hawaiian fishes*. University of Hawaii Press.

287 p.

———. 1970. *Digenetic trematodes of Hawaiian fishes*. Keigaku Publishing Co., Tokyo.
436 p.

Zimmerman, W.J. & B. Bergman. 1990. The *Gunnera* symbiosis: DNA restriction fragment length polymorphism and protein comparisons of *Nostoc* symbionts. *Microbial Ecol.* **19**: 291–302.