

Insect Field Collection Numbers of R.C.L. Perkins

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|------------|---------------|-------------------|-------------|--------------------|--|---|
| 1 | Oahu | Waianae mountains | Apr 1892 | 3000 | beaten from various trees | |
| 2 | Oahu | Waianae mountains | Apr 1892 | 3000 | dead, under bark; <i>Rhyncogonus</i> , 1 spec. | |
| 3 | Oahu | Waianae mountains | Apr 1892 | Above 3000 | | |
| 4 | Oahu | Waianae mountains | Apr 1892 | | fungus | |
| 5 | Oahu | Kaala mountains | 26 Mar 1892 | 1500 | under & in decaying oranges (II) | This appears to have been the second collection made by Perkins in Hawai'i (cf. collection no. 15 below) |
| 6 | Oahu | Waianae mountains | Apr 1892 | 2000–3000 | under bark & from dead wood of koa trees | |
| 7 | Oahu | Waianae mountains | Apr 1892 | | from kukui wood | |
| 8 | Oahu | Waianae mountains | Apr 1892 | | beaten from flowers & branches & from inside fruit of isolated trees | |
| 9 | Oahu | Near Honolulu | May 1892 | | forests, from hau trees | |
| 10 | Oahu | Waianae mountains | Apr 1892 | | casually taken whilst shooting | |
| 11 | Oahu | Kaala mountains | Apr 1892 | | decayed koa bark; 2 specimens (XI) | |
| 12 | Oahu | Waianae mountains | Apr 1892 | | various trees | |
| 13 | Oahu | Waianae mountains | Apr 1892 | 2000 | beaten from orange trees | |
| 14 | Oahu | Waianae mountains | Apr 1892 | 2000 | | |
| 15 | Oahu | Kaala mountains | Mar 1892 | 1500 | the <i>Scaphidium</i> from decaying trees (I) | Since the Roman numerals in the associated data of Perkins appear to be in chronological order, it is assumed that the date of collection for this field number must be on or before 26 March (cf. collection No. 5, which has Roman numeral II). |
| 16 | Oahu | Waianae mountains | Apr 1892 | | dead leaves | |
| 17 | Oahu | Waianae mountains | Apr 1892 | 2500 | a pair taken <i>in cop.</i> on unknown plant | |
| 18 | Oahu | Kaala mountains | — | | decayed wood | |
| 19 | Oahu | Waianae mountains | Apr 1892 | | the staph. from koa wood | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|--|---------------|-------------|--|---|
| 19* | Oahu | Lanihuli Ridge | Oct 1892 | | | |
| 20 | Oahu | Waianae mountains | Apr 1892 | 2000 | fungi & dead wood | |
| 21 | Oahu | Waianae mountains | Apr 1892 | 2000–3000 | beaten from dead koa branches | |
| 22 | Oahu | Waianae mountains | [Apr 1892] | [2000–3000] | beaten from dead koa branches | |
| 23 | Oahu | Waianae mountains | Apr 1892 | 2000 | from kukui tree | |
| 24 | Oahu | Waianae mountains | [Apr 1892] | 2000 | (XIV & XV) | from fungi on dead trees (see biological note XIV for no. 27) |
| 25 | Oahu | Waianae mountains | Apr 1892 | 1700 | | |
| 26 | Oahu | Waianae mountains | Apr 1892 | | kukui tree | |
| 27 | Oahu | Waianae mountains | Apr 1892 | 2000 | fungi on dead trees (XIV) | this collection was apparently made at the same time as collection no. 24 (see above) |
| 28 | Oahu | Waianae mountains | Apr 1892 | 3000 | beaten from shrub | |
| 29 | Oahu | Waianae mountains | Apr 1892 | | kukui tree | |
| 30 | Oahu | Forest near Honolulu | May 1892 | | beaten from native apple tree | |
| 30* | Oahu | Oahu mountains | winter months | 2500 | | |
| 31 | Oahu | Waianae mountains | Apr 1892 | | Carabidae under bark; the weevil (<i>Carpophilus</i>) in house, 3000 ft. | |
| 32 | Oahu | Kawailoa gulch; near head of left branch | Apr 1893 | | rocks in stream | |
| 33 | Oahu | Halemano | Jan 1893 | 2000 | from koa | [= Helemano] |
| 34 | Oahu | Kaala mountains | Dec 1892 | 2000 | in very wet decaying log of kukui, <i>Aleurites</i> | |
| 35 | Oahu | Waianae mountains | Apr 1892 | 2000 | | |
| 36 | Oahu | Waianae mountains | Apr 1892 | 2000 | | |
| 37 | Oahu | Waianae mountains | Mar 1893 | | | |
| 38 | Oahu | Konahuanui | Nov 1892 | 3000 | in stem of fern | |
| 39 | Oahu | Mountains near Honolulu | 27 Dec 1892 | 2000 | | |
| 40 | Oahu | Kaala mountains | Dec 1892 | 2000 | anobiids | |
| 41 | Oahu | Kawailoa gulch | Mar, Apr 1893 | | all from the gulch itself & very far up; about 56 spec. | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|---------------------------------------|----------------|-------------|---|---|
| 42 | Oahu | Lanihuli ridge, near Honolulu | 27 Oct 1892 | nearly 3000 | from leaves of trees (pair of <i>Disenochus</i>) | |
| 43 | Oahu | Mountains near Honolulu | Oct 1892 | 2500 | | [probably Nu'uaniu area] |
| 44 | Oahu | Ridges round Nuuanu Valley | Nov 1892 | | | |
| 45 | Oahu | Koolau range, far back of Waialua | Jan 1893 | 2000 | under stones in wet gulches | |
| 46 | Oahu | Lanihuli ridge | Oct 1892 | 2000 | in stems of various flag-like plants growing in tufts | [after 24 October 1892] |
| 47 | Oahu | Mountains near Honolulu | 27 Oct 1892 | 2000 | under bark | [Waiolani Ridge side of the Nu'uaniu Valley up to Lanihuli] |
| 48 | Oahu | Ridges round Nuuanu Valley | Nov 1892 | 2000 & up | | [before 6 November 1892] |
| 49 | Oahu | Head of Kawailoa gulch | Apr 1893 | | | |
| 50 | Oahu | Koolau range, mountains above Waialua | Mar 1893 | | beaten | |
| 51 | Oahu | Kaala mountains | Dec 1892 | 3000 | | |
| 52 | Oahu | Kaala mountains | Dec 1892 | 3000 | note/ extraordinarily hairy | |
| 53 | Oahu | Kaala mountains | Dec 1892 | 2500 | | |
| 54 | Oahu | Kawailoa | [Feb-Apr 1893] | | base of leaves of <i>Freycinetia</i> | |
| 55 | Oahu | Koolau range behind Waialua | Mar 1893 | 2000 | under stone in wet gulch | |
| 56 | Oahu | Kaala mountains | Jan 1893 | 2000 | | |
| 57 | Oahu | Kaala mountains | Dec 1892 | 3000 | | |
| 58 | Oahu | Kawailoa gulch | Feb 1893 | 2000 | | |
| 59 | Oahu | Mt. Kaala | Jan 1893 | | | |
| 60 | Oahu | Mt. Kaala | Dec 1892 | 2000 | | |
| 61 | Oahu | Mountains behind Honolulu | Nov 1892 | 2000 & up | | [Nu'uaniu Valley; before 6 November 1892] |
| 62 | Oahu | Kaala mountains | Dec 1892 | 2000 | from decaying kukui nuts | |
| 63 | Oahu | Mountains near Honolulu | Oct, Nov 1892 | above 2000 | | [Nu'uaniu Valley; between 24 Oct and 6 November 1892] |
| 64 | Oahu | Mt. Kaala | Dec 1892 | | high up mountain in gulches | |
| 65 | Oahu | Kawailoa gulch | Apr 1893 | | far up gulch | |
| 66 | Oahu | Kawailoa gulch | Apr 1893 | 2000 | under wet log | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|---|---------------------|-------------|---|---|
| 67 | Oahu | Mountains near Honolulu | Oct, Nov 1892 | 2500–3000 | <i>Heterhamphus</i> in lilies; carabids lillie & ferns | [Nu‘uanu Valley; between 24 Oct and 6 November 1892] |
| 68 | Oahu | Ridge near head of Kawaihoa gulch | Apr 1893 | | beaten from tree | |
| 69 | Oahu | Kaala mountains | Dec 1892 | | | |
| 70 | Oahu | Mountains near Honolulu | [Oct, Nov. 1892] | 2000 | from lillies | |
| 71 | Oahu | Mountains above Waialua | Mar 1893 | 2000 | | [probably the Ko‘olau range, not Waianaes] |
| 72 | Oahu | Mountains near Honolulu | | | at base of leaves of <i>Freycinetia</i> | |
| 73 | Oahu | Kaala mountains | Dec 1892 | 3000 | | |
| 74 | Oahu | Mountains behind Waialua | Feb [1893] | 1000 | | [originally transcribed in the Cambridge list as “II ‘92” which is impossible since Perkins did not arrive in Hawaii until March 1892; since this locality is where he was in 1893, it is assumed here that the date is correctly “II ‘93”] |
| 75 | Oahu | Kaala mountains | Dec 1892 | 3000 | | |
| 76 | Oahu | Mountains near Honolulu | 24 Oct 1892 | 2000+ | under bark with larvae preserved in spirit marked "B" | [Nu‘uanu Valley; after 24 October] |
| 77 | Oahu | Mountains behind Waialua, Kawaihoa gulch | Apr 1893 | 2000 | | |
| 78 | Lanai | behind Koele | Feb 1894 | | | [before 21 February 1894] |
| 79 | Lanai | Lanai gulch | Jan, Feb 1894 | about 2000 | carabid, 2 black legged under bark | [before 23 February 1894] |
| 80 | Lanai | Lanai gulch | Jan 1894 | 2000 | | |
| 81 | Oahu | Kaala | May, Jun 1894 | | | |
| 82 | Lanai | behind Koele | Feb 1894 | | | [before 23 February 1894] |
| 83 | Lanai | — | 1894 | about 2000 | | [before 23 February 1894] |
| 84 | Lanai | — | Dec 1893 | about 2000 | | [after 11 December 1893] |
| 85 | Lanai | — | Oct 1893 | sea level | | [probably an error in transcription for December; Perkins did not arrive on Lāna‘i until 11 December 1893] |
| 86 | Lanai | — | Jan 1894 | above 2000 | | |
| 87 | Lanai | behind Koele | Feb 1894 | | | [before 21 February 1894] |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|---------|--------------------------------|---------------|---------------------|---|---|
| 88 | Lanai | behind Koele | Jan 1894 | above 2000 | | |
| 89 | Lanai | near Koele | Jan 1894 | above 2000 | | |
| 90 | Lanai | windward side, looking to Maui | Feb 1894 | 2000 | all from pua trees (except <i>Oodemias</i>) chiefly anobiids | [before 21 February 1894] |
| 91 | Lanai | — | Jan 1894 | about 2000 | | |
| 92 | Lanai | — | Dec 1893 | about 2000 | | [after 11 December 1893] |
| 93 | Lanai | — | Jan 1894 | 3000 | | |
| 94 | Lanai | near Koele | Jan 1894 | | | |
| 95 | Molokai | Pelekunu | Oct 1893 | | | [after 9 October 1893] |
| 96 | Lanai | above Waipaa | Feb 1894 | about 3000 | | [before 21 February 1894] |
| 97 | Molokai | leeward side | Oct 1893 | sea level or nearly | | |
| 98 | Lanai | leeward | Jan, Feb 1894 | about 2000 | anobiids all from pua trees | [before 23 February 1894] |
| 99 | Lanai | near Waipaa | Feb 1894 | above 2000 | | [before 23 February 1894] |
| 100 | Lanai | near Waipaa | Feb 1894 | | | [before 23 February 1894] |
| 101 | Lanai | above Waipaa | Feb 1894 | about 3000 | Heteroptera | [before 23 February 1894] |
| 102 | Lanai | above Waipaa, looking to Maui | Feb 1894 | about 3000 | <i>Cis</i> & parasitic Hymenoptera | [before 23 February 1894] |
| 103 | Lanai | windward side | Feb 1894 | | anobiids in cop. off pua trees | [before 23 February 1894] |
| 104 | Lanai | near Waipaa | [Feb 1894] | about 2000 | | [before 23 February 1894] |
| 105 | Lanai | near Waipaa | [Feb 1894] | about 2000 | off pua trees, 2 very minute staphs. | [before 23 February 1894] |
| 106 | Molokai | Pelekunu | Oct 1893 | | | [after 9 October 1893] |
| 107 | Lanai | on main ridge | Feb 1894 | 3000 | | [before 23 February 1894] |
| 108 | Molokai | [Pelekunu] | Oct 1893 | 3000 | | [after 9 October 1893] |
| 109 | Lanai | — | — | [sea level] | ants & beetles, coast species under logs | [between 11 December 1893 and 23 February 1894] |
| 110 | Maui | Haleakala | 5 Apr 1894 | 5000 | the brown largest <i>Oodemias</i> from akala stem | |
| 111 | Maui | Haleakala | Apr 1894 | 5000 | | |
| 112 | Maui | Haleakala, Olinda | 31 Mar 1894 | 4000 | | |
| 113 | Maui | Haleakala | Apr 1894 | 5000 | by grubbing | |
| 114 | Maui | Haleakala | 20 Mar 1894 | 5000 | | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|------|--------|-------------------------------|---------------|---------------|--|-----------------------------|
| 115 | Maui | Haleakala | Mar 1894 | 5000 | the larger sp (<i>Cyclothorax</i>) in moss & on leaves of growing trees | |
| 115 | Maui | Haleakala | Mar 1894 | 4500 | the smaller (gen.?) [than above] under stones | |
| 115 | Maui | Haleakala | Mar 1894 | 5000 | single brown specimen | |
| 116 | Maui | Haleakala | Apr 1894 | 5000 | all under bark | |
| 117 | Maui | Haleakala | Apr 1894 | 4500 | | |
| 118 | Maui | [Wailuku] | [1894] | | top layer Wailuku mutilated odyneroid | [April or May] |
| 118 | Maui | Iao Valley | Apr, May 1894 | | bottom layer IV & V '94 | |
| 119 | Oahu | Pauoa | — | | | [possibly Pauoa flats area] |
| 120 | Maui | Haleakala | 6 Apr 1894 | 5000 | | |
| 121 | Maui | Haleakala, woods below Olinda | 18 Mar 1894 | above 5000 | | |
| 122 | Maui | Haleakala | Apr 1894 | | | |
| 123 | Maui | Haleakala | Mar 1894 | 5000 | | |
| 124 | Maui | Haleakala | 11 Apr 1894 | 9000 | 1 <i>Proterhinus</i> | |
| 124 | Maui | Haleakala | 11 Apr 1894 | 9000 | 3 thrips | |
| 124 | Maui | Haleakala | 11 Apr 1894 | nearly 10,000 | 2 staphs | |
| 125 | Maui | Haleakala | 5–6 Apr 1894 | 5000 | | |
| 126 | Maui | Haleakala | 9 Apr 1894 | 9000–10,000 | | |
| 127a | Maui | Haleakala | 15 Mar 1894 | 5000 | | |
| 127b | Maui | Haleakala | 16 Mar 1894 | 5000 | | |
| 127c | Maui | Haleakala | 18 Mar 1894 | 5000 | | |
| 128 | Maui | Haleakala | [1894] | above 5000 | | |
| 129 | Maui | Haleakala | 10 Apr 1894 | 5000 | top layer <i>Oodemus</i> (2 species) & <i>Proterhinus</i> from Akala stems | |
| 130 | Maui | Haleakala | Apr 1894 | 8000 | Diptera | |
| 130 | Maui | Haleakala | Apr 1894 | 9500 | carabid | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|---------|------------------------|----------------|--------------|------------------------------------|--|
| 130 | Mau | Haleakala | Apr 1894 | 9000 | the rest [of above] | |
| 131 | Lanai | above Koele, Mahana? | July 1894 | 3000 | | [before 13 July 1894] |
| 132 | Lanai | Halepaakai | July 1894 | | | [before 13 July 1894] |
| 133 | Lanai | Lanaihale | July 1894 | | | [before 13 July 1894] |
| 134 | Lanai | Halepaakai | July 1894 | | | [before 13 July 1894] |
| 135 | Lanai | Lanai summit | July 1894 | | | [before 13 July 1894] |
| 136 | Lanai | Halepaakai | July 1894 | | bugs | [before 13 July 1894] |
| 137 | Lanai | Halepaakai | Jan 1894 | | <i>Straussia</i> | |
| 138 | Hawaii | Kona | August 1894 | 3000 | | |
| 139 | Hawaii | Kilauea | July 1894 | | | [after 22 July 1894 arrival date] |
| 140 | Hawaii | Kona | Aug 1894 | 3000 | | [before 15 August 1894] |
| 141 | Molokai | — | 28 Aug 1893 | 4500 upwards | | [ridge above Waikolu Valley] |
| 142 | Hawaii | Kona | Aug 1894 | | | [before 15 August 1894] |
| 143 | Kauai | — | May 1894 | 4000 | | |
| 144 | Hawaii | Kilauea | July 1894 | 5000 | | [after 22 July 1894 arrival date] |
| 145 | Molokai | forest above Pelekunu | Sep 1893 | | | |
| 146 | Molokai | Kahanui | 18 Sep 1893 | | | |
| 147 | Molokai | — | 22 Sep 1893 | 5000 | | [forest near ridge above Waikolu Valley] |
| 148 | Molokai | above head of Pelekunu | Sep 1893 | 4500+ | | |
| 149 | Molokai | above Kalawao | 1 Aug 1893 | | | |
| 150 | Molokai | above Kalawao | 12 July 1893 | 4000 | | |
| 151 | Molokai | gulch | 5 July 1893 | 3500 or more | | [Makakupaia area] |
| 152 | Molokai | — | 19–20 Sep 1893 | 3000 | | |
| 153 | Molokai | Kahanui | 11 Aug 1893 | | | [either the locality or the date is wrong. Perkins was above Kalawao on this date; nowhere near the coast] |
| 154 | Kauai | above Waimea | May 1894 | 5000 | 6 <i>Apterocyclus</i> 3 found dead | [above Halemanu area] |
| 156 | Molokai | — | 6 Sep 1893 | 3000 | | [Makakupaia area] |
| 157 | Molokai | — | 9 Sep 1893 | 4500 | | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|------|---------|------------------|----------------|----------------|-----------------------------------|--------------------|
| 158 | Molokai | highest forest | 13 Sep 1893 | | | |
| 159 | Molokai | — | 11 Sep 1893 | 4500 & upwards | | |
| 160 | Molokai | Kalae | Aug 1893 | | | [1–11 August 1893] |
| 161 | Molokai | — | 23 Sep 1893 | 4500 | | |
| 162 | Molokai | highest forest | 17 Sep 1893 | | | |
| 163 | Molokai | — | 11 Sep 1893 | 4500 & upwards | ? same as 159 | |
| 164 | Molokai | — | 19–20 Sep 1893 | 3000 above | swept | |
| 165 | Molokai | head of Pelekunu | 23 Aug 1893 | | | |
| 166 | Molokai | above Kalawao | 8 Aug 1893 | | | |
| 167 | Molokai | Kalae | 7 Aug 1893 | | | |
| 168 | Molokai | Makakupaia | 26 Jul 1893 | | | |
| 169 | Molokai | — | 9–11 Aug 1893 | | | |
| 169* | Molokai | Kalaana | 12 May 1893 | 3000 above | fly parasitic on <i>Himatione</i> | |
| 170 | Molokai | Kaunakakai | Jul 1893 | sea level | | |
| 171 | Molokai | — | 8 Sep 1893 | 3000 | | |
| 172 | Molokai | Kalae | 7 Aug 1893 | | | |
| 173 | Molokai | above Pelekunu | Jul 1893 | | | |
| 174 | Molokai | — | 24 Jun 1893 | 3000 | | |
| 175 | Molokai | — | 22 May 1893 | | | |
| 176 | Molokai | — | 12 Jun 1893 | 3000 | | |
| 177 | Molokai | — | 5 Jun 1893 | above 3000 | | [Makakupaia area] |
| 178 | Molokai | — | 17 Jul 1893 | | | |
| 179 | Molokai | — | 9 Jun 1893 | | | [Makakupaia area] |
| 180 | Molokai | — | 21 Jun 1893 | 4500 | | |
| 181 | Molokai | — | 13 Jun 1893 | 4500 | | |
| 182 | Molokai | — | 18 Jun 1893 | 4500 | | |
| 183 | Molokai | — | May 1893 | 3500–4500 | Metromeni under stones | [Makakupaia area] |
| 184 | Molokai | — | 17 May 1893 | about 4000 | | |
| 185 | Molokai | — | 4 Jun 1893 | 3000 | | [Makakupaia area] |
| 186 | Molokai | — | 27 Jun 1893 | 5000 | all in flowers of lillies | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|---------|------------------|-------------|-------------|---|-------------------|
| 187 | Kauai | Waimea mountains | May 1894 | about 4000 | | |
| 188 | Molokai | — | 10 Jun 1893 | 4000 | | |
| 189 | Molokai | — | 29 May 1893 | 3000–4000 | | |
| 190 | Molokai | — | 27 May 1893 | 4000 | | |
| 191 | Molokai | — | 27 May 1893 | 4000 | | |
| 192 | Molokai | — | 22 Jun 1893 | 6000 | | |
| 193 | Molokai | — | 9 Jun 1893 | | Hemiptera | [Makakupaia area] |
| 194 | Hawaii | Kona | Sep 1892 | about 2500 | | |
| 195 | Hawaii | Kona | Jun 1892 | 3000 | beaten from koa | |
| 196 | Molokai | — | 11 Jun 1893 | | and other dates | |
| 197 | Molokai | — | 1893 | | odd specimens | |
| 198 | Hawaii | Kona | Sep 1892 | 3000 | | |
| 199 | Hawaii | Kona | Sep 1892 | 2500 | Staphs from <i>Freycinetia</i> | |
| 200 | Hawaii | Kona | Sep 1892 | 3000 | | |
| 201 | Hawaii | Kona | Oct 1892 | 3000 | | |
| 202 | Hawaii | Kona | Aug 1892 | 4000 | | |
| 203 | Hawaii | Kona | Sep 1892 | 3000 | cossonid from tree fern | |
| 204 | Hawaii | Kona | Sep 1892 | 5000 | note that the small homopteron, sometimes green, sometimes yellow is exactly like the ripe & unripe grass seeds enclosed, on which it rests | |
| 205 | Oahu | Honolulu | | | in house | |
| 206 | Hawaii | Kona | Sep 1892 | 2000+ | | |
| 207 | Hawaii | Kona | Sep 1892 | 3000+ | under bark of dead tree | |
| 208 | Hawaii | Kona | Sep 1892 | 2500 | from kukui trees | |
| 209 | Hawaii | Kona | 24 Jul 1892 | 4000 | | |
| 210 | Hawaii | Kona | Sep 1892 | 3000 | this extremely active runner beaten from dead <i>Acacia</i> | |
| 211 | Hawaii | Kona | Sep 1892 | 1500 | swept up, Hilo grass | |
| 212 | Hawaii | Kona | Sep 1892 | 3000 | | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|----------|-------------|-------------|--|--|
| 213 | Hawaii | Kona | 20 Jun 1892 | 2500 | | [Perkins's journal entry for this date says 4000 feet. He might have also collected at lower elevation locations on his way down to Ka'awaloa] |
| 214 | Hawaii | Kona | Sep 1892 | 3000 | <i>Cis</i> from fungus on <i>Acacia</i> | |
| 215 | Hawaii | Kona | Jul 1892 | 4000 | pair of Nitidulidae & a pair of cossonids in cop | |
| 216 | Hawaii | Kona | Sep 1892 | 2000 | 5 spec.; 1 very minute in very wet decayed log | |
| 217 | Hawaii | Kona | | about 3000 | beaten from various trees, the red <i>Proterhinus</i> & the sp. of Cossonidae from tree fern | |
| 218 | Hawaii | Kaawaloa | Sep 1892 | 1500 | flying round house at dusk | [no doubt near the Kalukalu area] |
| 219 | Hawaii | Kona | Sep 1892 | 3000 | | |
| 220 | Hawaii | Kona | Sep 1892 | 5000 | this fine sp. was taken from an old stunted koa growing alone in the midst of a belt of rough lava | |
| 221 | Hawaii | Kona | Sep 1892 | 2000 | | |
| 222 | Hawaii | Kona | Sep 1892 | 2500-3000 | | |
| 223 | Hawaii | Kona | Sep 1892 | 4000 | from decayed koa | |
| 224 | Hawaii | Kona | Jul 1892 | 4000 | No. 26, 2 males, 1 female | |
| 225 | Hawaii | Kona | Sep 1892 | 4000 above | Heteroptera & Homoptera both from bastard sandal tree | |
| 226 | Hawaii | Kona | Sep 1892 | 5000 | | |
| 227 | Hawaii | Kona | Jun 1892 | 3000 | outside burrow in trunk of a dead tree | [after 20 June 1892] |
| 228 | Hawaii | Kona | Sep 1892 | 2500-3000 | <i>Cis</i> beaten from forest trees other than <i>Acacia</i> , on climbers | |
| 229 | Hawaii | Kona | Jun 1892 | 2500 | Staphs. nitidulids from <i>Freycinetia</i> | [after 20 June 1892] |
| 230 | Hawaii | Kona | Sep 1892 | 3000 | from various trees | |
| 231 | Hawaii | Kona | Aug 1892 | 4000 above | | |
| 232 | Hawaii | Kona | Sep 1892 | 4000+ | from decayed koa | |
| 233 | Oahu | Halemano | Feb 1893 | 2000 | from koa | [= Helemano] |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|---------|---|-------------|-------------|--|----------------------|
| 234 | Molokai | Waikolu | 24 Sep 1893 | | | |
| 235 | Hawaii | Kona | Sep 1892 | 3000 | | |
| 236 | Oahu | near Honolulu | Oct 1892 | about 2000 | beaten from koa | |
| 237 | Oahu | Nuuanu Valley | Oct 1892 | | | |
| 238 | Oahu | Lanihuli ridge | Oct 1892 | | beaten | |
| 239 | Oahu | near head of the south branch of the Kawaihoa gulch | Apr 1893 | | | |
| 240 | Oahu | Halemano | Dec 1892 | 2000+ | | [= Helemano] |
| 241 | Oahu | Halemano | Jan 1893 | 2000+ | | [= Helemano] |
| 242 | Oahu | ridge of Nuuanu Valley, Konahuanui side | May 1892 | | beaten from dead branches of <i>Acacia koa</i> ; <i>Proterhinus</i> , <i>Holcobius</i> , <i>Cis</i> & c | |
| 243 | Hawaii | Kona | Jun 1892 | 3000 | beaten from trunks & branches of various trees | |
| 244 | Oahu | mountains near Honolulu | 27 Oct 1892 | 2000 | <i>Cis</i> beaten | |
| 245 | Hawaii | Kona | 27 Jul 1892 | 3500 | | [Pulehua area] |
| 246 | Hawaii | Kona | Jun 1892 | about 3000 | beaten from kukui | [after 20 June 1892] |
| 247 | Hawaii | Kona | 26 Jul 1892 | 4000 | | [Pulehua area] |
| 248 | Hawaii | Kona | Sep 1892 | 2000–3000 | beaten from trees | |
| 249 | Hawaii | Kona | Jul 1892 | | beaten from <i>Acacia</i> trees | |
| 250 | Maui | Haleakala | Apr 1894 | about 5000 | under bark of <i>Acacia</i> | |
| 251 | Maui | Haleakala | | | all the small carabids & Homoptera by grubbing; the <i>Metromenus</i> from lilly, 5000 ft; Apr 1894; one unique carabid with light elytra black spotted taken by grubbing 8 Apr 1894 | |
| 252 | Maui | Haleakala, Olinda | Apr 1894 | 3000–4000 | 60 <i>Metromeni</i> & 20 of the large carabid. the latter on dead branches of trees & under bark of <i>Acacia</i> | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|--------------------------|----------|-------------|--|------------------------------|
| 253 | Kauai | — | Jun 1894 | 4000 | | [Halemanu area] |
| 254 | Mauai | Haleakala | Apr 1894 | 9500–10,000 | | |
| 255 | Kauai | mountains above Waimea | May 1894 | 4000 | | [Halemanu area] |
| 256 | Kauai | mountains above Waimea | May 1894 | 4000 | large <i>Brachypeplus</i> under bark of same tree as Munro's cossonid. Other <i>Brachypeplus</i> from various trees & flowers of "haha". minute carabids by grubbing | [Halemanu area] |
| 257 | Kauai | mountains above Waimea | May 1894 | about 4000 | anobiids & staphs. 1 very dull dark <i>Brachypeplus</i> under koa bark, 1 browner species from flower of lilly | [Halemanu area] |
| 258 | Kauai | mountains above Waimea | May 1894 | about 4000 | | [Halemanu area] |
| 259 | Kauai | mountains above Waimea | May 1894 | about 4000 | chiefly <i>Proterhinus</i> | [Halemanu area] |
| 260 | Kauai | mountains above Makaweli | Jun 1894 | 2000–3000 | <i>Clytarlus</i> (2), <i>Apterocyclus</i> (1), Otorhynchid (several) & 1 <i>Itodacnus</i> | |
| 260 | Kauai | Hanapepe | Jun 1894 | | <i>Aegosoma</i> (1) | |
| 261 | Kauai | mountains above Waimea | May 1894 | about 4000 | 15 Munro's cossonids | [Halemanu area] |
| 262 | Kauai | mountains above Waimea | May 1894 | about 4000 | 100 carabids | [Halemanu area] |
| 263 | Kauai | mountains above Makaweli | May 1894 | 2000+ | | |
| 264 | Kauai | mountains above Waimea | May 1894 | 3000+ | chiefly anobiids | |
| 265 | Kauai | mountains above Waimea | May 1894 | 4000 | | [Halemanu area] |
| 266 | Kauai | — | Jun 1894 | about 4000 | carabids. <i>Clytarlus</i> (1), <i>Brachypeplus</i> (1), <i>Parandra</i> , large staphs (5), <i>Anotheorus</i> | [probably the Halemanu area] |
| 267 | Kauai | — | Jun 1894 | 3000 | 2 or 3 rare, solid black carabids beaten from dead sticks &c; also the <i>Eopenthes</i> | |
| 267 | Kauai | — | Jun 1894 | 4000 | | [Halemanu area] |
| 268 | Kauai | mountains above Waimea | Jun 1894 | 3000 | | |
| 269 | Hawaii | Kona | 1892 | | overlooked specimens | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|--------------------------|-------------|-------------|---|----------------------|
| 270 | Hawaii | Kona | Jun 1892 | 3000+ | remains of longicorn, very abundant, probably wrong season as I saw no live spec. | [after 20 June 1892] |
| 271 | Kauai | mountains above Waimea | May 1894 | 4000 | about 45 carabids | [Halemanu area] |
| 272 | Oahu | Waialua, coast | March 1892 | | from underground nest (ants) (VII) | |
| 273 | Kauai | mountains above Waimea | May 1894 | | above: 3 carabids far up Waimea River under stones; below: minute <i>Cis</i> , Staphs, &c | |
| 274 | Maui | Haleakala, Olinda woods | Apr 1894 | 3000–4000 | 139 anobiids & a few other things. of the two large sp. separated at bottom of tube, the black punctured one is under bark of smooth barked <i>Acacia</i> ; the pubescent sp. (?genus) from kolea | |
| 275 | Hawaii | Kona | Jun 1892 | 2000–3000 | | [after 20 June 1892] |
| 276 | Hawaii | Kona | Jun 1892 | about 3000 | beaten from Ohia | [after 20 June 1892] |
| 277 | Hawaii | Kona | Aug 1892 | about 4000 | elater & Cossonidae | [Pulehua area] |
| 278 | Hawaii | Kona | Jun 1892 | 2000–3000 | | [after 20 June 1892] |
| 279 | Hawaii | Kona | 29 Jun 1892 | 4000 | staph. from fungus; others from dead wood | |
| 280 | Kauai | mountains above Makaweli | Jun 1894 | above 2000 | | |
| 281 | Kauai | mountains above Makaweli | Jun 1894 | above 2000 | | |
| 282 | Oahu | Kawailoa | Apr 1893 | | fern stems | |
| 283 | Hawaii | Kona | Sep 1892 | 2500 | from decaying lehua tree | |
| 284 | Hawaii | Kona | 25 Jul 1892 | about 4000 | chiefly Rhynchota | [Pulehua area] |
| 285 | Kauai | — | Jun 1894 | about 4000 | Staphs | [Halemanu area] |
| 286 | Hawaii | Kona | Jun 1892 | 3000 | waterbugs | [after 20 June 1892] |
| 287 | Hawaii | Kona | 19 Aug 1892 | about 4000 | Coleoptera & Hymenoptera | [Pulehua area] |
| 288 | Hawaii | Kona | 14 Jul 1892 | 4000 | one small sp. (single spec.) | [Pulehua area] |
| 289 | Hawaii | Kona | | 4000 | No. 21 4 spec. excessively minute under paper | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|-------------------------------|-------------|-------------|---|--|
| 290 | Hawaii | Kona | 1 Jul 1892 | | from dead wood, 1 very minute species | |
| 291 | Oahu | right [side] of Nuuanu Valley | May 1892 | | beaten from dead leaves of a <i>Pandanus</i> | |
| 292 | Hawaii | Kona | Aug 1892 | 4000 | from fungi on koa tree, 5 spec. very minute | [Pulehua area] |
| 293 | Hawaii | Hualalai | 5 Aug 1892 | about 5000 | Rhynchota &c | |
| 294 | Hawaii | Kona | Jun 1892 | 4000+ | | [after 20 June 1892] |
| 295 | Hawaii | Kona | Jun 1892 | 4500 | from dead wood | [after 20 June 1892] |
| 296 | Hawaii | Kona | Jun 1892 | about 3000 | | [after 20 June 1892] |
| 297 | Hawaii | Kona | 4 Jul 1892 | about 5000 | | |
| 298 | Hawaii | Kona | 21 Jul 1892 | 3500 | | |
| 299 | Hawaii | Kona | Sep 1892 | 4000+ | | |
| 300 | Hawaii | Kona | 4 Jul 1892 | about 5000 | | |
| 301 | Hawaii | Kona | 1 Jul 1892 | about 4000 | on fungi | |
| 302 | Hawaii | Kona | Sep 1892 | 4000+ | | |
| 303 | Hawaii | Kona | 29 Jun 1892 | about 4000 | | [Pulehua area] |
| 304 | Hawaii | Kona | 9 Aug 1892 | about 4000 | | |
| 305 | Hawaii | Kona | 30 Jun 1892 | about 5000 | from dead wood of several trees | [Pulehua area] |
| 306 | Hawaii | Kona | 29 Aug 1892 | 4000 | No. 27 | |
| 307 | Hawaii | Kona | 10 Aug 1892 | 4000 | | |
| 308 | Hawaii | Kona | 1 Jul 1892 | about 4000 | on dead trunks | |
| 309 | Hawaii | Kona | Sep 1892 | about 2000 | all from flowers of <i>Freycinetia</i> | |
| 310 | Hawaii | Mt. Hualalai | 5 Aug 1892 | about 5000 | 5 very minute staphs | [in saddle between Hualalai and Mauna Loa] |
| 311 | Hawaii | Kona | Jun 1892 | 4500 | beaten from the "maile" plant | |
| 312 | — | — | — | — | found in bottom of box containing mostly Kona things | [probably 1892] |
| 313 | Hawaii | Kona | Sep 1892 | about 2000 | find minute beetles (? <i>Cis</i>) & small Hymenoptera | |
| 314 | Hawaii | Kona | 25 Jul 1892 | about 4000 | | |
| 315 | Hawaii | Kona | 30 Jun 1892 | nearly 5000 | | [Pulehua area] |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|--------------|-------------|--------------|---|---|
| 316 | Hawaii | Kona | 9 Aug 1892 | about 4000 | | |
| 317 | Hawaii | Kona | Jul 1892 | about 4000 | odd specimens | |
| 318 | Hawaii | Kona | 4 Jul 1892 | about 5000 | | |
| 319 | Hawaii | Kona | Jul 1892 | about 4000 | | |
| 320 | Hawaii | Kona | 30 Jun 1892 | nearly 5000 | from flowers & dead wood | [in forest above Pulehua area] |
| 321 | Hawaii | Kona | 30 Jun 1892 | nearly 5000 | large sp. from spider's web; pair taken in cop, two very small staphs | |
| 322 | Hawaii | Kona | 4 Jul 1892 | about 5000 | | [original transcription has "IV 7.92" which probably is a misinterpretation of "4 Jul 1892" since Perkins did not arrive in Kona until June 1892] |
| 323 | Hawaii | Kona | 1 Aug 1892 | 6000 | Hymenoptera & Hemiptera | |
| 324 | Hawaii | Kealakekua | 26 Jun 1892 | on the beach | | |
| 325 | Hawaii | Kona | 11 Aug 1892 | 4000 | | |
| 326 | Hawaii | Kona | 18 Jul 1892 | 30004000 | | |
| 327 | Hawaii | Kona | 16 Jul 1892 | about 4000 | | |
| 328 | Hawaii | Kona | 1 Jul 1892 | about 4000 | from fungi | |
| 329 | Hawaii | Kona | 9 Aug 1892 | about 4000 | No. 25 | |
| 330 | Hawaii | Kona | Jun 1892 | 4500 | from fungi & dead wood, manu specimens small | [after 20 June 1892] |
| 331 | Hawaii | Kona | 30 Jun 1892 | nearly 5000 | from dead branches & trunks | [Pulehua area] |
| 332 | Hawaii | Mt. Hualalai | 5 Aug 1892 | about 5000 | | |
| 333 | Hawaii | Kona | Aug 1892 | about 4000 | | |
| 334 | Hawaii | Kona | 21 Jul 1892 | 3500 | | |
| 335 | Hawaii | Kona | 21 Jul 1892 | 3500 | | |
| 336 | Hawaii | Kona | Jun 1892 | 4000 | | [after 20 June 1892] |
| 337 | Hawaii | [Kona] | 5 Aug 1892 | about 5000 | | |
| 338 | Hawaii | Kona | 1 Jul 1892 | 4000 | under bark | |
| 339 | Hawaii | Kona | 1 Aug 1892 | 4000+ | | |
| 340 | Hawaii | Kona | 30 Jun 1892 | nearly 5000 | dry dead wood of various trees | [Pulehua area] |
| 341 | Hawaii | Kona | 25 Jul 1892 | | | [Pulehua area] |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|---------|----------------------------------|---------------|-----------------------|---|---|
| 342 | Hawaii | Kona | 1 Aug 1892 | | Hemiptera &c | [Pulehua to Kanahaha] |
| 343 | Hawaii | Kona | 20 Jul 1892 | | | [Pulehua area] |
| 344 | Hawaii | Mt. Hualalai | 5 Aug 1892 | about 5000 | | [saddle area between Hualalai and Mauna Loa; Perkins's journal entry for this date says he collected up to 6000 ft; it is unlikely he summited the peak, which is 8000 ft.] |
| 345 | Molokai | — | 11 Jun [1893] | | upper layer; all exceedingly minute | [Makakupaia area] |
| 345 | Molokai | — | 12 Jun [1893] | | lower layer | [Makakupaia area] |
| 346 | Maui | Haleakala | 11 Apr 1894 | | "Mauna" | |
| 347 | Maui | Haleakala | 9 Apr 1894 | slightly above 10,000 | top layer: 20 <i>Cyclothorax</i> unique? very shiny, nearly punctured (2nd largest species) brown black | |
| 347 | Maui | Haleakala | 9 Apr 1894 | nearly 10,000 | the rest [= not top layer] | |
| 348 | Maui | Haleakala | March 1894 | 5000 | | |
| 349 | Maui | Iao Valley | Mar 1894 | | | |
| 350 | Maui | Haleakala | 28 Mar 1894 | 4500 | | |
| 351 | Maui | Haleakala (Olinda) | 1894 | 4000 | fragments of beetles | |
| 352 | Maui | Haleakala | Apr 1894 | 5000 | <i>Brachypeplus</i> under <i>Acacia</i> bark | |
| 353 | Maui | Haleakala | 9 Apr 1894 | about 9500 | 39 spec. of <i>Cyclothorax</i> | |
| 354 | Maui | Haleakala | 1 Apr 1894 | 5000+ | | |
| 355 | Maui | Haleakala | Mar 1894 | 5000+ | | |
| 356 | Maui | Haleakala | Apr 1894 | 5000 | | |
| 357 | Maui | Haleakala | Apr 1894 | 4000 | coleops | |
| 358 | Maui | Haleakala | Mar 1894 | 5000 | | |
| 359 | Maui | Haleakala | Mar 1894 | 5000 | | |
| 360 | Maui | Iao Valley | Apr, May 1894 | | | |
| 361 | Maui | Haleakala, from Olinda downwards | Apr 1894 | 4000–3000 | the <i>Proterhinus</i> from "maile" stem; the large <i>Oodemus</i> & the flat <i>Brachypeplus</i> under koa bark; the <i>Fornax</i> under log | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|-------------------------|---------------|------------------|---|-----------------|
| 362 | Maui | Haleakala (Olinda) | Apr 1894 | | <i>Cyclothorax</i> under koa bark; <i>Disenochus</i> under log | |
| 363 | Maui | Haleakala | 28 Mar 1894 | 4500 | | |
| 364 | Maui | Haleakala (Olinda) | 31 Mar 1894 | 4000 | find 3 spms. of ponaid ant | |
| 365 | Kauai | — | Jul 1894 | about 4000 | <i>Brachypeplus</i> & anobiids | [Halemanu area] |
| 366 | Maui | Haleakala | Apr 1894 | 4500 | <i>Proterhinus</i> from “kolea” tree | |
| 367 | Maui | Haleakala | Apr 1894 | 4500 | The <i>Proterhinus</i> from smoothed bark <i>Acacia</i> | |
| 368 | Maui | West Maui | — | about 2000 | anobiid | |
| 368 | Maui | West Maui | — | 4000 | staphs & depressed <i>Brachypeplus</i> from <i>Freycinetia</i> ; single species of <i>Brachypeplus</i> from lilly | |
| 369 | Maui | Haleakala | 10 Apr 1894 | 5000 | <i>Brachypeplus</i> from flowers of tree <i>Lobelia</i> ; pair of minute black carabid under stones | |
| 370 | Maui | Haleakala | Apr 1894 | 5000+ | pair of <i>Proterhinus</i> in cop | |
| 371 | Maui | Haleakala | 1 Apr 1894 | about 5000 | | |
| 372 | Maui | Haleakala, Olinda woods | Apr 1894 | 4000 & downwards | | |
| 373 | Maui | Haleakala | Mar 1894 | 5000–6500 | big carabid in & above forests under stones, stumps & under bark (many spec. (310) | |
| 374 | Maui | Haleakala | 11 Apr 1894 | 7000–10,000 | <i>Barypristus</i> + ? gen. | |
| 375 | Maui | Haleakala | Mar, Apr 1894 | 5000 | | |
| 376 | Hawaii | Kilauea, near crater | Jul 1894 | | | |
| 377 | Maui | sandhills near Kahuku | Mar 1894 | | <i>Rhyncogonus</i> only | |
| 378 | Maui | Haleakala | Mar 1894 | 5000–6000 | 70 <i>Oodemas</i> ; 1 <i>Anotheorus</i> ; 2 <i>Heterhamphus</i> ; 34 <i>Dryophthorus</i> | |
| 379 | Maui | West Maui | Apr 1894 | 4000 | Coleops | |
| 380 | Maui | Haleakala | Apr 1894 | 7000 | top layer: find minute specimens | |
| 380 | Maui | Haleakala | Apr 1894 | 9000–10,000 | lower layer: all <i>Oodemas</i> | |
| 381 | Maui | Haleakala | Mar 1894 | 5000+ | | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|--------------------------|------------|-------------|--|--------------|
| 382 | Maui | Haleakala | Mar 1894 | 5000–6000 | The bronzy species <i>Barypristus</i> (54 specimens) always above the forest under stone. The <i>Blackburnia</i> (26 specimens) under logs, stems of tree fern, burnt stumps, &c in the forest & above it, but where forest once stood | |
| 383 | Maui | Haleakala | Mar 1894 | 4500–6000 | <i>Cyclothorax</i> from various situations | |
| 384 | Maui | Haleakala | Mar 1894 | 4500–5500 | 74 <i>Metromeni</i> , all in forest under bark (dead or alive), moss, logs, stones, &c; several species but no constancy of habitat. The large spec. beaten from dead tree | |
| 385 | Maui | Haleakala | Apr 1894 | about 5000 | contains 92 <i>Metromeni</i> , all under bark (mostly of <i>Acacia</i>) | |
| 386 | Hawaii | Kona | Oct 1892 | 3000 | from tree-ferns | |
| 387 | Hawaii | Kona | Sep 1892 | 3000 | | |
| 388 | Hawaii | Kona | Sep 1892 | 3000 | | |
| 389 | Oahu | mountains behind Waialua | Jan 1893 | about 800 | beaten | |
| 390 | Oahu | Halemano | Jan 1893 | 2000 | under bark | [= Helemano] |
| 391 | Oahu | Kaala mountains | Dec 1892 | 2000 | very minute <i>Homalota</i> 1 sp. | |
| 392 | Oahu | Kaala mountains | Dec 1892 | 2500 | | |
| 393 | Hawaii | Kona | | | | |
| 394 | Hawaii | Kona | Sep 1892 | 3000 | | |
| 395 | Hawaii | Kona | Sep 1892 | 3000 | | |
| 396 | Hawaii | Kona | Sep 1892 | 3000 | | |
| 397 | Hawaii | Kona | Sep 1892 | 3000 | | |
| 398 | Oahu | Honolulu | Oct 1892 | | at light | |
| 399 | Oahu | Nuuanu Valley, on ridges | Oct 1892 | | beaten | |
| 400 | Oahu | Nuuanu Valley, on ridges | [Oct 1892] | | on dead <i>Freycinetia</i> leaves | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|---------|---|-------------|-------------------|---|------------------------------------|
| 401 | Hawaii | Kona | 10 Sep 1892 | 4000 | | [Pulehua area] |
| 402 | Oahu | Mountains near Honolulu | 1892 | 2000+ | | |
| 403 | Oahu | Waianae Mts. | [1892] | 2000 | Heteroptera; 3 spec. beaten from native apple | |
| 404 | Hawaii | Kona, Kaawaloa | [1892] | 1500 | <i>P. boetica</i> & Coleoptera | |
| 405 | Kauai | — | May 1894 | sea level to 2500 | aculeates | |
| 406 | Oahu | Mountains behind Honolulu | Apr 1892 | 2000 | 2 sp. of Hemiptera | |
| 407 | Oahu | Waialua, coast | Mar 1892 | | Hemiptera (IX) | |
| 408 | Maui | | | | aculeates | |
| 409 | Lanai | | | | aculeates | |
| 410 | Kauai | | [1894] | 4000 | | [Halemanu area] |
| 411 | Oahu | Waialua | Mar 1892 | | Male, female, worker ant; Heteroptera | |
| 412 | Molokai | above Pelekunu, high forest | | | minute insects, chiefly staphs | |
| 413 | Maui | Haleakala | Mar 1894 | 5000 | <i>Disenochus?</i> under logs in forest; <i>Barypristus?</i> under stones above [forest]; <i>Cyclothorax</i> under bark slightly bronzy sp. also in moss | |
| 414 | Maui | Haleakala | Mar 1894 | 5000 | | |
| 415 | Oahu | Kawailoa gulch, far up | Apr 1893 | | in stem of tree fern | |
| 416 | Oahu | mountains behind Waialua, high up | Feb 1893 | | from dead tree | |
| 417 | Oahu | Nuuanu Valley, 4 miles out [from downtown Honolulu] | Oct 1892 | | | |
| 418 | Oahu | Waianae Mts. | Apr 1892 | | under bark of dead kukui tree | |
| 419 | Hawaii | Kona | Sep 1892 | 4000 | 3 beetles, dead bug & carabid under stone | |
| 420 | Oahu | Waianae Mts. | Apr 1892 | | Carabidae in cop | [Wai'anae (lee) side of the range] |
| 421 | Oahu | Kaala mountains | Mar 1892 | 1000–2000 | beetles under koa tree | |
| 422 | Oahu | Kaala mountains, gulches | Mar 1893 | 2000 | I could only find fragments of <i>Blackburnia</i> as sent | |
| 423 | Oahu | Nuuanu Valley | Nov 1892 | | in dead stem of guava | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|------------------------------------|---------|---|----------------|----------------|--|--|
| 424 | Molokai | Kahanui | Sep 1893 | | tipulid prey of <i>Crabro</i> with white mandibles & scape | |
| 425 | Oahu | Waianae Mts. | Apr 1892 | | | [Wai'anae (lee) side of the range] |
| 426 | Oahu | Waianae Mts. | Apr 1892 | about 3000 | | [Wai'anae (lee) side of the range] |
| 427 | Oahu | Kaala mountains | Mar 1892 | up to 1500 | | [after 26 March 1892] |
| 428 | Oahu | in the mountains near Waialua but locality lost | [1892] | | | |
| 429 | Molokai | — | May 1893 | near sea level | attached to the kiawe tree (an introduced <i>Acacia</i>) | |
| 430 | Oahu | Halemano | Jan 1893 | 2000+ | larvae in spirit "A" | [= Helemanu] |
| 431 | Hawaii | Kona | Aug 1892 | 4000 | 3 sp. of Heteroptera found in crevices of rotten logs | |
| 432 | Oahu | Honolulu | Oct & Nov 1892 | 2000+ | | |
| 433 | Hawaii | Kona | 4 Jul 1892 | 3000 | | |
| 434 | Hawaii | Kona | 1892 | | | |
| 435 | Hawaii | Kona | Jun 1892 | 3000 | parasitic on short eared owl | [after 20 June 1892] |
| Second Expedition 1895-1897 | | | | | | |
| | | | | | | |
| No. | Island | Locality | Date | Elev. | Other | Notes |
| 501 | Kauai | Halemanu | May 1895 | 4000 | coleops & aculeates | [there were no collection numbers between 435 and 501] |
| 502 | Kauai | Halemanu | May 1895 | | <i>Odynerus, Otiorhynchus, &c</i> | [about 4000 feet elevation] |
| 503 | Kauai | Halemanu | May 1895 | 4000 | anobiids &c | |
| 504 | Kauai | Halemanu | May 1895 | 4000 | coleops &c | |
| 505 | Kauai | Kaholuamano | Apr 1895 | | coleops &c | [about 3800 ft.] |
| 506 | Kauai | Makaweli | Apr 1895 | near sea level | Hymenoptera, aculeates | [between 2 and 9 April 1895] |
| 507 | Kauai | Halemanu | May 1895 | 4000 | Hymenoptera, aculeates | |
| 508 | Kauai | Kaholuamano | 16 Apr 1895 | | coleops | [about 3800 ft.] |
| 509 | Kauai | Kaholuamanu | Apr 1895 | | chiefly coleops, 1 very minute <i>Oligota</i> , also <i>Psocus</i> | [about 3800 ft.] |
| 510 | Kauai | Halemanu | May 1895 | 4000 | | [before 22 May 1895] |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|----------------------|-------------|---------------|---|---|
| 511 | Kauai | Kaholuamano | 12 Apr 1895 | | coleops, 28 spec. | [about 3800 ft.] |
| 512 | Kauai | Halemanu | May 1895 | | 5 pair anobiids in cop | |
| 513 | Kauai | Makaweli | May 1895 | coast to 2000 | aculeates | |
| 514 | Kauai | Halemanu | May 1895 | 4000 | <i>Odynerus, Mimesa, Rhyncogonus</i> | |
| 515 | Kauai | Makaweli | Apr 1895 | 2000+ | | [between 3 and 11 April 1895] |
| 516 | Kauai | Kaholuamano | 15 Apr 1895 | | coleops | [about 3800 ft.] |
| 517 | Kauai | Kaholuamano | 13 Apr 1895 | | coleops | [about 3800 ft.] |
| 518 | Kauai | Kaholuamano | Apr 1895 | | 3 agrions | [about 3800 ft.] |
| 519 | Kauai | Kaholuamano | Apr 1895 | | coleops & 2 Odyneri | [about 3800 ft.] |
| 520 | Kauai | Makaweli, near coast | May 1895 | | 2 dragonflies | [probably 23 May. Perkins left the next day for Honolulu] |
| 521 | Kauai | Kaholuamano | 15 Apr 1895 | | coleops, also <i>Crabro</i> & 1 <i>Odynerus</i> | [about 3800 ft.] |
| 522 | Kauai | Kaholuamano | Apr 1895 | | Hymenoptera aculeates | [about 3800 ft.] |
| 523 | Kauai | Kaholuamano | Apr 1895 | | coleops, some very small &c | [about 3800 ft.] |
| 524 | Kauai | Kaholuamano | 15 Apr 1895 | | coleops, small &c | [about 3800 ft.] |
| 525 | Kauai | Halemanu | 25 May 1895 | 4000 | | [the date must be a transcription error since Perkins left Halemanu on 22 May 1895] |
| 526 | Kauai | Kaholuamanu | Apr 1895 | | <i>Proterhinus</i> &c | [about 3800 ft.] |
| 527 | Kauai | Kaholuamanu | Apr 1895 | | coleops, chiefly carabids | [about 3800 ft.] |
| 528 | Kauai | Kaholuamanu | Apr 1895 | | coleops, many minute & also apterous parasitic Hymenoptera &c | [about 3800 ft.] |
| 529 | Kauai | Kaholuamanu | 13 Apr 1895 | | coleops | [about 3800 ft.] |
| 530 | Kauai | Halemanu | May 1895 | 4000 | chiefly coleops, 150–200 spec. small | [between 7 and 22 May 1895] |
| 531 | Hawaii | Kilauea | Aug 1895 | | | |
| 532 | Hawaii | Kilauea | Aug 1895 | | | |
| 533 | Hawaii | Kilauea | Sep 1895 | | chiefly aculeates, 1 small <i>Dryophthorus</i> & 1 ichneumon | [before 19 September 1895] |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|---------------------------------|----------|-------------|--|--------------------------|
| 534 | Hawaii | Kilauea | Jun 1895 | | aculeates &c | [after 12 June 1895] |
| 535 | Hawaii | Kilauea | Aug 1895 | | | |
| 536 | Hawaii | Kilauea | Jul 1895 | | | |
| 537 | Hawaii | Kilauea | Aug 1895 | | <i>Clytarlus</i> of koa | |
| 538 | Hawaii | Kilauea | Aug 1895 | | <i>Eopenthes</i> | |
| 539 | Hawaii | Kilauea | Aug 1895 | | | |
| 540 | Hawaii | Kilauea | Aug 1895 | | | |
| 541 | Oahu | Waianae mountains, lee side | Feb 1896 | 2000–3000 | Carabidae & Hymenoptera | |
| 542 | Oahu | Waianae mountains, lee side | Feb 1896 | | Carabidae | |
| 543 | Oahu | Waianae mountains, leeward side | Feb 1896 | 2000+ | <i>Forficula</i> , 3 sp. Carabidae | |
| 544 | Oahu | Waianae mountains, lee side | Feb 1896 | 2000+ | Carabidae | |
| 545 | Oahu | Waianae mountains, lee side | Feb 1896 | 2000–3000 | <i>Forficula</i> , Hymenoptera, Carabidae | |
| 546 | Oahu | Waianae mountains, leeward side | Feb 1896 | 2000–3000 | Carabidae | |
| 547 | Oahu | Waianae mountains, leeward side | Feb 1896 | about 3000 | <i>Proterhinus</i> &c | |
| 548 | Oahu | Waianae mountains, coast | Feb 1896 | | 1 small <i>Scymnus</i> ? with the big beetle | |
| 549 | Kauai | | Oct 1895 | 4000 | 1 <i>Parandra</i> | [probably Halemanu area] |
| 550 | Hawaii | above Hilo | Dec 1895 | 1800 | <i>Colpodiscus</i> &c | |
| 551 | Kauai | | Oct 1895 | 4000 | coleops | [probably Halemanu area] |
| 552 | Hawaii | Hilo | Jan 1896 | 2000+ | under bark | [before 28 January 1896] |
| 553 | Oahu | Waianae mountains, lee side | Feb 1896 | 2000–3000 | very minute sp. | |
| 554 | Kauai | | Oct 1895 | 4000 | <i>Oligota</i> | [probably Halemanu area] |
| 555 | Hawaii | Hilo | Dec 1895 | 1800 | <i>Oligota</i> | |
| 556 | Oahu | Waianae mountains, lee side | Feb 1896 | 3000 | pair of <i>Proterhinus</i> in cop | |
| 557 | Hawaii | Hilo | Jan 1896 | 2000 | found on decaying stem of tree <i>Lobelia</i> | [before 28 January 1896] |
| 558 | Hawaii | above Hilo | Dec 1895 | 1800 | <i>Brachypeplus</i> &c | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|---|----------|-------------|---|----------------------------|
| 559 | Hawaii | above Hilo, Amaula | Dec 1895 | 2000 | <i>Brachypeplus</i> , coleops | |
| 560 | Kauai | | Oct 1895 | 4000 | | [probably Halemanu area] |
| 561 | Hawaii | Hilo, Kaumana | Jan 1896 | 2000 | large <i>Prosopis</i> & Heteroptera &c | [before 28 January 1896] |
| 562 | Hawaii | Hilo, Kaumana | Jan 1896 | 2000 | | [before 28 January 1896] |
| 563 | Kauai | | Oct 1895 | 4000 | coleops | [probably Halemanu area] |
| 564 | Kauai | | Oct 1895 | 4000 | | [probably Halemanu area] |
| 565 | Hawaii | above Hilo | Dec 1895 | 1800 | aculeates, <i>Eopenthes</i> &c | |
| 566 | Hawaii | above Hilo | Dec 1895 | 1800 | <i>Xyletobius</i> | |
| 567 | Hawaii | above Hilo | Dec 1895 | 1800 | staphs &c | |
| 568 | Hawaii | Kilauea | Jul 1895 | | | |
| 569 | Hawaii | Olaa | ?1895 | | coleops, Hymenoptera, <i>Forficula</i> , 2 sp. <i>Prosopis</i> &c | |
| 570 | Hawaii | Olaa | Jul 1895 | | <i>Eopenthes</i> , also Diptera | |
| 571 | Oahu | Honolulu mountains | 1895 | 2000+ | <i>Heterhamphus</i> | [after 23 March 1895] |
| 572 | Hawaii | Olaa | Jun 1895 | | <i>Eopenthes</i> , <i>Colpodiscus</i> , <i>Prosopis</i> &c | [after 16 June 1895] |
| 573 | Hawaii | Kilauea | Sep 1895 | | agriions | [before 19 September 1895] |
| 574 | Hawaii | Kilauea | Aug 1895 | | large black <i>Fornax</i> , cistelid &c | |
| 575 | Hawaii | Kilauea | Jul 1895 | | | |
| 576 | Hawaii | Kilauea | Jul 1895 | | anobiids, <i>Fornax</i> &c | |
| 577 | Hawaii | Kilauea | Aug 1895 | | | |
| 578 | Hawaii | Kilauea | Jul 1895 | | | |
| 579 | Hawaii | Kilauea | Aug 1895 | | | |
| 580 | Hawaii | Kilauea | | | <i>Clytarlus</i> of ohia lehua (<i>Metrosideros</i>) | |
| 581 | Hawaii | Kilauea | | | <i>Clytarlus</i> , large <i>Fornax</i> &c | |
| 582 | Maui | Haleakala | May 1896 | 5000 | | |
| 583 | Maui | Haleakala | May 1896 | 5000 | in cop | |
| 584 | Hawaii | Kilauea | Jun 1895 | 4000 | <i>Clytarlus</i> , <i>Eopenthes</i> , cossonid, larger staph, aculeates &c | |
| 585 | Hawaii | Kilauea, Hilo road, 1.5 miles [from Kilauea] | Aug 1895 | | <i>Clytarlus</i> of koa, Carabidae | |
| 586 | Hawaii | Kilauea | Aug 1895 | | | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|---------|--|----------|-------------|--|--|
| 587 | Hawaii | Kilauea | Jul 1895 | | aculeates &c | |
| 588 | Molokai | Molokai mountains | Jun 1896 | | | |
| 589 | Molokai | Molokai mountains | Jun 1896 | about 4000 | contains inter alia several <i>Blackburnia</i> & 1 <i>Atrachynemis</i> | |
| 590 | Molokai | Molokai mountains | Jun 1896 | | common coleops, <i>Forficula</i> &c | |
| 591 | Molokai | Molokai mountains | Jun 1896 | 4500–5000 | Carabidae all from boggy mountain top, mostly under moss on trees | |
| 591 | Molokai | Molokai mountains | Jun 1896 | 3000 | everything else [other than that above] | |
| 592 | Molokai | Molokai mountains, boggy plateau below the densest forest | Jun 1896 | about 4000 | contains several good things (Carabidae) | |
| 593 | Molokai | Molokai mountains | Jun 1896 | | | |
| 594 | Molokai | Molokai mountains | Jun 1896 | | | |
| 595 | Molokai | Molokai mountains, on side of stream on bottom of gulch | Jun 1896 | about 4000 | Carabidae | |
| 596 | Molokai | Molokai mountains, boggy plateau below the densest forest | Jun 1896 | about 4000 | Carabidae | |
| 597 | Maui | Haleakala | May 1896 | 4000+ | | [all May 1896 Haleakala (Nos. 597– 625) are on the Olinda side] |
| 598 | Maui | Haleakala | May 1896 | 4000+ | | |
| 599 | Maui | Haleakala | May 1896 | 4000+ | | |
| 600 | Maui | Haleakala | May 1896 | 5000 | | |
| 601 | Maui | Haleakala | May 1896 | 4800+ | | |
| 602 | Maui | Haleakala, above forest | May 1896 | | 3 sp. Carabidae | |
| 603 | Maui | Haleakala | May 1896 | 5000 | | |
| 604 | Maui | Haleakala | May 1896 | 5000 | | |
| 605 | Maui | Haleakala | May 1896 | 5000 | | |
| 606 | Maui | Haleakala | May 1896 | 5000 | | |
| 607 | Maui | Haleakala | May 1896 | 5000 | | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|----------------------|----------|-------------|---|------------------------|
| 608 | Maui | Haleakala | May 1896 | 5000 | | |
| 609 | Maui | Haleakala | May 1896 | 5000 | | |
| 610 | Maui | Haleakala | May 1896 | 5000 | | |
| 611 | Maui | Haleakala | May 1896 | 5000 | | |
| 612 | Maui | Haleakala | May 1896 | 5000 | | |
| 613 | Maui | Haleakala | May 1896 | 5000 | | |
| 614 | Maui | Haleakala | May 1896 | 5000 | | |
| 615 | Maui | Haleakala | May 1896 | 5000 | | |
| 616 | Maui | Haleakala | May 1896 | 5000 | | |
| 617 | Maui | Haleakala | May 1896 | 5000 | <i>Fornax</i> bred from long thin wireworm larvae | |
| 618 | Maui | Haleakala | May 1896 | 5000 | | |
| 619 | Maui | Haleakala | May 1896 | 4000 | | |
| 620 | Maui | Haleakala | May 1896 | 4000 | | |
| 621 | Maui | Haleakala | May 1896 | 4000 | | |
| 622 | Maui | Haleakala | May 1896 | 4000 | | |
| 623 | Maui | Haleakala | May 1896 | 4000 | | |
| 624 | Maui | Haleakala | May 1896 | 7000–10,000 | | |
| 625 | Maui | Iao Valley | May 1896 | | | |
| 626 | Oahu | Mountains near | Jul 1896 | | | |
| 627 | Hawaii | Kilauea? | Sep 1896 | | | |
| 628 | Maui | Wailuku & Iao Valley | | | Hymenoptera aculeates & cossonids | |
| 629 | Maui | Olinda | May 1896 | 3000–4000 | | |
| 630 | Hawaii | Kona | 1896 | 4000 | | |
| 631 | Kauai | high plateau | Aug 1896 | | | |
| 632 | Hawaii | | Aug 1896 | 4000 | | |
| 633 | Hawaii | | | 4000 | <i>Clytarlus, Odynerus</i> | |
| 634 | Hawaii | Kilauea | Aug 1895 | | aculeates | |
| 635 | Hawaii | Olaa | Nov 1896 | | aculeates Hymenoptera | |
| 636 | Maui | Haleakala | Oct 1896 | 5000+ | | |
| 637 | Hawaii | Volcano, Hilo Road | Aug 1896 | | Koebele | [Volcano Village area] |
| 638 | Hawaii | Kilauea | | | | |
| 639 | Hawaii | Kilauea | | | 115 aculeates &c | |
| 640 | Kauai | | Jul 1896 | 4000 | Rhynchota, Hymenoptera &c | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|---|---------------|-------------|--|-------------------------|
| 641 | Kauai | | Jan, Feb 1897 | 2000–3000 | | |
| 642 | Hawaii | Kilauea | Dec 1896 | | | |
| 643 | Hawaii | Hilo | Feb 1897 | | earwigs | |
| 644 | Hawaii | Kilauea | Jun 1895 | | aculeates | |
| 645 | Hawaii | above Hilo | Dec 1895 | 1800 | 1 <i>Eopenthes</i> & staphs | [Kaumana area] |
| 646 | Hawaii | Kilauea | Jun 1895 | | | |
| 647 | Maui | Mt. Lahaina | Jan 1897 | 3000 | | |
| 648 | Kauai | | Jan, Feb 1897 | 2000–3000 | (= 641) | |
| 649 | Hawaii | [Kilauea area] | Sep 1896 | 4000 | Hymenoptera aculeates | |
| 650 | Maui | Lahaina, coast | Jan 1897 | | | |
| 651 | Oahu | Honolulu | Nov 1896 | | | |
| 652 | Kauai | coast | Jan 1897 | | | |
| 653 | Oahu | Honolulu mountains | Mar 1897 | 2000 | | |
| 654 | Oahu | Waianae coast | Jan 1897 | | | |
| 655 | Maui | Haleakala crater | Oct 1896 | | | |
| 656 | Hawaii | Kilauea | Aug 1896 | | | |
| 657 | Hawaii | Olaa | Dec 1896 | | | |
| 658 | Hawaii | Kona | Dec 1896 | | coll. Koebele | |
| 659 | Hawaii | Kilauea | | | | |
| 660 | — | on the SS Kinau running from Honolulu to Hawaii | 1896 | | | |
| 661 | Maui | Haleakala | Oct 1896 | 5000 | contains amongst others a unique carabid (near <i>Mauna</i>) found under lichen on stem of <i>Pelea</i> | |
| 662 | Oahu | Waianae coast | 1896 | | | |
| 663 | Kauai | Makaweli, near coast | May 1895 | | | [between 22 and 25 May] |
| 664 | Kauai | Lihue | 1896 | | | [14 July–9 August] |
| 665 | Maui | Haleakala, above forest | May 1896 | | | [Olinda side] |
| 666 | Oahu | Waianae coast | Nov 1895 | | | |
| 667 | Oahu | Mountains near Honolulu | | 2000–3000 | <i>Chalcis</i> (variegated) & coccinellid introduced by A. Koebele | |
| 668 | Kauai | Makaweli | Feb 1897 | 2500 | | |
| 669 | Oahu | Waianae, lee side | Feb 1896 | 2000–3000 | | |
| 670 | Oahu | Waianae, lee side | | 2000–3000 | anobiids &c | |
| 671 | Kauai | Lihue | Jul 1896 | | | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|-------------------------|------------|-------------|--|--------------------------|
| 672 | Oahu | Waianae, coast | Feb 1896 | | | |
| 673 | Hawaii | Kilauea | Dec 1896 | | | |
| 674 | Maui | Lahaina | Jan 1897 | 2000+ | | |
| 675 | Maui | Lahaina, coast | Jan 1897 | | | |
| 676 | Kauai | — | Aug 1897 | 4000 | duplicate Carabidae | [probably Halemanu area] |
| 677 | Hawaii | Kilauea | Dec 1896 | | male with long plate above clypeus & long forked process to 8th vent. seg. female with long cheeks | |
| 678 | Hawaii | Kilauea | [Dec 1896] | | male with short plate above clypeus & hardly forked process to 8th vent. seg. female with shorter cheeks | |
| 679 | Maui | Haleakala | Oct 1896 | 4500–5000 | small <i>Oodemas</i> &c | |
| 680 | Maui | Haleakala | Oct 1896 | 4000+ | | |
| 681 | Oahu | Honolulu mountains | Sep 1896 | 2000 | | |
| 682 | Kauai | high plateau | Aug 1896 | | | [before 9 August] |
| 683 | Hawaii | Kilauea | Aug 1896 | | | [after 18 August] |
| 684 | Hawaii | Kilauea | Dec 1896 | | | |
| 685 | Hawaii | Kilauea | Sep 1896 | | | |
| 686 | Hawaii | Kilauea | Aug 1896 | | | [after 18 August] |
| 687 | Hawaii | Kilauea | | | about 40 <i>Ophius</i> all resting on 2 adjoining leaves 2/3 or more of those present escaped | |
| 688 | Hawaii | Olaa | Sep 1896 | | | [possibly 2000 ft.] |
| 689 | Maui | Iao Valley | Sep 1896 | | | |
| 690 | Kauai | Makaweli | Jan 1897 | 2000+ | | |
| 691 | Hawaii | Kilauea | Jul 1895 | | | |
| 692 | Hawaii | Kilauea | Jun 1895 | | | |
| 693 | Oahu | Mountains near Honolulu | Jul 1895 | | | |
| 694 | Hawaii | Olaa | Aug 1896 | 2000 | | |
| 695 | Hawaii | Kilauea | Sep 1896 | 4000 | | [near crater] |
| 696 | Oahu | Honolulu mountains | Aug 1896 | | | |
| 697 | Hawaii | Kaumana, Hilo | Jan 1897 | 2000 | many spec. some very minute | |

| | | | | | | |
|-------------------------------------|---------------|-----------------------------|---------------|--------------|--|---|
| 698 | Maui | Lahaina | Jan 1897 | | <i>Cis</i> & Homoptera from mts; the rest from the plains; the minute chalcid from scale | |
| 698 | Oahu | Honolulu | | | longicorn | |
| 699 | Maui | Haleakala | Oct 1896 | 4000–5000 | small insects | |
| 700 | Maui | Haleakala | Oct 1896 | | chiefly from flow of crater | |
| 701 | Kauai | Makaweli | Jan, Feb 1897 | | <i>Plagithmysus aequalis</i> | |
| 702 | Kauai | ?probably | 1896 | 4000 | | |
| 703 | Kauai | Makaweli | Feb 1897 | 2500 | | [if the elevation is correct, this is referring to the Makaweli “section” of Kauai and not the coastal Makaweli town] |
| 704 | Kauai | | Aug 1896 | 4000 | | [Halemanu area] |
| 705 | Maui | Haleakala crater | Oct 1896 | | dipteron sp.? Trypetidae bred from flowers of silversword | |
| | | | | | | |
| Third Expedition (1900–1901) | | | | | | |
| | | | | | | |
| No. | Island | Locality | Date | Elev. | Other | Notes |
| 750 | Oahu | Wahiawa | Apr 1901 | 1800 | | [there were no numbers between 705 and 750] |
| 751 | Oahu | Waianae mountains, Mokuleia | Apr 1901 | 2000 | | |
| 752 | Oahu | Waianae mountains, Mokuleia | May 1901 | 2000 | | |
| 753 | Oahu | mountains near Honolulu | 1900 | | | |
| 754 | Oahu | Honolulu | Mar 1901 | | | |
| 755 | Oahu | Honolulu | Mar 1901 | | | |
| 756 | Oahu | Honolulu mountains | Feb 1901 | | | |
| 757 | Oahu | Honolulu | 1900 | | | |
| 758 | Oahu | Honolulu | 1900 | | | |
| 759 | Oahu | Waialua, coast | May 1901 | | | |
| 760 | Oahu | Waialua | Sep 1900 | | | |
| 761 | Oahu | Waialua | Sep 1900 | | | |
| 762 | Oahu | Waialua | Mar 1901 | | | |
| 763 | Oahu | Waialua | Sep 1900 | | | |
| 764 | Hawaii | — | — | | | [unknown date and locality] |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|-------------------------------|---------------|-------------|---|---|
| 765 | Oahu | [probably Honolulu mountains] | Sep 1900 | | presumably Honolulu Mts. the boy who wrote the list left out the locality | |
| 766 | Oahu | Honolulu | Feb 1901 | | | |
| 767 | Oahu | Honolulu | Nov 1901 | | | |
| 768 | Oahu | Kawailoa | Apr 1901 | 1800 | | |
| 769 | Oahu | Honolulu mountains | Jun 1901 | 2000 | | |
| 770 | Oahu | Honolulu (in the town) | Jul 1901 | | | |
| 771 | Kauai | — | — | | | |
| 772 | Maui | Haleakala | 1900 | 3000 | | |
| 773 | Oahu | Waialua | Jun 1901 | | if native beetles the elevation is 1800 | [the 1800 ft locality is near Kawailoa] |
| 774 | Oahu | head of Pauoa, near Honolulu | Nov 1900 | | | |
| 775 | Maui | Iao Valley | Sep 1901 | | | |
| 776 | Oahu | Honolulu coast | Jan 1901 | | | |
| 778 | Oahu | Wahiawa | Apr 1901 | | | |
| 779 | Hawaii | Hualalai | Sep 1900 | | | |
| 780 | Oahu | Waianae coast | Jan 1901 | | | |
| 781 | Oahu | Back of Tantalus | Jun 1901 | | | |
| 782 | Oahu | Konahuanui ridge | Nov 1900 | | | |
| 783 | Oahu | Konahuanui ridge | Dec 1900 | | | |
| 784 | Oahu | back of Tantalus | Nov 1900 | | <i>Pipturus, Pisonia</i> | |
| 785 | Oahu | Honolulu mountains | Aug 1900 | | | |
| 786 | Oahu | Honolulu mountains | Jul 1900 | | | |
| 787 | Oahu | Honolulu mountains | Dec 1900 | | | |
| 788 | Oahu | Honolulu mountains | Dec 1900 | | | |
| 789 | Oahu | Honolulu mountains | Jun 1900 | | | |
| 790 | Oahu | Honolulu mountains | Sep 1900 | | | |
| 791 | Oahu | back of Tantalus | Dec 1900 | | <i>Pipturus</i> | |
| 792 | Oahu | Honolulu end of Koolau range | Dec 1900 | | | |
| 793 | Oahu | Honolulu end of Koolau range | Nov, Dec 1900 | | | |
| 794 | Oahu | back of Tantalus | Nov, Dec 1900 | | | |
| 795 | Hawaii | Kona | | | | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|------------------------------|----------|-------------|-----------------------------------|-------|
| 796 | Oahu | Waialua coast | 1900 | | | |
| 797 | Hawaii | Kona | 1900 | | | |
| 798 | Oahu | Waialua coast | 1900 | | | |
| 799 | Oahu | Honolulu | Oct 1900 | | | |
| 800 | Oahu | Honolulu mountains | Dec 1900 | | <i>Cis</i> bred from kukui fungus | |
| 801 | Oahu | — | | | | |
| 802 | Kauai | Lihue | 1900 | | | |
| 803 | Oahu | Nuuanu Pali | Dec 1900 | | | |
| 804 | Oahu | Honolulu mountains | Jul 1900 | | <i>Cis</i> with 1 Hymenoptera | |
| 805 | Oahu | Honolulu mountains | | | | |
| 806 | Oahu | Waialua coast | Oct 1901 | | | |
| 807 | Hawaii | Kona | | | | |
| 808 | Hawaii | North Kona | | | | |
| 809 | Maui | Haleakala | | | | |
| 810 | Oahu | Konahuanui ridge | Feb 1901 | | all on <i>Pelea</i> | |
| 811 | Oahu | Koolau, high mountains | Feb 1901 | | flower of <i>Astelia</i> | |
| 812 | Hawaii | Hualalai | 1900 | | | |
| 813 | Oahu | Honolulu mountains | summer | | | |
| 814 | Hawaii | North Kona, Hualalai | 1900 | | | |
| 815 | Hawaii | Olaa | | | | |
| 816 | Oahu | head of Pauoa | Dec 1900 | | | |
| 817 | Maui | — | 1900 | | | |
| 818 | Oahu | head of Pauoa Valley | Dec 1900 | | | |
| 819 | Oahu | back of Tantalus | Nov 1900 | | <i>Pipturus</i> | |
| 820 | Oahu | back of Tantalus | Nov 1900 | | <i>Pisonia</i> | |
| 821 | Oahu | Honolulu end of Koolau range | Dec 1900 | | | |
| 822 | Oahu | — | Dec 1900 | | <i>Pelea</i> | |
| 823 | Hawaii | — | | | | |
| 824 | Oahu | Honolulu mountains | Jan 1901 | | | |
| 825 | Oahu | Honolulu mountains | Dec 1900 | | Homoptera, all <i>Pipturus</i> | |
| 826 | Maui | Haleakala | 1901 | 2000–3000 | | |
| 827 | Oahu | Honolulu mountains | Jul 1900 | | on "kalia" | |
| 828 | Oahu | Waialua end of Koolau range | Sep 1900 | | | |
| 829 | Kauai | Waimea watershed | Apr 1901 | | | |
| 830 | Hawaii | — | Nov 1900 | | bred | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|------------------------------|---------------|-------------|---|-------|
| 831 | Hawaii | — | Oct 1900 | | bred | |
| 832 | Oahu | Honolulu mountains | Dec 1900 | | <i>Cis & Proterhinus</i> on <i>Pipturus</i> | |
| 833 | Oahu | Honolulu mountains | Dec 1900 | | kadua tree | |
| 834 | Oahu | Honolulu city | Jun 1900 | | | |
| 835 | Oahu | [Nuuanu] Pali | Dec 1900 | | <i>Dracaena aurea</i> | |
| 836 | Oahu | Honolulu | Jul 1900 | | | |
| 837 | Oahu | Honolulu | | | bred from <i>Pipturus</i> | |
| 838 | Oahu | Honolulu mountains | 31 Jul 1900 | | bred from <i>Pipturus</i> | |
| 841 | Oahu | Koolau range | Jul 1900 | 2000 | | |
| 844 | Kauai | — | 1900 | | | |
| 845 | Mauai | Haleakala | 1900 | 3000 | | |
| 846 | Oahu | Koolau range | 1900 | | | |
| 847 | Oahu | Honolulu mountains | Dec 1900 | | | |
| 848 | Oahu | Honolulu | Nov 1900 | | | |
| 849 | Oahu | Honolulu city | | | | |
| 850 | Oahu | Honolulu, back of Tantalus | Nov 1900 | | | |
| 851 | Oahu | Waialua | Apr 1900 | | | |
| 852 | Mauai | Haleakala | — | 5000 | | |
| 853 | Oahu | Honolulu | Apr 1900 | | | |
| 854 | Oahu | Honolulu | Jan 1901 | | on tamarinds | |
| 855 | Oahu | Honolulu | Sep 1900 | | | |
| 856 | Oahu | Honolulu mountains | Dec 1900 | | on <i>Pipturus</i> | |
| 857 | Oahu | north west Koolau | Aug 1901 | | | |
| 858 | Oahu | north west Koolau | May 1901 | | | |
| 859 | Oahu | north west Koolau | Aug 1901 | | | |
| 860 | Oahu | Honolulu mountains | 1900 | | | |
| 861 | Oahu | Honolulu mountains | Dec 1900 | | | |
| 862 | Oahu | Honolulu mountains | Jun 1901 | | | |
| 863 | Oahu | north west Koolau | Jul 1901 | | | |
| 864 | Oahu | north west Koolau | May 1901 | | | |
| 865 | Oahu | north Koolau range | Aug 1901 | | | |
| 866 | Oahu | Honolulu mountains | Jan 1900 | | | |
| 867 | Oahu | Konahuanui ridge | Mar 1901 | | | |
| 868 | Oahu | south east coast | Jan 1901 | | | |
| 869 | Oahu | Honolulu mountains | Nov 1900 | | <i>Pipturus</i> | |
| 870 | Oahu | Honolulu end of Koolau range | Nov, Dec 1900 | | | |

| No. | Island | Locality | Date | Elev. (ft.) | Other | Notes |
|-----|--------|-------------------------|---------------|-------------|-----------------------------|-------|
| 871 | Oahu | Konahuanui ridge | Nov 1900 | | | |
| 872 | Oahu | Honolulu mountains | Sep 1900 | | | |
| 873 | Oahu | Waialua | Sep 1900 | 2000 | | |
| 874 | Maui | Haleakala | — | 3000 | | |
| 875 | Oahu | | Mar 1901 | | | |
| 876 | Oahu | Honolulu mountains | Oct 1900 | | | |
| 877 | — | missing | | | | |
| 878 | Oahu | Waialua | | | | |
| 879 | Maui | Haleakala | Sep 1901 | 2000–3000 | | |
| 880 | Maui | Haleakala | 1901 | 2000–3000 | | |
| 881 | Maui | Haleakala, saw mill | 1901 | | | |
| 882 | Kauai | Lihue | | | | |
| 883 | Oahu | Honolulu mountains | Nov, Dec 1900 | | | |
| 884 | Oahu | high Honolulu mountains | Dec [1900] | | | |
| 885 | Oahu | Honolulu mountains | Sep 1900 | | | |
| 886 | Oahu | Honolulu mountains | Jun, Jul 1900 | | | |
| 887 | Oahu | Koolau | | 2500 | on koa | |
| 888 | Oahu | Honolulu mountains | Aug 1900 | | | |
| 889 | Oahu | | | | Koebele | |
| 890 | | ?Oahu or ?Hawaii | | | doubtful | |
| 891 | Oahu | Waianae | Apr 1900 | | | |
| 892 | — | — | — | | <i>Fornax</i> bred from koa | |
| 893 | Oahu | north west Koolau | Mar 1901 | | | |
| 894 | Hawaii | Hualalai | 1900 | | | |
| 895 | Oahu | Koolau range | Aug 1900 | | | |
| 896 | Oahu | Honolulu | Dec 1900 | | bred | |
| 897 | Maui | Haleakala | 1900 | | | |
| 898 | Hawaii | — | | | bred | |
| 899 | Hawaii | — | | 4000 | | |
| 900 | Oahu | back of Tantalus | | | | |
| 901 | Oahu | Honolulu? | | | | |
| 902 | Oahu | back of Tantalus | | | | |
| 903 | — | no label | | | | |
| 904 | Oahu | Honolulu mountains | Dec 1900 | | | |
| 905 | Oahu | Tantalus | Dec 1900 | | bred | |
| 906 | Hawaii | Kona | Oct 1900 | | from <i>Euphorbia</i> | |

