

86

PLANTS OF THE MANUA ISLANDS


BY
T. G. YUNCKER

BERNICE P. BISHOP MUSEUM
BULLETIN 184

HONOLULU, HAWAII
PUBLISHED BY THE MUSEUM
1945

ISSUED APRIL 17, 1945

Am
101
B54
no. 184



General Library
Anthrop. Div. - Jan. 1941
W.H.
3-29-46
51592

Plants of the Manua Islands

By T. G. YUNCKER

DEPARTMENT OF BOTANY, DEPAUW UNIVERSITY

INTRODUCTION

The volcanic islands of Ofu, Olosega, and Tau are known collectively as Manua, now part of American (Eastern) Samoa. They are situated between 169 and 170 degrees west longitude and 14 and 15 degrees south latitude and are about 67 miles east of Pago Pago on the island of Tutuila. Tau, containing about 14 square miles, is the largest and attains an altitude of over 900 meters. It is more or less dome shaped with precipitous coasts on the northern and western sides. Ofu and Olosega, only a few miles distant, are rugged and picturesque and are separated from each other by a narrow and shallow strait.

The islands lie outside the regular trade routes and are rarely visited by ships other than occasional small trading vessels or those making official governmental visits. The only white persons residing there are a few connected with the United States Navy Department, which has jurisdiction over the islands, and occasional transient visitors. There are about 2,500 natives living on the three islands, who subsist principally on sea food, bananas, taro, breadfruit, and coconuts. Breadfruit trees grow in large numbers in and about every village. Taro, bananas, and coconuts are extensively cultivated in plantations. Tinned foods are also used to some extent, salmon being a favorite.

The original vegetation has now been exterminated to a large extent on the lowlands and lower hillsides to provide for coconut and banana plantations. It still persists, however, at the higher altitudes and in ravines and other areas too rough for cultivation. The vegetation in these regions, especially on Tau, is exceedingly dense and travel off the trails is difficult.

A number of botanists have made collections on the islands. Among the older collections are those made by Reinecke, Powell, Brackenridge, and McMullen. More recently, collections have been made by D. W. Garber, between 1921 and 1925; by Anna and Wray Harris, mostly during 1938; and by me. I spent several weeks on the islands of Tau and Ofu during October and November of 1939. Christophersen in his Flowering Plants of Samoa (B. P. Bishop Mus., Bull. 128 and 154, 1935 and 1938) cites the collections made by Garber. Aside from this, no attempt at a complete listing of the plants has been made. The Garber and the Harris collections included in the present list, as well as most of the other scattered citations, are in the Bernice P. Bishop Museum herbarium and have been examined by me. I have not seen collections

made by earlier visitors to the islands. They are now mostly in European herbaria and, with a few exceptions, are not included. During the summer of 1939, Paul L. Guest of Honolulu made a survey of the economic trees of American Samoa, including those of Manua. His manuscript report in the Bishop Museum library is not, unfortunately, supported by herbarium materials. However, it has been checked against this list and his observations have been included wherever they extend the range or report a species overlooked by other botanists.

The present list includes 37 species of mosses, 53 species of pteridophytes, 95 species of monocotyledons, and 236 species of dicotyledons, or a total of 421 species. More careful botanizing, particularly at high altitudes on Tau, will probably bring to light several species hitherto overlooked.

The spelling of native names, obtained from natives or from the publications of Christophersen and of Setchell (*Vegetation of Tutuila Island, Carnegie Inst., Pub. 20, 1924*), have been checked by Dr. Peter H. Buck.

The islands were visited and the collections made while I was on an appointment as Bishop Museum Fellow in Yale University. I am deeply indebted to Dr. Peter H. Buck, Director of Bishop Museum, for placing the facilities of the Museum at my disposal and for assistance in arranging the itinerary; to Captain E. W. Hanson, then Governor of American Samoa, and the other officers and men of the United States Navy stationed in the islands for transportation to Manua from Pago Pago and for the many kindnesses and courtesies which were shown me while in Samoa; and to Chiefs Tufele and Lafite of Tau who placed a guest house at my disposal during my stay on the island and who assisted in my work in every way possible.

I am also indebted to a number of specialists who have cooperated in the identifications of the specimens and have contributed bibliographic data. Miss Marie Neal, botanist at Bishop Museum, assisted in the preliminary identification of much of the material and helped in making comparisons with specimens in the Museum herbarium; Mr. E. B. Bartram contributed the list of mosses; Dr. E. B. Copeland named the ferns, except those collected by Garber and Harris and identified by Dr. Carl Christensen; Dr. F. R. Fosberg identified a number of specimens, particularly those in the Rubiaceae and Ebenaceae, and in addition made a number of nomenclatorial suggestions; Mrs. Agnes Chase named the grasses; Dr. L. O. Williams identified the orchids; Mr. Hugh O'Neill identified the sedges; Dr. E. D. Merrill and Dr. A. C. Smith named a number of sterile or otherwise obscure specimens; Dr. Leon Croizat supplied the description of a new species of *Securinera*; and Dr. H. W. Rickett and Mr. C. V. Morton contributed nomenclatorial data. To all who have assisted in any way I wish to express my thanks.

BRYOPHYTA

MUSCI¹

This list of species from the Manua islands represents a substantial and interesting addition to the moss flora of Samoa. With two minor exceptions, the collections are all from the larger island of Tau and no doubt give a fairly complete outline of the mosses of this outlying region. At least nine species are new to Samoa, as far as I know; and four reductions to synonymy are proposed based on the information at hand.

One cannot fail to be impressed with the strong west to east course of geographical distribution shown in the list of species. This is especially true of *Calymperes*, a genus with marked insular tendencies, and may indicate a continuous land connection from Malaysia through Polynesia at no very remote period.

FAMILY FISSIDENTACEAE

Genus *Fissidens* Hedwig

Fissidens mangarevensis Montagne, Voy. au Pôle Sud, Crypt., 344, 1845.

Tau: on tree trunk in forest, alt. about 150 m., Oct. 1, 1939, *Yuncker 9154*; Palapala, gulch above Palapala Hill, on mountainside, damp native forest, banks of dry stream on trunks of tree fern and other trees, 1.5 miles from shore, alt. 270-300 m., *W. Harris 54b*.

New to Samoa. Distribution: Mangareva, Austral Islands, Rapa, Fiji, Solomon Islands.

FAMILY LEUCOBRYACEAE

Genus *Leucophanes* Hampe

Leucophanes candidum (Hornschuch) Lindberg, Oefvers. Vet. Akad. Förh., 602, 1854.

Tau: north slope of Lepue, near summit of cone, under cover of forest, on lava boulder, one third mile from shore, alt. 285 m., *W. Harris 1759*.

Distribution: Ceylon, Malaysia, New Guinea, Philippines, Fiji.

Leucophanes pungens Fleischer, unpublished manuscript.

Tau: Tunoa Peak, summit, ridge, medium damp, one eighth mile from shore, alt. 189 m., growing on dead logs in forest, *W. Harris*.

Distribution: Fiji.

Genus *Exodictyon* Cardot

Exodictyon scabrum (Mitten) Cardot, Rev. Bryol., 7, 1899.

Tau: Faleulu draw, hillside, in native forest on living tree trunks, damp, 1 mile from shore, alt. 480 m., *W. Harris*.

¹ Contributed by Edwin B. Bartram.

***Exodictyon dentatum* (Mitten) Cardot, Rev. Bryol., 7, 1899.**

Tau: Faleulu Valley, south of Muao's house, on damp dead wood, light forest, hillside, moderate slope, 0.75 mile to shore, alt. 210 m., *W. Harris* 404; Palapala, gulch above Palapala Hill, on mountainside, damp native forest, banks of a dry stream, on trunks of tree ferns and other trees 1.5 miles from shore, alt. 270-300 m., *W. Harris* 53.

Distribution: Tahiti, New Hebrides, New Caledonia.

FAMILY CALYMPERACEAE**Genus *Calymperes* Swartz**

***Calymperes hyophylaceum* C. Müller**, in Besch. Essai Calymp. Sci. Nat., Ann. VIII, Bot. 1: 287, 1895.

Tau: Papatea, South Cove, lower edge of forest, steep hillside, crater rim, 5-30 feet from shore, alt. 1.5-6 m., growing on lower trunks of coconut tree and exposed foot of forest tree, *A. Harris* 418; Papatea, South Cove, on tuff at shore, under lower edge of strand forest, damp crater rim, steep hillside, 5-30 feet from shore, *A. Harris* 464; upper Luma plantation, east of pig fence, along Vaosa trail, plantation in amphitheater plateau, moderate slope, damp, 2-10 feet above ground on trunks of coconut and breadfruit trees, 0.75 mile from shore, alt. 90 m., *W. Harris* 444a.

New to Samoa. Distribution: Sumatra, Java, Philippines.

***Calymperes samoanum* Bescherelle**, Essai Calymp. Sci. Nat., Ann. VIII, Bot. 1: 302, 1895.

Tau: Judds Crater, northwest rim, on trunks and limbs of trees, in damp, rather open forest, 0.75 mile to shore, alt. 150 m., *W. Harris* 413, 414; on coconut tree, alt. 250-300 m., *Yuncker* 9243.

Endemic.

***Calymperes strictifolium* (Mitten) Roth, Hedwigia** 51: 127, 1911.

***Syrrhopodon tuberculosus* Dixon and Theriot**, Linn. Soc. Bot., Jour. 43: 303, 1916.

Tau: Tavalogi Ridge, west end, ridge summit, under light forest cover, moss taken from north face of tuff capping, medium damp, one eighth mile from sea, alt. 143 m., *W. Harris*; Tavalogi Ridge, west end, north face, summit of ridge, cliff face, under forest, medium dry, on tuff rock, one eighth mile from shore, alt. 100 m., *W. Harris*; Faleulu Valley, one eighth mile south of Muao's house, growing on dead logs and fallen wood lying on moist soil under cover of light forest, hillside, gentle slope, 0.75 mile from shore, alt. 213 m., *Harris* 436b.

Upon examination of part of the type collection of this species, the highly characteristic tuberculate costa and leaf margins at once suggested a close relationship with *S. tuberculosus* Dixon and Theriot. Further comparisons failed

to reveal any noteworthy distinctions. Dixon has compared the type material of his species with *C. strictifolium* and agrees that they are conspecific. Under the circumstances, *S. tuberculosus* safely can be reduced to synonymy. It is odd that Mitten failed to mention the striking character that at once distinguishes this very individual species.

Distribution: Borneo, Philippines, Fiji.

***Calymperes porrectum* Mitten, Linn. Soc. Bot., Jour., 172, 1869.**

Calymperes salakense Bescherelle, Essai Calymp. Sci. Nat., Ann. VIII, Bot. 1: 302, 1895.

Tau: on coconut tree trunk, along Luma-Faleasao trail, alt. 45-50 m., Sept. 30, 1939, *Yuncker 9140*; Faleulu Valley, one eighth mile south of Muao's house, growing on dead logs and fallen wood lying on moist soil under cover of light forest, hillside, gentle slope, 0.75 mile from shore, alt. 213 m., *W. Harris 436a*.

As far as I can see, the Samoan collections agree perfectly with *C. salakense* Bescherelle. Fleischer evidently recognized this agreement (*Laubmoosflora v. Java*, p. 252) but he erroneously cites *C. porrectum* Bescherelle instead of *C. porrectum* Mitten. As Mitten's name was published in 1869, it has unquestioned priority and *C. salakense* Bescherelle should be reduced to synonymy.

Distribution: Malacca, Java, New Guinea, Philippines.

***Calymperes Dozyanum* Mitten, Linn. Soc., Jour., Suppl. 1 (Musc. Ind. Or.): 42, 1859.**

Tau: on coconut tree trunk, along Luma-Faleasao trail, alt. 45-50 m., Sept. 30, 1939, *Yuncker 9140a*.

New to Samoa. Distribution: Ceylon, Sumatra, Java, Borneo, Philippines.

***Calymperes moluccense* Schwaegrichen, Suppl. 2(2): 99, pl. 127, 1824.**

Tau: upper Luma plateau, east of pig fence, along Vaosa trail, plantation in amphitheater plateau, moderate slope, damp, 2-10 feet above ground on trunks of coconut and breadfruit trees, 0.75 mile to shore, alt. 90 m., *W. Harris 444*.

New to Samoa. Distribution: Moluccas, Fiji.

***Calymperes Vreisii* Bescherelle, Essai Calymp. Sci. Nat., Ann. VIII, Bot. 1: 307, 1895.**

Tau: just south of Falemai, on face of cliffs, 6 m. below top, Fitiuta trail, on surface of massive lava boulders, 800 feet from beach, alt. 36 m., *W. Harris 682b*.

New to Samoa. Distribution: Celebes, Marquesas Islands.

***Calymperes tahitense* (Sullivant) Mitten, Linn. Soc. Bot., Jour. 10: 172, 1869.**

Tau: Judds Crater, northwest rim, on trunks and limbs of trees in damp, rather open forest, 0.75 mile to shore, alt. 168 m., *Harris 413a*; Faleulu Valley,

one eighth mile south of Muao's house, 0.75 mile from shore, alt. 204 m., growing on trunks of live trees, 1-8 feet above ground in medium damp, open forest, on gentle slope, foot of mountains, *W. Harris* 423; Faleulu Valley, woodland south of Muao's plantation, 0.75 mile from shore, alt. 204 m., growing on damp lava rock bed of small intermittent stream, on hillside in damp, rather open forest, moderate slope, *W. Harris* 439; upper Luma plantation, east of pig fence, along Vaosa trail, plantation in amphitheater plateau, moderate slope, damp, 2-10 feet above ground, on trunks of coconut and breadfruit trees, 0.75 mile from shore, *W. Harris* 444b.

Distribution: Java, Borneo, Philippines, Fiji, Tahiti.

***Calymperes serratum* A. Braun, in C. Müller, Syn. 1: 527, 1849.**

Tau: Palapala gulch above Palapala Hill, on mountainside, damp native forest, banks of dry stream, on trunks of tree ferns and other trees, 1.5 miles from shore, alt. 270-300 m., *W. Harris* 58.

Distribution: Malaysia, East China, Philippines, New Caledonia, Fiji.

FAMILY POTTIACEAE

Genus **Hyophila** Bridel

***Hyophila involuta* (Hooker) Jaeger, Adumb. 1: 202, 1871-1872.**

Hyophila samoana Mitten, Sam. M., 193, 1871.

Tau: Faleasao, old beach flats, middle section behind Falepopo, growing on tuff at lower edge of talus, under plantation trees, damp, 450 feet from shore, alt. 4-5 m., *W. Harris*.

I have not seen the original of *H. samoana* Mitten, but the present collection is apparently typical and differs in no important particulars from *H. involuta*. I have expressed the thought (Mosses of the Philippines, Philip. Jour. Sci. 68: 114, 1939) that *H. involuta* in a broad sense probably has a wide distribution and an extensive synonymy.

Distribution: India, eastern China, Philippines, and probably wider.

FAMILY DREPANOPHYLLACEAE

Genus **Mniomalia** C. Müller

***Mniomalia semilimbata* (Mitten) C. Müller, Mus. Godeffroy, Jour. 5: 60, 1873-1874.**

Tau: Luma, cliffs east of dispensary, open forest, medium damp, 1,500 feet from shore, alt. 39 m., *W. Harris* 51; Judds Crater, midway between bottom of pit and northwest rim, under forest and plantation cover, on talus below cliff, on rotten log, damp, alt. 135 m., *W. Harris*.

Distribution: Sumatra, Borneo, Philippines, New Guinea.

FAMILY SPIRIDENTACEAE

Genus *Spiridens* Nee*Spiridens aristifolius* Mitten, Linn. Soc. Bot., Jour. 10: 193, 1869.

Tau: on tree in moist forest, west slope of Mt. Vaoaimanu, alt. 450 m., Oct. 5, 1939, *Yuncker 9258*; Faleulu Valley, one eighth mile south of Muao's house, on dead logs and sticks lying on moist soil, gentle slope, hillside, 0.75 mile from shore, alt. 216 m., *W. Harris 114*; Faleulu Draw, damp shallow draw on steep mountainside, in forest, growing on tree trunks 1-8 feet above ground, 1.5 miles from shore, alt. 540 m., *W. Harris*.

Distribution: Fiji.

FAMILY ORTHOTRICHACEAE

Genus *Macromitrium* Bridel*Macromitrium semipellucidum* Dozy and Molkenboer, Sci. Nat., Ann. 2: 311, 1844.

Tau: Palapala, gulch above Palapala Hill, on mountainside, damp native forest, banks of dry stream, on trunks of tree ferns and other trees, 1.5 miles from shore, alt. 270-300 m., *W. Harris 56*; Faleulu Valley, one eighth mile south of Muao's house, growing on trunks of living trees in light forest, damp situation, hillside, gentle slope, 1.5 miles from shore, alt. 210 m., *A. Harris 182*.

Distribution: Sumatra, Borneo, Philippines, New Guinea.

FAMILY PTEROBRYACEAE

Genus *Garovaglia* Endlicher*Garovaglia Powellii* Mitten, Linn. Soc. Bot., Jour. 10: 169, 1869.

Tau: Palapala, gulch above Palapala Hill, on mountainside, damp native forest, banks of dry stream, on trunks of tree ferns and other trees, 1.5 miles from shore, alt. 270-300 m., *W. Harris 54a*; Faleulu Valley, one eighth mile south of Muao's house, growing on trunks of living trees in light forest, damp situation, hillside, gentle slope, one to one eighth mile from shore, alt. 210 m., *A. Harris 181*.

The spines on the back of the leaf are confined to the edges of the plicae and easily overlooked. The best way to view them is to put part of the stem under a cover glass when some of the leaves will show the edges of the folds in profile.

Endemic.

FAMILY METEORIACEAE

Genus *Floribundaria* C. Müller*Floribundaria aeruginosa* (Mitten) Fleischer, Hedwigia 44: 302, 1905.

Tau: trail to peak back of Luma, at side of gulch, on damp rocks, alt. 300 m., *Garber 715a*; Luma cliffs, east of dispensary, on large lava boulders,

partly shaded, medium damp, south side of trail under open forest, 1,500 feet from shore, alt. 39 m., *W. Harris 128.*

Olosega: Olosegauta-Piumafua Ridge, on mossy tree trunk, alt. 510 m., *Garber 1080a.*

Distribution: Fiji, Tahiti.

FAMILY NECKERACEAE

Genus **Himantocladium** (Mitten) Fleischer

Himantocladium implanum (Mitten) Fleischer, Laubmfl. Java 3: 886, 1907.

Tau: Palapala, gulch above Palapala Hill, on mountainside, damp native forest, banks of dry stream, on trunks of tree ferns and other trees, 1.5 miles from shore, alt. 270-300 m., *W. Harris 57.*

Distribution: Fiji, Rapa.

Genus **Pinnatella** (C. Müller) Fleischer

Pinnatella Kuhliana (Bryol. Jav.) Fleischer, Hedwigia 45: 80, 1906.

Tau: Afuli, Fagamalo beach trail, forest cover is strand trees (mostly *fau* and *futu*), moss from lava boulders in talus below shore cliffs, 30 feet from upper beach line, alt. 6 m., *W. Harris.*

Distribution: Sumatra, Java, Celebes, Ceram, New Guinea, Tahiti.

FAMILY HOOKERIACEAE

Genus **Callicostella** (C. Müller) Mitten

Callicostella papillata (Montagne) Mitten, Linn. Soc. Jour., Suppl. 1 (Musc. Ind. Or.): 136, 1859.

Tau: Faleulu Valley, one eighth mile south of Muao's house, growing on dead logs and fallen wood lying on moist soil under cover of light forest, hillside, gentle slope, 0.75 mile from shore, alt. 213 m., *W. Harris 436d.*

Distribution: India, Sumatra, Java, Borneo, Philippines, Pacific islands.

FAMILY THUIDIACEAE

Genus **Pelekium** Mitten

Pelekium velatum Mitten, Linn. Soc. Bot., Jour. 10: 176, 1869.

Tau: Tavalogi Ridge, west end, ridge summit, under light forest cover, moss taken from north face of tuff capping, medium damp, one eighth mile from sea, alt. 143 m., *W. Harris.*

Distribution: Malay Peninsula, Siam, Malaysia, Philippines.

Genus **Thuidium** Bruch and Schimper

Thuidium samoanum Mitten, Linn. Soc. Bot., Jour. 10: 186, 1869.

Tau: Afuli-Fagamalo beach trail, talus on steep hillside below shore cliffs,

strand forest mainly *fau* and *futu* trees, 30 feet from shore, alt. 6 m., *W. Harris* 684a; Afuli-Fagamalo beach trail, forest cover is strand trees (mostly *fau* and *futu*), moss from lava boulders in talus below shore cliffs, 30 feet from upper beach line, alt. 6 m., *W. Harris*; Faleulu Valley, woodland, south of Muao's plantation, 0.75 mile from shore, alt. 204 m., growing on damp lava bed of small intermittent stream, on hillside in damp rather open forest, moderate slope, *W. Harris* 438.

Distribution: Solomon Islands, Fiji.

Thuidium cymbifolium (Dozy and Molkenboer) Bryol. Java, Bryol. Java 2: 115, 1865.

Tau: Palapala stream above Palapala Hill, gulch on north slope of Vaoaimanu Mt., below Olomanu, on rock on stream bank under native forest cover, 1.5 miles to shore, alt. 270-300 m., *W. Harris* 395.

Distribution: Malaysia, Philippines, Formosa, Lord Howe Island, New Guinea, China, Japan.

FAMILY BRACHYTHECIACEAE

Genus **Rhynchostegium** Bruch and Schimper

Rhynchostegium vitianum Bartram and Dixon, B. P. Bishop Mus., Occ. Papers 11(20): 18, 1936.

Tau: Faleulu Valley, woodland south of Muao's plantation, 0.75 mile from shore, alt. 204 m., growing on damp lava rock bed of intermittent stream, on hillside in damp rather open forest, moderate slope, *W. Harris* 437a.

New to Samoa. Distribution: Fiji, Austral Islands.

FAMILY SEMATOPHYLLACEAE

Genus **Acporium** Mitten

Acporium sigmatodontium (C. Müller) Fleischer, Laubmfl. Java 4: 1281, 1918.

Tau: Palapala, gulch above Palapala Hill, on mountainside, damp native forest, banks of dry stream, on trunks of tree ferns and other trees, 1.5 miles from shore, alt. 270-300 m., *W. Harris* 54.

Apart from the relatively shorter setae there are apparently no other characters distinguishing *A. brevisetulum* (C. Müller) Jaeger from *A. sigmatodontium*. The variety *brachypodium* (Dixon) Bartram is a parallel form, so I think that *A. brevisetulum* may be safely dropped.

Distribution: Ceylon, Sumatra, Java, New Guinea, Philippines, Tahiti.

Genus *Taxithelium* Spruce

Taxithelium samoanum (Mitten) Brotherus, Engler and Prantl, Nat. Pflanzenfam. I, 3 : 1091, 1908.

Tau: Faleulu Valley, one eighth mile south of Muao's house, growing on dead logs and fallen wood lying on moist soil under cover of light forest, hillside, gentle slope, 0.75 mile from shore, alt. 213 m., *A. Harris 176a*.

Endemic.

Taxithelium tenuisetum (Sullivant) Mitten, Fl. Vit., 397, 1871.

Tau: Faleulu Valley, one eighth mile south of Muao's house, growing on dead logs and fallen wood lying on moist soil under cover of light forest, hillside, gentle slope, 0.75 mile from shore, alt. 213 m., *W. Harris 436c*.

Ofu: on rotten log, along southeastern shore trail, alt. 5 m., Nov. 26, 1939, *Yuncker 9517*.

Distribution: Fiji, Niue Island.

FAMILY HYPNACEAE

Genus *Ectropothecium* Mitten

Ectropothecium sodale (Sullivant) Mitten, Linn. Soc. Bot., Jour. 10 : 180, 1869.

Tau: Faleulu Valley, one eighth mile south of Muao's house, growing on dead logs and fallen wood lying on moist soil under cover of light forest, hillside, gentle slope, 0.75 mile from shore, alt. 213 m., *W. Harris 156, 436, 403; A. Harris 176*.

Distribution: Fiji, Tahiti.

Ectropothecium incubans (Reinwardt and Hornschuch) Jaeger var. *saberulum* (Brotherus) Fleischer, Laubmfl. Java 4 : 1410, 1922.

Tau: Palapala, gulch above Palapala Hill, on mountainside, damp native forest, banks of dry stream, on trunks of tree ferns and other trees, 1.5 miles from shore, alt. 270-300 feet, *W. Harris 55*.

New to Samoa. Distribution: Java, Fiji.

Ectropothecium adnatum Brotherus, unpublished manuscript.

Tau: Faleulu Valley, Muao's plantation, south of house, in rocky soil under coconut, breadfruit, and banana trees, medium damp soil, hillside, moderate slope, 0.5 mile from shore, alt. 195 m., *W. Harris 375*; Faleulu Valley, one eighth mile south of Muao's house, 0.75 mile from shore, alt. 204 m., growing on trunks of live trees, 1-8 feet above ground, in medium damp open forest, on gentle slope, foot of mountains, *W. Harris 422*; Faleulu Valley, one eighth mile south of Muao's house, growing on trunks of living trees in light forest, damp situation, hillside, gentle slope, one eighth mile from shore, alt. 210 m.,

A. Harris 180; Papatea, South Cove, alt. 1.5-6 m., taken on tuff, damp lower edge of strand forest, 5-30 feet from shore, *W. Harris 407*; on old log in forest south of Siufaga, alt. 300 m., Oct. 4, 1939, *Yuncker 9236, 9244*.

New to Samoa. Distribution: Solomon Islands, Fiji, Niue Island.

Ectropothecium cyathothecium (C. Müller) Brotherus, Engler and Prantl, Nat. Pflanzenfam. I, 3 : 1066, 1908.

Tau: Faleulu Valley, one eighth mile south of Muao's house, on dead logs and sticks lying on moist soil, gentle slope, hillside, 0.75 mile from shore, alt. 216 m., *W. Harris 116*.

Distribution: Fiji.

Ectropothecium malacoblastum (C. Müller) Jaeger, Adumb. 2 : 526, 1877-1878.

Tau: Faleulu Valley, woodland south of Muao's plantation, 0.75 mile from shore, alt. 204 m., growing on damp lava rock bed of small intermittent stream on hillside in damp rather open forest, moderate slope, *W. Harris 437, 138*; Faleulu, Muao's plantation, alt. 168 m., growing on rock wall under cover of plantation, 0.5 mile from shore, *A. Harris 160*; Palapala stream above Palapala Hill, gulch on north slope of Mt. Vaoaimanu below Olomanu, on rock on stream bank under native forest cover, 0.5 mile to shore, alt. 270-300 m., *W. Harris 394*.

Distribution: Fiji.

Genus **Vesicularia** C. Müller

Vesicularia tahitensis (Ångström) Brotherus, Engler and Prantl, Nat. Pflanzenfam. I, 3 : 1094, 1908.

Tau: Judds Crater, northwest rim, spur rim, south exposure, under light forest cover of tree ferns and other trees, cliff face, growing on volcanic cinders, 0.5 mile from shore, alt. 165 m., *W. Harris 1764b*.

New to Samoa. Distribution: Tahiti, Rapa, Pitcairn Island.

Vesicularia inflectens (Bridel) C. Müller, Engl. Bot. Jahrb. 23 : 330, 1892.

Tau: south side of South Cove, Papatea, rocky point, foot of hill, lower edge of strand forest at beach, growing on tuff rock, 5-20 feet from shore, alt. 3-4.5 m., *W. Harris 313*; just south of Falemai, on face of cliffs, 20 feet below top, Fitiuta trail, on surface of massive lava boulders, 800 feet from beach, alt. 36 m., *W. Harris 682c*; Judds Crater, bottom of pit, on lava boulders, under cover of shrubs, fern and mixed plantation and forest trees, 0.5 mile from shore, alt. 114 m., *W. Harris*; Afuli-Fagamalo beach trail, forest cover is strand trees (mostly *fau* and *futu*), moss from lava boulders in talus below shore cliffs, 30 feet from upper beach line, alt. 6 m., *W. Harris*.

Distribution: Hong Kong, Borneo to Rapa and Austral Islands.

PTERIDOPHYTA

PSILOPHYTINAE

FAMILY PSILOTACEAE

Genus *Psilotum* Swartz

Psilotum complanatum Swartz, Syn. Fil., 118, 414, pl. 4, 1806.

Tau: epiphytic on tree fern, in moist ravine, alt. 200 m., Sept. 25, 1939, *Yuncker 9016*; alt. 250 m., Sept. 27, 1939, *Yuncker 9075*; Amouli trail, alt. 180 m., Sept. 16, 1921, *Garber 626*; trail to peak back of Tau, alt. 360 m., Jan. 28, 1922, *Garber 716*; Faleulu Valley, in light forest, alt. 210 m., July 27, 1938, *W. Harris 158*.

LYCOPODINAE

FAMILY LYCOPODIACEAE

Genus *Lycopodium* Linnaeus

Lycopodium carinatum Desvaux, in Poiret, Encycl. Méth., Suppl. 3: 555, 1814.

Tau: pendent from tree branches in damp ravine south of Siufaga, alt. 350 m., Sept. 27, 1939, *Yuncker 9073*.

Lycopodium Phlegmaria Linnaeus, Sp. Pl., ed. 2, 1564, 1763.

Tau: epiphytic on rotten tree trunk in damp ravine, alt. 200 m., Sept. 27, 1939, *Yuncker 9071, 9074*. Native name, *la'au fai pale*.

Lycopodium squarrosum Forster f., Fl. Ins. Austr. Prodr. 86, no. 479, 1786.

Tau: epiphytic on tree branch in damp ravine, alt. 250 m., Sept. 27, 1939, *Yuncker 9072*; on rotten log in forest on west slope of Mt. Vaoaimanu, alt. 450 m., Oct. 5, 1939, *Yuncker 9247*. Native name, *la'u talatala*.

FAMILY SELAGINELLACEAE

Genus *Selaginella* Spring

?***Selaginella Reineckei*** Hieronymus, in Engler and Prantl, Nat. Pflanzenfam. I, 4: 678, 1902.

Tau: in moist, shady ravine on slope south of Siufaga, alt. 150 m., Sept. 27, 1939, *Yuncker 9069*; in deep forest, slope south of Siufaga, alt. 150 m., Oct. 1, 1939, *Yuncker 9159*; in deep, moist forest 2 miles south of Siufaga, alt. 300 m., Oct. 1, 1939, *Yuncker 9151*; in deep moist ravine on south slope of Siufaga, alt. 150 m., Sept. 25, 1939, *Yuncker 9030*; Amouli trail, alt. 165 m., Sept. 16, 1921, *Garber 618*; trail back of Luma, alt. 240 m., Jan. 28, 1922, *Garber 713*.

Ofu: on moist forest floor, main ridge, alt. 250 m., Nov. 27, 1939, *Yuncker 9537*. Native name, *la'au vao*.

Selaginella Whitmeei Baker, Jour. Bot. 23: 24, 1885.

Tau: Powell 38; Amouli trail, alt. 180 m., Sept. 16, 1921, Garber 625.

FILICINAE

FAMILY OPHIOGLOSSACEAE

Genus **Ophioglossum** Linnaeus

Ophioglossum pendulum Linnaeus, Sp. Pl., ed. 2, 2: 1518, 1763.

Tau: epiphyte in moist ravine on mountain slope south of Siufaga, alt. 200 m., Sept. 27, 1939, Yuncker 9078; epiphyte, Amouli trail, alt. 135-150 m., Sept. 16, 1921, Garber 616; 632. Native name, *laugapapa* (also used for *Asplenium nidus* L.).

FAMILY MARATTIACEAE

Genus **Angiopteris** Hoffmann

Angiopteris evecta (Forster) Hoffmann var. **Vaupelii** Hieronymus, Hedwigia 61: 250, 1919.

Tau: plateau back of Faleasao, alt. 180 m., Aug. 4, 1921, Garber 579. Abundant on all islands, Guest. Native name, *oli oli*.

Genus **Marattia** Swartz

Marattia Smithii Mettenius, Kuhn, Ferh. zool. bot. Fes. 19: 584, 1869.

Olosega: top of Piumafua mountain, alt. 600 m., June 17, 1925, Garber 1053.

FAMILY GLEICHENIACEAE

Genus **Dicranopteris** Bernhardi

Dicranopteris linearis (Burman) Underwood, Torrey Bot. Club, Bull. 34: 250, 1907.

Ofu: trailing over edge of cliff, slope back of Ofu village, alt. 240 m., Nov. 25, 1939, Yuncker 9450.

Olosega: Olosegauta-Piumafua ridge, alt. 480 m., June 17, 1925, Garber 1078. Native names, *vaomu*, *vao taumanamana*.

FAMILY SCHIZAEACEAE

Genus **Schizaea** Smith

Schizaea dichotoma (Linnaeus) Smith, Ac. Turin, Mém. 5: 422, pl. 9, fig. 9, 1793.

Tau: in deep forest on western slope of Mt. Vaoaimanu, alt. 300 m., Oct. 5, 1939, Yuncker 9262; Amouli trail, alt. 180 m., Sept. 16, 1921, Garber 626a.

Ofu: on moist forest floor on main ridge, alt. 350 m., Nov. 27, 1939, *Yuncker 9525*; Feaiia Point, alt. 90 m., May 25, 1925, *Garber 974*.

Olosega: Olosegauta-Piumafua ridge, alt. 420 m., June 17, 1925, *Garber 1083*.

FAMILY HYMENOPHYLLACEAE

Genus *Trichomanes* Linnaeus

Trichomanes apiifolium Presl, Hym., 16, 44, 1843.

Tau: on decayed tree trunk, below peak back of Tau, alt. 525 m., Jan. 28, 1922, *Garber 722*; on tree trunk, alt. 630 m., Jan. 28, 1922, *Garber 740*; on the ground, alt. 630 m., Jan. 28, 1922, *Garber 743*; at alt. 600 m., Jan. 28, 1922, *Garber 757*.

Olosega: Olosegauta-Piumafua ridge, alt. 540 m., June 17, 1925, *Garber 1042*.

Trichomanes assimile Mettenius, Kuhn, Linnaea 35: 386, 1868.

Tau: in damp moss on tree trunk, trail to peak back of Tau, alt. 420 m., Jan. 28, 1922, *Garber 718*.

Trichomanes bipunctatum Poiret, in Lamarck, Encycl. Méth. 8: 69, 1808.

Tau: on mossy tree trunk, below peak back of Tau, alt. 630 m., Jan. 28, 1922, *Garber 741*.

Trichomanes Boryanum Kunze, Farrnkr. 1: 237, pl. 97, 1847.

Tau: on wet rocks in shady ravine on slope two miles south of Siufaga, alt. 50 m., Sept. 25, 1939, *Yuncker 9017*; Amouli trail, alt. 165 m., Sept. 16, 1921, *Garber 621*; trail to peak back of Luma, alt. 300 m., Jan. 28, 1922, *Garber 715*.

Trichomanes dentatum van den Bosch, Nederl. Kruidk. Arch. 51: 182, 1861.

Olosega: *Reinecke 66*; on mossy tree trunk, top of Piumafua mountain, alt. 630 m., June 17, 1925, *Garber 1052*; on mossy tree trunk, Piumafua ridge, alt. 570 m., June 17, 1925, *Garber 1057*.

Trichomanes Endlicherianum Presl, Epim. Bot., 10, pl. 5A, 1849-51.

Tau: *Powell 155*.

Ofu: *Reinecke 43d*.

Trichomanes humile Forster f., Fl. Ins. Austr. Prodr., 12, 1786.

Tau: on moist rocks and tree trunks along Fitiuta trail, Sept. 26, 1939, *Yuncker 9057*; on rotten log in moist, shady ravine, alt. 45 m., Sept. 25, 1939, *Yuncker 9001*; on volcanic cinders, northwest rim of Judds Crater, alt. 165 m., June 17, 1937, *W. Harris 1764a*; northwest rim of Judds Crater, on trunks of trees, alt. 170 m., July 27, 1938, *W. Harris*; on face of cliffs just south of Falemai, Fitiuta trail, alt. 36 m., April 3, 1937, *W. Harris 682a*; gulch above Palau-

pala Hill, alt. 270-300 m., July 10, 1938, *W. Harris*; on face of lava cliff behind Luma, alt. 45 m., July 24, 1938, *W. Harris*; on damp mossy rock at base of tree along rocky stream bed, Amouli trail, alt. 180 m., Sept. 16, 1921, *Garber* 628.

Olosega: Olosegauta-Piumafua ridge, on mossy tree trunk, alt. 510 m., June 17, 1925, *Garber* 1081.

Trichomanes intermedium van den Bosch, Nederl. Kruidk. Arch. 5: 179, 1861.

Tau: below peak back of Tau, alt. 585 m., Jan. 28, 1922, *Garber* 729; gulch above Palapala Hill, on tree trunks, alt. 270-300 m., July 10, 1938, *W. Harris*.

Olosega: Olosegauta-Piumafua ridge, alt. 570 m., June 17, 1925, *Garber* 1063.

Trichomanes pallidum Blume, Enum., 225, 1828.

Manua: *Nightingale* (type of *T. glaucofuscum* Hooker, Night. Ocean. Sketch., App. 131, 1835, in Kew).

Trichomanes saxifragoides Presl, Hym. 16, 39, 1843.

Tau: in damp moss on tree fern trunk, slope back of Tau, alt. 645 m., Jan. 28, 1922, *Garber* 744.

Manua: *Powell* 181.

Genus **Hymenophyllum** Smith

Hymenophyllum formosum Brackenridge, U. S. Expl. Exped., Bot. 16: 268, pl. 32, fig. 3, 1854.

Tau: below the peak back of Tau, alt. 570 m., Jan. 28, 1922, *Garber* 759.

Olosega: on mossy tree trunk near top of Piumafua mountain, June 17, 1925, *Garber* 1060.

Hymenophyllum polyanthos Swartz, Jour. Bot. Schrad. 2: 100 (1800) 1801.

Tau: epiphytic, in dense, wet forest on west slope of Mt. Vaoaimanu, alt. 450 m., Oct. 5, 1939, *Yuncker* 9256; on damp tree trunk below peak back of Tau, alt. 585 m., Jan. 28, 1922, *Garber* 727; also at 615 m., *Garber* 737; Faleulu draw, on hillside on trees, alt. 480 m., Sept. 24, 1936, *W. Harris*.

Manua: *Reinecke* 175.

Hymenophyllum praetervisum Christ, Bot. Jahrb. 23: 338, 1896.

Manua: *Reinecke* 88c.

FAMILY CYATHEACEAE

Genus **Alsophila** R. Brown

?**Alsophila extensa** (Forster) R. Brown, Prodr. Fl. Nov. Holl. 1: 158, 1810.

Abundant on all islands as reported by Guest in 1939. Native name, *oli oli*.

FAMILY POLYPODIACEAE

Genus *Dryopteris* Adanson

Dryopteris dissecta (Forster) O. Kuntze, Rev. Gen. 2: 812, 1891.

Tau: on moist bank along Luma-Faleasao trail, alt. 50 m., Sept. 30, 1939, *Yuncker* 9137; in damp, rocky soil, hillside, Faleulu valley, alt. 210 m., July 27, 1938, *W. Harris* 383; on lava rock, open forest, cliffs behind Luma village, alt. 39 m., July 24, 1938, *W. Harris* 48; on lava rock in talus at foot of cliff behind Luma village, alt. 8 m., July 13, 1938, *W. Harris* 347.

Ofu: on rocky cliff, slope back of Ofu village, alt. 30 m., Nov. 25, 1939, *Yuncker* 9478.

Dryopteris gongyloides (Schkuhr) O. Kuntze, Rev. Gen. 2: 811, 1891.

Tau: in low, marshy area, alt. 5 m., Oct. 7, 1939, *Yuncker* 9297.

Dryopteris setigera (Blume) O. Kuntze, Rev. Gen. 2: 813, 1891.

Tau: on old lava outcropping, slope south of Siufaga village, alt. 250 m., Oct. 4, 1939, *Yuncker* 9239; on face of lava cliff, behind Luma village, alt. 45 m., July 24, 1938, *W. Harris* 386.

Ofu: in thicket along southeastern shore trail, alt. 5 m., Nov. 26, 1939, *Yuncker* 9498.

Genus *Tectaria* Cavanilles

Tectaria Stearnsii Maxon, Biol. Soc. Washington, Proc. 36: 175, pl. 17, fig. a, 1923.

Ofu: on dry bank along trail on slope back of Ofu village, alt. 240 m., Nov. 25, 1939, *Yuncker* 9455. Native names, *vao tuaniu*, *la'au fai pale*.

Genus *Athyrium* Roth

Athyrium accedens (Blume) Milde, Bot. Zeit., 353, 1870.

Tau: on lava outcropping on slope 1-1.5 mile south of Siufaga, alt. 240 m., Oct. 4, 1939, *Yuncker* 9237; in coconut, breadfruit and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 147.

Genus *Blechnum* Linnaeus

Blechnum orientale Linnaeus, Sp. Pl., 1077, 1753.

Tau: in open area along trail on slope south of Siufaga, alt. 60 m., Sept. 25, 1939, *Yuncker* 9011.

Ofu: on clay bank along trail on slope back of Ofu village, alt. 240 m., Nov. 25, 1939, *Yuncker* 9454. Native name, *lau magamaga*.

Genus *Asplenium* Linnaeus

Asplenium laserpitiifolium Lamarck, Encycl. Méth. 2: 310, 1786.

Tau: in dark, wet, cavern-like depression in old lava out-cropping, alt. 250 m., Oct. 4, 1939, *Yuncker* 9240; on lava rocks, cliffs behind Luma village, alt. 39 m., July 24, 1938, *W. Harris* 41. Native name, *vao tuaniu*.

***Asplenium Nidus* Linnaeus, Sp. Pl., 1079, 1753.**

Tau: on lava cliff behind Luma, alt. 45 m., July 24, 1938, *Wray Harris* 123.

Ofu: on rocky cliff, slope back of Ofu village, alt. 45 m., Nov. 25, 1939, *Yuncker* 9474. Native name, *launapapa*.

Genus *Lomagramma* J. Smith

***Lomagramma cordipinna* Holttum, Gard. Bull. Straits Settlem. 9(2): 202, 1937.**

Tau: in damp forest, Faleulu valley, alt. 210 m., July 27, 1938, *W. Harris* 209.

Genus *Nephrolepis* Schott

***Nephrolepis biserrata* (Swartz) Schott, Gen. Fil., ad. pl. 3, 1834.**

Tau: epiphytic on tree in thicket about two miles south of Siufaga, alt. 150 m., Oct. 1, 1939, *Yuncker* 9156.

***Nephrolepis hirsutula* (Forster) Presl, Tent. Pterid., 79, 1836.**

Tau: on moist shady bank along Fitiuta trail, Sept. 26, 1939, *Yuncker* 9044; in moist, rocky ravine on slope about two miles south of Siufaga, alt. 50 m., Sept. 25, 1939, *Yuncker* 9019; on lava rock, cliffs back of Luma village, alt. 39 m., July 24, 1938, *W. Harris* 127; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 146.

Genus *Microlepia* Presl

?***Microlepia* species.**

Tau: on lava cliff, behind Luma village, alt. 45 m., July 24, 1938, *W. Harris* 384.

Genus *Lindsaya* Dryander

***Lindsaya decomposita* Willdenow, Sp. Pl. 5: 425, 1810.**

Tau: in rich soil of wet, dense forest on west slope of Mt. Vaoaimanu, alt. 500 m., Oct. 5, 1939, *Yuncker* 9259, 9257; epiphytic on tree in wet forest on slope about two miles south of Siufaga, alt. 300 m., Oct. 1, 1939, *Yuncker* 9152.

Genus *Pteris* Linnaeus

***Pteris comans* Forster, Fl. Ins. Austr. Prodr., 79, 1786.**

Tau: in low wet area south of Siufaga, alt. 3 m., Sept. 29, 1939, *Yuncker* 9093; south side of South Cove, Papatea, on tuff, alt. 3-5 m., July 1938, *W. Harris* 315.

***Pteris ensiformis* Burman f., Fl. Ind., 230, 1768.**

Tau: on face of lava cliff, back of Luma village, alt. 45 m., July 24, 1938, *W. Harris* 28.

Ofu: on rocky cliff, slope back of Ofu village, alt. 45 m., Nov. 25, 1939, *Yuncker* 9479. Native name, *lau manamana*.

Pteris tripartita Swartz, Jour. Bot. Schrad. 2:67 (1800) 1801.

Tau: in coconut plantation, Faleulu valley, alt. 16 m., July 27, 1938, *A. Harris* 144.

Ofu: *Garber* 998; in thicket along southeastern shore trail, alt. 5 m., Nov. 26, 1939, *Yuncker* 9501. Native names, *oli oli, lagasese*.

Genus **Acrostichum** Linnaeus

Acrostichum aureum Linnaeus, Sp. Pl., 1069, 1753.

Tau: within a few feet of the sea on a rocky point in Luma, Nov. 5, 1922, *Garber* 782.

Ofu: in soil pocket in lava rocks near the sea along the northwestern shore, alt. 5 m., Nov. 28, 1939, *Yuncker* 9561. Native names, *lau taputa, sa'ato*.

Genus **Humata** Cavanilles

Humata heterophylla (Smith) Desvaux, Prodr. fam. foug., 323, 1825.

Tau: epiphytic on trees in shady, damp ravine, south of Siufaga, alt. 150 m., Sept. 27, 1939, *Yuncker* 9068. Native name, *vao sosolo*.

Genus **Davallia** Smith

Davallia solida (Forster) Swartz, Jour. Bot. Schrad. 2:87 (1800) 1801.

Tau: clinging to tree along Fitiuta trail, Sept. 26, 1939, *Yuncker* 9050.

Ofu: clinging to coconut trees, slope behind Ofu village, alt. 150 m., Nov. 25, 1939, *Yuncker* 9456, 9457. Native name, *vao sosolo i le niu*.

Genus **Arthropteris** J. Smith

Arthropteris oblitterata (R. Brown) J. Smith, Cat. Cult. Ferns, 62, 1827.

Tau: clinging on trees and rocks, alt. 5 m., Oct. 7, 1939, *Yuncker* 9292; open woodland, on trunks of living trees, alt. 210 m., July 27, 1938, *A. Harris* 388; on damp mossy rocks, in bed of intermittent stream, Faleulu Valley, alt. 204 m., July 27, 1938, *W. Harris* 231; on lava rock, talus at foot of cliff behind Luma village, alt. 8 m., July 13, 1938, *W. Harris* 348; on lava cliff behind Luma village, alt. 45 m., July 24, 1938, *W. Harris* 44.

Genus **Cyclophorus** Desvaux

Cyclophorus lanceolatus (Linnaeus) Alston, Jour. Bot. 69:102, 1931.

Tau: epiphytic on trees along Fitiuta trail, Sept. 26, 1939, *Yuncker* 9055.

Ofu: clinging to coconut tree, slope back of Ofu village, alt. 150 m., Nov. 25, 1939, *Yuncker* 9458.

Genus **Polypodium** Linnaeus

Polypodium blechnoides (Greville) Hooker, Sp. Fil. 4:180, 1862.

Tau: epiphytic on trees in moist ravine on slope south of Siufaga, alt. 150 m., Sept. 27, 1939, *Yuncker* 9077.

Polypodium Scolopendria Burman f., Fl. Ind., 232, 1768.

Tau: terrestrial and epiphytic, near shore, alt. 5 m., Sept. 25, 1939, *Yuncker 9023*; on lava cliff behind Luma, alt. 45 m., July 24, 1938, *W. Harris 126*.

Ofu: on trees and rocks, slope back of Ofu village, alt. 20 m., Nov. 25, 1939, *Yuncker 9471*. Native name, *fiso vao*.

Polypodium vitiense Baker, Jour. Bot. 17: 298, 1879.

Tau: on old lava rock wall, slope south of Siufaga, alt. 150 m., Oct. 4, 1939, *Yuncker 9238*.

Genus ***Prosaptia*** Presl***Prosaptia contigua*** (Forster) Presl, Tent. Pterid., 166, 1836.

Tau: epiphytic on trees in dense, wet forest on west slope of Mt. Vaoai-manu, alt. 450 m., Oct. 5, 1939, *Yuncker 9260*.

Genus ***Antrophyum*** Kaulfuss***Antrophyum alatum*** Brackenridge, U. S. Expl. Exped., Bot. 16: 64, 1854.

Tau: on rotten log in thicket near Siufaga, alt. 5 m., Oct. 7, 1939, *Yuncker 9290*; epiphytic on trees in moist ravine along Fitiuta trail, Sept. 26, 1939, *Yuncker 9056*; in damp, rocky soil under light forest cover, hillside, Faleulu valley, alt. 210 m., July 27, 1938, *W. Harris 262, 382X*.

Antrophyum angustatum Brackenridge, U. S. Expl. Exped., Bot. 16: 63, 1854.

Ofu: on rocks in damp forest on central ridge, alt. 300 m., Nov. 27, 1939, *Yuncker 9539*. Native name, *lau talatalo*.

Genus ***Vaginularia*** Fée***Vaginularia paradoxa*** (Fée) Mettenius, Mus. Bot. Lugd.-Bat., Ann. 4: 174, 1868-69.

Tau: epiphytic on trees in moist ravine south of Siufaga, alt. 150 m., Sept. 27, 1939, *Yuncker 9070*; on damp, dead wood, in woodland, Faleulu valley, alt. 210 m., July 27, 1938, *W. Harris 405*; on trunks of living trees, light forest, Faleulu valley, alt. 210 m., July 27, 1938, *A. Harris 188*; on lava rock, cliffs back of Luma village, alt. 39 m., July 24, 1938, *W. Harris 133*.

SPERMATOPHYTA

MONOCOTYLEDONEAE

FAMILY PANDANACEAE

Genus ***Freycinetia*** Gaudichaud***Freycinetia Hombronii*** Martelli, in Rechinger, Wiss. Forschungsreise nach Samoa, 56, 1910.

Tau: on lava boulder in talus at foot of cliff, cliffs behind Luma village,

July 13, 1938, *W. Harris* 259. This specimen is sterile but it agrees well with specimens of this species in the Bishop Museum herbarium.

Freycinetia Reinecke Warburg, in Reinecke, Fl. Samoa-Inseln, 578, pl. 8, fig. B, 1898.

Tau: climbing in trees in ravine on slope south of Siufaga, alt. 200 m., Sept. 27, 1939, *Yuncker* 9092; near top of peak back of Tau, alt. 660 m., fruit, Jan. 28, 1922, *Garber* 750. Native names, 'ie'ie, lau 'ie'ie.²

Genus *Pandanus* Linnaeus

Pandanus calostigma Martelli f. *samoana* Martelli, B. P. Bishop Mus., Occ. Papers 10(13) : 9, pl. 2, figs. 1-2, 1934.

Ofu: in old coconut grove, old village site of Mafafa, alt. 4-5 m., May 26, 1925, *Garber* 984.

Pandanus tauensis Martelli, B. P. Bishop Mus., Occ. Papers 10(13) : 14, pls. 3, 4, 9, 1934.

Tau: on sandy beach at Faga, alt. 3-4 m., Sept. 11, 1921, *Garber* 665.

Guest reports it as scarce on Ofu. Native name, lau paogo.

Pandanus tectorius Parkinson var. *laevis* (Kunth) Warburg, Pflanzenr. 3 (IV.9) : 48, 1900.

Tau: near dwelling, Luma, cultivated, Dec. 15, 1921, *Garber* 662 (apparently this is the specimen cited by Martelli as *Garber* 665). Native names, 'ie toga, lau fala.

FAMILY GRAMINEAE

Genus *Schizostachyum* Nees

Schizostachyum glaucifolium (Ruprecht) Munro, Linn. Soc., Trans. 26 : 137, 1868.

Tau: small clump near the trail on west slope of Mt. Vaoaimanu, alt. 120 m., Oct. 5, 1939, *Yuncker* 9268.

Ofu: small clump in ridge forest above Ofu village, alt. 360 m., Nov. 27, 1939, *Yuncker* 9521.

Olosega: Piumafua Ridge, alt. 500 m., June 17, 1925, *Garber* 1041. Native names, 'ofe, 'ofe samoae.

Genus *Eragrostis* Beauvois

Eragrostis amabilis (Linnaeus) Wight and Arnott, in Hooker and Arnott, Bot. Beechey Voy., 251, 1841.

Tau: on old coral rock about dwellings, Oct. 2, 1939, *Yuncker* 9169; Luma, near swampy area, Aug. 5, 1938, *W. Harris* 3; Luma, in sandy soil near the beach, July 20, 1938, *W. Harris* 339. Native name, fali funa.

² Lau (leaf) is now often used before the specific name of a plant.

Genus **Centotheca** Desvaux

Centotheca latifolia (Linnaeus) Triniius, Fund. Agrost., 141, 1820.

Tau: in marshy area south of Siufaga, alt. 3 m., Sept. 29, 1939, *Yuncker 9100*; along Luma-Faleasao trail, alt. 50 m., Sept. 30, 1939, *Yuncker 9135*; Sani Ridge, back of Fitiuta, alt. 125 m., Aug. 9, 1921, *Garber 589*; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris 120*; Faleulu Valley, open woods, alt. 204 m., July 27, 1938, *W. Harris 213, 220*.

Olosega: trail back of Olosegauta, alt. 300 m., June 17, 1925, *Garber 1035*. Native names, *sefa, vao fali, lau 'ofe'ofe*.

Genus **Lepturus** R. Brown

Lepturus repens (Forster) R. Brown, Prodr. Fl. Nov. Holl., 207, 1810.

Ofu: in soil pockets, basaltic rocks along northwestern shore, alt. 5 m., Nov. 28, 1939, *Yuncker 9548*.

Genus **Eleusine** Gaertner

Eleusine indica (Linnaeus) Gaertner, Fruct. et Semin. 1: 8, 1788.

Tau: in sandy soil near the beach, Sept. 29, 1939, *Yuncker 9115*; along Luma-Faleasao trail, alt. 50 m., Sept. 30, 1939, *Yuncker 9136*; sandy soil along trail behind Luma, Aug. 5, 1938, *W. Harris 2*; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1939, *W. Harris 251*; on calcareous beach, Papatea, July 15, 1938, *A. Harris 72*.

Ofu: in sandy soil near the beach, Nov. 25, 1939, *Yuncker 9487*; in sandy soil along southeastern shore trail, alt. 5 m., Nov. 26, 1939, *Yuncker 9499*. Native names, *sefa, ta'ata'a*.

Genus **Cynodon** L. Richard

Cynodon Dactylon (Linnaeus) Persoon, Syn. Pl. 1: 85, 1805.

Tau: under coconut trees in loose, medium-dry beach sand, Luma, July 20, 1938, *W. Harris 336*.

Genus **Digitaria** Heister

Digitaria chinensis Hornemann, Hort. Hafn., Suppl., 8, 1819.

Tau: along Luma-Faleasao trail, alt. 50 m., Sept. 30, 1939, *Yuncker 9125*; Luma, on lava cliff, alt. 45 m., July 24, 1938, *W. Harris 30*; in talus, foot of cliff, alt. 7-8 m., July 13, 1938, *W. Harris 353*; south side of South Cove, Papatea, on tuff at beach, July 15, 1938, *W. Harris 317*.

Ofu: in sandy soil near the beach, alt. 2 m., Nov. 25, 1939, *Yuncker 9488*.

Digitaria Henryi Rendle, Linn. Soc. Bot., Jour. 36: 323, 1904.

Tau: Papatea, in sand near the beach, July 15, 1938, *W. Harris 71*.

Ofu: along rocky shore, northwestern side of island, alt. 5 m., Nov. 28, 1939, *Yuncker 9550*.

Genus **Stenotaphrum** Trinius

Stenotaphrum subulatum Trinius, Acad. Pétersb., Mém. VI, 3(2) : 190, 1835.
Tau: in low marshy area south of Siufaga, Sept. 29, 1939, *Yuncker* 9109.

Genus **Paspalum** Linnaeus

Paspalum conjugatum Bergius, Helvet. Phys. Math., Act. 7 : 129, pl. 8, 1762.
Tau: weed in banana plantation on mountain slope south of Siufaga, alt. 300 m., Oct. 4, 1939, *Yuncker* 9233; Papatea, sandy soil near the beach, July 15, 1938, *W. Harris* 95; Luma, in talus, foot of cliff, alt. 7-8 m., July 13, 1938, *W. Harris* 354.

Ofu: in sandy soil near the beach, Nov. 25, 1939, *Yuncker* 9489. Native names, *vailima*, *vailima matafao*.

Paspalum vaginatum Swartz, Prodr. Veg. Ind. Occ., 21, 1788.

Tau: Luma, in sandy soil near the beach, July 20, 1938, *W. Harris* 305; Papatea, in sandy soil near the beach, July 15, 1938, *A. Harris* 73.

Genus **Panicum** Linnaeus

Panicum ambiguum Trinius, Acad. Pétersb., Mém. VI, 3(2) : 243, 1835.
Tau: in sandy soil near the beach south of Siufaga, Sept. 29, 1939, *Yuncker* 9114; sandy soil, Luma, July 20, 1938, *W. Harris* 334.

Ofu: in sandy soil near the beach back of Ofu village, Nov. 25, 1939, *Yuncker* 9486.

Panicum pilipes Nees and Arnott, Hook. Jour. Bot. Kew Miscel. 2 : 97, 1850.

Tau: Faleulu, weed in coconut plantation, alt. 180 m., July 27, 1938, *W. Harris* 166.

Ofu: plantation, Feaiia Point, alt. 90 m., May 25, 1925, *Garber* 975.

Genus **Oplismenus** Beauvois

Oplismenus compositus (Linnaeus) Beauvois, Ess. Agrost., 54, 1812.
Tau: Luma, on lava cliff, alt. 45 m., July 24, 1938, *W. Harris* 31.

Oplismenus undulatifolius (Arduino) Roemer and Schultes var. **imbecillus** (R. Brown) Hackel, Govern. Labor. Manila, Pub. 35 : 81, 1906.

Ofu: in sandy soil along the southeastern shore trail, alt. 5 m., Nov. 26, 1939, *Yuncker* 9501A.

Genus **Pennisetum** L. Richard

Pennisetum macrostachyum (Brongniart) Trinius, Acad. Pétersb., Mém. VI, 3(2) : 177, 1835.

Tau: cultivated ornamental, Siufaga, Oct. 3, 1939, *Yuncker* 9214.

Genus **Cenchrus** Linnaeus

Cenchrus echinatus Linnaeus, Sp. Pl., 1050, 1753.

Tau: in sandy soil near the shore, Sept. 29, 1939, *Yuncker* 9110; on cal-

careous beach, Papatea, July 15, 1938, *A. Harris* 70; under coconut trees in loose, medium-dry beach sand, Luma, July 20, 1938, *W. Harris* 307, 335.

Ofu: in sandy soil near the beach back of Ofu village, Nov. 25, 1939, *Yuncker* 9485. Native names, *vao tuitui*, *vao papalagi*.

Genus *Thuarea* Persoon

Thuarea involuta (Forster) Roemer and Schultes, Syst. 2: 808, 1817.

Ofu: in sandy soil near the beach, Nov. 25, 1939, *Yuncker* 9491.

Genus *Misanthus* Andersson

Misanthus japonicus (Thunberg) Andersson, Oefvers. Vet. Akad. Förh. 12: 166, 1856.

Tau: along trail, Luma, January 11, 1922, *Garber* 705. Native names, *u*, *fiso*.

Genus *Saccharum* Linnaeus

Saccharum officinarum Linnaeus, Sp. Pl., 54, 1753.

Tau: small patch noted near Luma, but not collected (Yuncker). Native name, *tolo*.

Genus *Coix* Linnaeus

Coix lachryma-jobi Linnaeus, Sp. Pl., 972, 1753.

Tau: in marshy area near Luma, Oct. 6, 1939, *Yuncker* 9283. Native name, *sagisagi*.

FAMILY CYPERACEAE

Genus *Cyperus* Linnaeus

Cyperus brevifolius (Rottboell) Hasskarl, Cat. Hort. Bogor. Alt., 24, 1844.

Tau: weed in taro patch, Luma, Oct. 6, 1939, *Yuncker* 9276; in moist soil on plateau back of Siufaga, alt. 45 m., Oct. 2, 1939, *Yuncker* 9186; on lava cliff behind Luma, alt. 45 m., July 24, 1938, *W. Harris* 91. Native name, *tuse*.

Cyperus compressus Linnaeus, Sp. Pl., 46, 1753.

Tau: plateau back of Luma, alt. 100 m., Dec. 15, 1921, *Garber* 659; in sand along trail behind Luma, Aug. 5, 1938, *W. Harris* 4.

Ofu: in sandy soil near the beach, Nov. 28, 1939, *Yuncker* 9546.

Cyperus cyperoides (Linnaeus) O. Kuntze, Rev. Gen. 3(2): 333, 1898.

Tau: along the Luma-Faleasao trail, alt. 50 m., Sept. 20, 1939, *Yuncker* 9132; plateau back of Luma, alt. 125 m., Dec. 15, 1921, *Garber* 647. Native name, *ta'ata'a*.

Cyperus cyperoides (Linnaeus) O. Kuntze subsp. ***cyperinus*** (Retzius) Kü-kenthal, Engl. Bot. Jahrb. 59: 46, 1924.

Tau: plateau back of Luma, alt. 125 m., Dec. 15, 1921, *Garber* 649; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938,

W. Harris 121; on lava cliffs behind Luma, alt. 45 m., July 24, 1938, *W. Harris* 32.

Cyperus javanicus Houttuyn, Nat. Hist. II, 13: Aawyz. Plaat. [1], pl. 88, fig. 1, 1782.

Tau: in low marshy area, Oct. 6, 1939, *Yuncker* 9285.

Ofu: on talus slope along the northwest shore, Nov. 28, 1939, *Yuncker* 9549; in low ground near the beach, Nov. 25, 1939, *Yuncker* 9484. Native name, *selesele*.

Cyperus Kyllingia Endlicher, Cat. Hort. Acad. Vindob. 1: 94, 1842.

Tau: south side of South Cove, Papatea, on tuff rock at lower edge of strand forest near the shore, July 15, 1938, *W. Harris* 318.

Cyperus Seemannianus Boeckeler var. *altus* Kükenthal, B. P. Bishop Mus., Bull. 128: 18, 1935.

Tau: Faleulu Valley, rocky soil on hillside, alt. 195 m., July 27, 1938, *W. Harris* 199.

Cyperus Seemannianus Boeckeler var. *monostachys* (Boeckeler) Kükenthal, B. P. Bishop Mus., Bull. 128: 18, 1935.

Tau: Sani Ridge back of Fitiuta, alt. 125 m., Sept. 8, 1921, *Garber* 590; plateau back of Luma, alt. 150 m., Dec. 15, 1921, *Garber* 657.

Cyperus Seemannianus Boeckeler aff. var. *perdensus* Kükenthal, B. P. Bishop Mus., Bull. 128: 18, fig. 2, 1935.

Tau: south side of South Cove, Papatea, on tuff at lower edge of strand forest, July 15, 1938, *W. Harris* 319.

Genus *Eleocharis* R. Brown

Eleocharis dulcis (Burman f.) Triniius, ex Henschel, Vita Rumph., 186, 1833.

Tau: in low marshy area near Siufaga, Oct. 7, 1939, *Yuncker* 9296. Native name, *utuutu*.

Genus *Fimbristylis* Vahl

Fimbristylis annua (Allioni) Roemer and Schultes, Syst. 2: 95, 1817.

Tau: in low, marshy area south of Siufaga, Sept. 29, 1939, *Yuncker* 9103; in swamp near Siufaga, Oct. 7, 1939, *Yuncker* 9295.

Fimbristylis cymosa R. Brown var. *pycnocephala* (Hillebrand) Kükenthal, B. P. Bishop Mus., Bull. 128: 20, 1935.

Tau: lava rock along the sea, Fitiuta, Aug. 10, 1921, *Garber* 608.

Ofu: on spray-swept rocks near the sea, Nov. 28, 1939, *Yuncker* 9557. Native name, *mutia*.

Fimbristylis diphylla (Retzius) Vahl, Enum. 2: 289, 1806.

Tau: plateau back of Luma, alt. 125 m., Dec. 15, 1921, *Garber* 648.

Genus **Rynchospora** Vahl

Rynchospora corymbosa (Linnaeus) Britton, New York Acad. Sci., Trans. 11 : 84, 1892.

Tau: in low marshy area, Oct. 6, 1939, *Yuncker* 9286; in low marshy area south of Siufaga, Sept. 29, 1939, *Yuncker* 9102.

Olosega: taro swamp, back of Olosega village, June 23, 1925, *Garber* 1090. Native names, *lau selesele*, *vao selesele*.

Genus **Mapania** Aublet

Mapania longifolia (Boeckeler) C. B. Clarke, Kew Bull., add. ser. 8 : 54, 1908.

Tau: below peak back of Tau, alt. 600 m., Jan. 28, 1922, *Garber* 739. Native name, *lau fala vao*.

Mapania macrocephala (Gaudichaud) K. Schumann, Engl. Bot. Jahrb. 13 : 265, 1891.

Tau: in moist ravine south of Siufaga, alt. 100 m., Sept. 27, 1939, *Yuncker* 9084.

Genus **Scleria** Bergius

Scleria lithosperma (Linnaeus) Swartz, Prodr. Veg. Ind. Occ., 18, 1788.

Tau: in moist shady ravine on slope about two miles south of Siufaga, alt. 150 m., Sept. 27, 1939, *Yuncker* 9080; Luma-Faleasao trail, alt. 40 m., Jan. 8, 1922, *Garber* 690; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 214.

Ofu: along the ridge trail back of Ofu village, alt. 150 m., Nov. 25, 1939, *Yuncker* 9470. Native name, *popo vao*.

Scleria scrobiculata Nees, Wight, Contrib., 117, 1834.

Tau: in low marshy area south of Siufaga, Sept. 29, 1939, *Yuncker* 9106; along the trail near the shore south of Siufaga, Sept. 25, 1939, *Yuncker* 9004; plateau back of Luma, alt. 100 m., Dec. 15, 1921, *Garber* 644.

Ofu: on rocky slope behind Ofu village, alt. 180 m., Nov. 25, 1939, *Yuncker* 9465. Native name, *tafa tolu*.

FAMILY PALMAE

Genus **Eupritchardia** O. Kuntze

Eupritchardia pacifica (Seemann and Wendland) O. Kuntze, Rev. Gen. 3(2) : 323, 1898.

Ofu: cultivated, Aloafao, alt. 8 m., June 6, 1925, *Garber* 1012. Native name, *niu piu*.

Genus **Cocos** Linnaeus

Cocos nucifera Linnaeus, Sp. Pl., 1188, 1753.

Tau: cultivated, Luma, Jan. 10, 1922, *Garber* 703. Extensively cultivated in several varieties on all islands (Yuncker). Native name, *niu*.

FAMILY ARACEAE

Genus *Epipremnum* Schott

?*Epipremnum pinnatum* (Linnaeus) Engler, Pflanzenr. 37(IV.23 Ba): 60, fig. 25, 1908.

Tau: climbing in trees in thicket near the shore south of Siufaga (specimen sterile), Oct. 7, 1939, Yuncker 9288; on the ground and scrambling over rocks, Fagamalo beach, Jan. 7, 1922, Garber 677. Native name, *lau manamana*.

Genus *Amorphophallus* Blume

Amorphophallus campanulatus (Roxburgh) Blume, Mus. Hist. Nat. Paris, Ann. 3: 366, 1834.

Tau: in bush near Siufaga, alt. 3 m., Sept. 21, 1922, Garber 774. Native name, *teve*.

Genus *Colocasia* Schott

Colocasia esculenta (Linnaeus) Schott, Melet. Bot. 1: 18, 1832.

Ofu: cultivated, Ofu village, alt. 8 m., June 22, 1925, Garber 1100; cultivated on all the islands in a number of varieties (Yuncker). Native name, *talo*.

FAMILY BROMELIACEAE

Genus *Ananas* Adanson

Ananas comosus (Linnaeus) Merrill, Interpret. Herb. Amboin., 133, 1917.

Large plantations on Tau, Guest. Native name, *fala 'aina*.

FAMILY COMMELINACEAE

Genus *Commelina* Plumier

Commelina diffusa Burman f., Fl. Ind., 18, pl. 7, fig. 2, 1768.

Tau: in moist soil, along trail on west slope of Mt. Vaoaimanu, alt. 300 m., Oct. 5, 1939, Yuncker 9273.

Olosega: wet talo plantation, June 23, 1925, Garber 1088. Native name, *ma'u'u toga*.

Genus *Aneilema* R. Brown

Aneilema vitiense Seemann, Fl. Vit., 314, pl. 96, 1868.

Tau: along the trail in moist soil, west slope of Mt. Vaoaimanu, Oct. 5, 1939, Yuncker 9274; on old lava wall, south of Siufaga, alt. 300 m., Oct. 4, 1939, Yuncker 9230; Sani Ridge back of Fitiuta, alt. 100 m., Aug. 9, 1921, Garber 588. Native name, *ma'u'u toga*.

FAMILY LILIACEAE

Genus *Cordyline* Commerson

Cordyline terminalis (Linnaeus) Kunth, Berl. Akad., Abh. 30, 1820.

Tau: in thicket along the trail near Fitiuta, alt. 50 m., Sept. 26, 1939, Yuncker 9047; along the Luma-Faleasao trail, alt. 50 m., Sept. 30, 1939,

Yuncker 9128; Fitiuta trail, alt. 80 m., Aug. 4, 1921, *Garber* 554; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 136. Native name, *ti*.

Genus **Asparagus** Linnaeus

Asparagus plumosus Baker, Linn. Soc. Bot., Jour. 14: 613, 1875.

Tau: cultivated ornamental, Siufaga, Oct. 3, 1939, *Yuncker* 9221. Native name, *lau lisi*.

FAMILY AMARYLLIDACEAE

Genus **Zephyranthes** Herbert

Zephyranthes rosea (Sprengel) Lindley, Bot. Reg. 10: 821, 1824.

Tau: cultivated ornamental, Siufaga, Oct. 3, 1939, *Yuncker* 9220. Also observed occasionally as an escape.

Genus **Crinum** Linnaeus

Crinum asiaticum Linnaeus, Sp. Pl., 292, 1753.

Tau: cultivated ornamental, Siufaga, Oct. 3, 1939, *Yuncker* 9219. Native name, *lau talatalo*.

FAMILY TACCACEAE

Genus **Tacca** Forster

Tacca pinnatifida Forster, Charact. Gen., Pl. 70, pl. 35, 1776.

Tau: in sandy soil near the shore, Sept. 25, 1939, *Yuncker* 9027; plateau back of Luma, alt. 125 m., Dec. 15, 1921, *Garber* 653.

Ofu: in sandy soil near the shore, along southeastern trail, Nov. 26, 1939, *Yuncker* 9505. Native name, *masoa, pia*.

FAMILY DIOSCOREACEAE

Genus **Dioscorea** Plumier

Dioscorea bulbifera Linnaeus, Sp. Pl., 1033, 1753.

Tau: climbing in thicket on plateau back of Siufaga, alt. 45 m., Oct. 2, 1939, *Yuncker* 9181.

Ofu: climbing in thicket on main ridge, alt. 300 m., Nov. 27, 1939, *Yuncker* 9526. Native name, *soi*.

Dioscorea pentaphylla Linnaeus, Sp. Pl. 1032, 1753.

Tau: climbing in thicket two miles south of Siufaga, alt. 150 m., Oct. 1, 1939, *Yuncker* 9155. Native name, *pilita*.

FAMILY MUSACEAE

Genus **Musa** Linnaeus

Musa paradisiaca Linnaeus, Sp. Pl., 1043, 1753.

The banana is extensively cultivated on all of the islands in a number of varieties as an important item of food. Native name, *fa'i*.

FAMILY ZINGIBERACEAE

Genus *Alpinia* Roxburgh

Alpinia purpurata (Vieillard) K. Schumann, Pflanzenr. 20(IV.46): 323, 1904.

Tau: on plateau near Siufaga, alt. 45 m., Oct. 2, 1939, *Yuncker 9177*.

Ofu: weed in coconut plantation near Ofu village, alt. 100 m., Nov. 25, 1939, *Yuncker 9460*. Native name, *teuila*.

Genus *Curcuma* Linnaeus

?*Curcuma longa* Linnaeus, Sp. Pl., 2, 1753.

Tau: near dwelling, Siufaga, Oct. 3, 1939, *Yuncker 9218*.

Ofu: Aloafao, alt. 8 m., June 6, 1921, *Garber 1009*. Native name, *ago*.

Genus *Zingiber* Adanson

Zingiber Zerumbet (Linnaeus) Smith, Exot. Bot. 2: 103, pl. 112, 1804.

Tau: Luma-Faleasao trail, alt. 40 m., Jan. 8, 1922, *Garber 689*.

Ofu: along ridge trail above Ofu village, alt. 300 m., Nov. 27, 1939, *Yuncker 9544*. Native name, 'avapui'.

FAMILY CANNACEAE

Genus *Canna* Linnaeus

Canna humilis Bouché, Linnaea 8: 153, 1833.

Tau: plateau back of Siufaga, alt. 100 m., Sept. 27, 1922, *Garber 781*; margin of wet taro patch near Siufaga, Oct. 6, 1939, *Yuncker 9282*. Native name, *fagamanu*.

FAMILY ORCHIDACEAE

Genus *Habenaria* Willdenow

Habenaria tradescantifolia Reichenbach f., in Seemann, Fl. Vit., 293, 1868.

Tau: terrestrial, in dense wet forest on slope two miles south of Siufaga, alt. 300 m., Oct. 1, 1939, *Yuncker 9150*; in deep, shady ravine about one mile south of Siufaga, alt. 50 m., Sept. 27, 1939, *Yuncker 9000A*.

Genus *Platylepis* Blume

Platylepis heteromorpha Reichenbach f., Otia Bot. Hamb. 1: 52, 1878.

Olosega: terrestrial herb, Olosegauta-Piumafua ridge, alt. 550 m., June 17, 1925, *Garber 1073*.

Genus *Erythrodes* Blume

Erythrodes species.

Tau: terrestrial herb, in dense rain forest on west slope of Mt. Vaoaimanu, alt. 500 m., Oct. 5, 1939, *Yuncker 9269*.

Genus **Cheirostylis** Blume**Cheirostylis** species.

Tau: terrestrial herb, in dense rain forest on western slope of Mt. Vaoaimanu, alt. 450 m., Oct. 5, 1939, *Yuncker 9265*.

Genus **Vrydagzynea** Blume**Vrydagzynea** species.

Tau: herb, on mossy rock, Amouli trail, alt. 175 m., Sept. 16, 1921, *Garber 630*.

Genus **Malaxis** Solander**Malaxis resupinata** (Forster) O. Kuntze, Rev. Gen., 673, 1891.

Ofu: in rich soil, forest floor on ridge back of Ofu village, alt. 400 m., Nov. 27, 1939, *Yuncker 9522*.

Genus **Microstylis** Nuttall**Microstylis** species.

Tau: epiphyte, top of peak back of Tau, alt. 700 m., Jan. 28, 1922, *Garber 753*.

Genus **Liparis** L. C. Richard**Liparis** species.

Tau: terrestrial herb, in dense rain forest on west slope of Mt. Vaoaimanu, alt. 500 m., Oct. 5, 1939, *Yuncker 9266*.

Genus **Agrostophyllum** Blume**Agrostophyllum megalurum** Reichenbach f., in Seemann, Fl. Vit., 296, 1868.

Tau: epiphyte, in dense rain forest on west slope of Mt. Vaoaimanu, alt. 450 m., Oct. 5, 1939, *Yuncker 9246*.

Genus **Appendicula** Blume**Appendicula bracteosa** Reichenbach f., in Seemann, Fl. Vit., 299, 1868.

Tau: epiphyte, in dense rain forest on west slope of Mt. Vaoaimanu, alt. 500 m., Oct. 5, 1939, *Yuncker 9267*; epiphyte, on damp tree trunk, below peak back of Tau, alt. 600 m., Jan. 28, 1922, *Garber 730*; epiphyte on trunk of tree fern in gulch, trail to peak back of Luma, alt. 300 m., Jan. 28, 1922, *Garber 714*.

Genus **Glomera** Blume**Glomera Reineckeana** (Kraenzlin) Schlechter, in Fedde, Repert. Nov. Spec.

9: 96, 1910.

Tau: epiphyte, top of peak back of Tau, on fallen tree trunk, alt. 650 m., Jan. 28, 1922, *Garber 748*.

Genus **Phaius** Loureiro**Phaius Tankervilleae** (Banks) Blume, Mus. Bot. Lugd.-Bat. 2: 177, 1856.

Tau: terrestrial, in dense rain forest on western slope of Mt. Vaoaimanu, alt. 300 m., Oct. 5, 1939, *Yuncker 9261*.

Genus *Calanthe* R. Brown

Calanthe furcata Bateman, Bot. Reg. 24: Misc. 28, 1838.

Tau: terrestrial, in dense, wet forest on slope two miles south of Siufaga, alt. 300 m., Oct. 30, 1939, *Yuncker* 9149.

Calanthe triplicata (Willemet f.) Ames, Philip. Jour. Sci., Bot. 2: 326, 1907.

Olosega: Olosegauta-Piumafua ridge, alt. 500 m., June 17, 1925, *Garber* 1036, 1077.

***Calanthe* species.**

Tau: terrestrial, in dense rain forest on west slope of Mt. Vaoaimanu, alt. 600 m., Oct. 5, 1939, *Yuncker* 9248.

Genus *Spathoglottis* Blume

Spathoglottis plicata Blume, Bijdr. Fl. Jav., 401, fig. 76, 1825.

Tau: terrestrial, in thicket one mile south of Siufaga, alt. 5 m., Sept. 25, 1939, *Yuncker* 9007A; in open grassy area south of Siufaga, alt. 50 m., Sept. 25, 1939, *Yuncker* 9007; bush trail back of Luma, alt. 200 m., Jan. 28, 1922, *Garber* 710, 712.

Ofu: in open grassy area on ridge back of Ofu village, alt. 400 m., Nov. 27, 1939, *Yuncker* 9527.

Olosega: Olosegauta-Piumafua ridge, alt. 500-600 m., June 17, 1925, *Garber* 1054. Native name, *niu vao*.

Genus *Dendrobium* Swartz

Dendrobium biflorum (Forster) Swartz, Nova Acta Soc. Sci. Upsal. II, 6: 84, 1799.

Tau: on fallen tree trunk, plateau back of Faleasao, alt. 150 m., Aug. 4, 1921, *Garber* 575.

Dendrobium tipuliferum Reichenbach f., Gard. Chron. 1: 72, 1877.

Tau: epiphyte, in dense rain forest on western slope of Mt. Vaoaimanu, alt. 450 m., Oct. 5, 1939, *Yuncker* 9264.

Dendrobium Tokai Reichenbach f., Hamb. Gardenz. 21: 293, 1865.

Tau: epiphytic, dark damp ravine two miles south of Siufaga, alt. 150 m., Sept. 27, 1939, *Yuncker* 9066; on dead tree, Amouli trail, alt. 150 m., Sept. 16, 1921, *Garber* 619. Native name, *laumafafia*.

Dendrobium Vaupelianum Kraenzlin, Notizbl. Bot. Gart. Berlin 5: 109, 1909.

Tau: epiphyte, in dense forest on slope about two miles south of Siufaga, alt. 150 m., Oct. 1, 1939, *Yuncker* 9067, 9160; on branch of tree in dark damp ravine on slope between one and two miles south of Siufaga, alt. 150 m., Sept. 27, 1939, *Yuncker* 9067; on breadfruit tree, Fitiuta trail, Aug. 4, 1921, *Garber* 557; on breadfruit tree, plateau back of Luma, alt. 150 m., Dec. 15, 1921, *Garber* 655. Native name, *laumafafia*.

Genus Pseuderia Schlechter

Pseuderia ramosa L. Williams, Harvard Univ., Bot. Mus. Leaflet 7: 140, 1939.

Tau: epiphyte, in dense rain forest on west slope of Mt. Vaoaimanu, alt. 300 m., Oct. 5, 1939, *Yuncker 9251*.

Pseuderia species.

Tau: epiphyte, trail to peak back of Tau, alt. 400 m., Jan. 28, 1922, *Garber 717*.

Genus Bulbophyllum Thouars

Bulbophyllum Christophersenii L. Williams, Harvard Univ., Bot. Mus. Leaflet 7: 143, 1939.

Tau: epiphyte, in deep wet forest on slope two miles south of Siufaga, alt. 300 m., Oct. 1, 1939, *Yuncker 9153*; epiphyte, in damp forest on west slope of Mt. Vaoaimanu, alt. 360 m., Oct. 5, 1939, *Yuncker 9249*.

Bulbophyllum longiscapum Rolfe, Kew Bull., 45, 1896.

Tau: epiphyte, in dark damp ravine two miles south of Siufaga, alt. 150 m., Sept. 27, 1939, *Yuncker 9076*.

Bulbophyllum species.

Tau: epiphyte, in dense moist forest on west slope of Mt. Vaoaimanu, alt. 450 m., Oct. 5, 1939, *Yuncker 9272*.

Bulbophyllum species.

Tau: epiphyte, in dense rain forest on west slope of Mt. Vaoaimanu, alt. 450 m., Oct. 5, 1939, *Yuncker 9250*.

Genus Phreatia Lindley

Phreatia Reineckeana Schlechter, in Fedde, Repert. Nov. Spec. 9: 109, 1911.

Tau: epiphyte, on dead limb of tree, Amouli trail, alt. 175 m., Sept. 16, 1921, *Garber 629*.

Phreatia samoensis (Kraenzlin) Schlechter, in Fedde, Repert. Nov. Spec. 3: 320, 1907.

Tau: epiphyte, on tree trunk, slope of peak back of Tau, alt. 650 m., Jan. 28, 1922, *Garber 745*.

Phreatia upoluensis Schlechter, in Fedde, Repert. Nov. Spec. 3: 319, 1907.

Tau: epiphyte, in open forest along the trail one mile south of Siufaga, alt. 10 m., Sept. 25, 1939, *Yuncker 9000*.

Genus Thrixspermum Loureiro**Thrixspermum species.**

Tau: epiphyte, in dense moist forest on west slope of Mt. Vaoaimanu, alt. 450 m., Oct. 5, 1939, *Yuncker 9271*.

Genus **Taeniophyllum** Blume**Taeniophyllum** species.

Tau: epiphyte, near South Point, Fitiuta trail, alt. 150 m., Oct. 12, 1921,
Garber 641.

DICOTYLEDONEAE

FAMILY CASUARINACEAE

Genus **Casuarina** Linnaeus**Casuarina equisetifolia** Linnaeus, *Amoen. Acad.* **4**: 143, 1759.

A few trees on Tau and Ofu, Guest. Native name, *toa*.

FAMILY PIPERACEAE

Genus **Piper** Linnaeus**Piper Graeffei** Warburg, *Engl. Bot. Jahrb.* **25**: 610, 1898.

Tau: climbing on trees along Fitiuta trail, Sept. 26, 1939, *Yuncker 9040, 9042, 9048.*

Ofu: clinging to tree along southeastern shore trail, Nov. 26, 1939, *Yuncker 9510.* Native name, *fue manogi*.

Piper Graeffei Warburg var. **cordatum** Warburg, *Engl. Bot. Jahrb.* **25**: 610, 1898.

Tau: clinging on tree in rocky ravine on slope south of Siufaga, alt. 100 m., Sept. 25, 1939, *Yuncker 9013*; clinging on trees along bush trail back of Luma village, alt. 225 m., Jan. 28, 1922, *Garber 711* (?). Native name, *'ava'ava aitu*.

Piper Graeffei Warburg var. (?).

Ofu: clinging on trees, leaves broadly ovate, glabrous, 7-9-nerved, slope back of Ofu village, alt. 180 m., Nov. 25, 1939, *Yuncker 9447.*

Piper Macgillivrayi C. DeCandolle var. (?) **subrotundifolium** C. DeCandolle, *Akad. Wiss. Wien, Denkschr.* **85**: 264, 1910.

Tau: clinging on tree south of Siufaga, Sept. 29, 1939, *Yuncker 9118*; along Amouli trail, Sept. 16, 1921, *Garber 638.* Native names, *'avasa*, *'ava'ava aitu*.

Piper methysticum Forster f., *De Plant Esculent.*, 76, 1786.

Tau: near dwelling, Siufaga, Oct. 3, 1939, *Yuncker 9205*; plateau back of Luma village, alt. 150 m., Dec. 15, 1921, *Garber 654.*

Ofu: edge of plantation on ridge behind Ofu village, alt. 20 m., Nov. 27, 1939, *Yuncker 9538.* Native names, *'ava*, *'ava inu*.

Genus **Peperomia** Ruiz and Pavon**Peperomia tutuilana** Yuncker, B. P. Bishop Mus., Bull. 143: 33, fig. 18, 1937.Tau: on rocks along the Fitiuta trail, Sept. 26, 1939, *Yuncker 9031*. Native name, *la'au sosolo i le ma'a*.

FAMILY ULMACEAE

Genus **Trema** Loureiro**Trema orientalis** (Linnaeus) Blume var. *viridis* Lauterbach, Engl. Bot. Jahrb. 50: 321, fig. 2-D, 1913.Tau: along Fitiuta trail, Sept. 26, 1939, *Yuncker 9046*; Fitiuta trail back of Luma, alt. 100 m., Dec. 16, 1921, *Garber 670*.Ofu: in forest on main ridge, alt. 100 m., Nov. 25, 1939, *Yuncker 9461*. Abundant on all islands, Guest. Native name, *magele*.

FAMILY MORACEAE

Genus **Broussonetia** L'Heritier**Broussonetia papyrifera** (Linnaeus) Ventenat, Tabl. Regn. Veg. 3: 547, 1794.Tau: cultivated, Siufaga, Oct. 3, 1939, *Yuncker 9204*.Olosega: plantation back of Olosegauta, alt. 350 m., *Garber 1071*. Cultivated on all the islands, Guest. Native names, *u'a, siapo*.Genus **Artocarpus** Forster**Artocarpus altilis** (Parkinson) Fosberg, Washington Acad. Sci., Jour. 31: 95, 1941.Ofu: cultivated, June 1925, *Garber 1003, 1004, 1005, 1006, 1007, 1016, 1017, 1018*. The breadfruit is cultivated in all of the villages. The natives recognize a number of varieties on the basis of the shape and size of the leaves or shape and quality of the fruit. Native name, *'ulu*.Genus **Ficus** Linnaeus**Ficus ciliata** Warburg, in Engl. Bot. Jahrb. 25: 615, 1898.Tau: in thicket near the beach south of Siufaga, Oct. 4, 1939, *Yuncker 9242*; along trail near Fitiuta, Sept. 26, 1939, *Yuncker 9033*; in thicket on slope two miles south of Siufaga, Oct. 1, 1939, *Yuncker 9145*.**Ficus Godeffroyi** Warburg, in Engl. Bot. Jahrb. 25: 613, 1898.Olosega: Olosegauta-Piumafua ridge, alt. 500-600 m., June 17, 1925, *Garber 1055, 1058, 1064*. Moderate on Tau, Ofu, and Olosega, Guest. Native name, *aoa*.**Ficus obliqua** Forster f., Fl. Ins. Austr. Prodr., 77, 1786.Ofu: in forest on ridge behind Ofu village, alt. 360 m., Nov. 27, 1939, *Yuncker 9543*; Nuu Islet, June 4, 1925, *Garber 992*.

***Ficus prolixa* Forster f., Fl. Ins. Austr. Prodr., 77, 1786.**

Tau: in thicket on plateau above Siufaga, alt. 45 m., Oct. 2, 1939, *Yuncker 9180*. Moderate on Tau, Ofu and Olosega, Guest.

?*Ficus scabra* Forster f., Fl. Ins. Austr. Prodr., 76, 1786.

Ofu: in thicket on slope behind Ofu village, Nov. 25, 1939, *Yuncker 9473*. Native name, *mati*.

***Ficus tinctoria* Forster f., Fl. Ins. Austr. Prodr., 76, 1786.**

Tau: in open forest along trail near Fitiuta, Sept. 29, 1939, *Yuncker 9052*; trail back of Luma, alt. 30 m., Aug. 21, 1921, *Garber 544*.

Ofu: Alaufau [Aloafao], June 5, 1925, *Garber 996*. Abundant on all the islands, Guest. Native name, *mati*.

FAMILY URTICACEAE

Genus *Fleurya* Gaudichaud***Fleurya interrupta* (Linnaeus) Gaudichaud, Bot. Freyc. Voy., 497, 1826.**

Tau: plateau back of Faleasao, alt. 150 m., Aug. 4, 1921, *Garber 576*; south side of South Cove, Papatea, on tuff near the beach, July 15, 1938, *W. Harris 320*; Papatea, in sand near the beach, July 15, 1938, *W. Harris 261*.

Ofu: at plantation house, Tua, June 5, 1925, *Garber 1001*; Aloafao, June 6, 1925, *Garber 1013*. Native names, *ogoogo*, *vao mageso*.

Genus *Elatostema* Forster***Elatostema basiandrum* Reinecke, in Engl. Bot. Jahrb. 25: 622, 1898.**

Tau: in stream bed, gulch on side of Palapala Hill, damp forest, alt. 300 m., July 10, 1938, *W. Harris 190*.

***Elatostema strictum* Reinecke, in Engl. Bot. Jahrb. 25: 625, 1898.**

Tau: on rocks in moist ravine, alt. 200 m., Sept. 27, 1939, *Yuncker 9081*.

Genus *Procris* Commerson***Procris pedunculata* (Forster) Weddell, DC Prodr. 16(1): 191, 1869.**

Tau: in moist thicket along Fitiuta trail, Sept. 26, 1939, *Yuncker 9051*; Fitiuta trail, Aug. 10, 1921, *Garber 607*; on lava cliff behind Luma, alt. 45 m., July 24, 1938, *W. Harris 389*; south side of South Cove, Papatea, on tuff at beach, July 15, 1938, *W. Harris 320*; in sand near the beach, Papatea, July 15, 1938, *W. Harris 261*. Native names, *fua lole*, *lole vao*, *fua feti'iti'i*.

Genus *Pipturus* Weddell***Pipturus aff. argenteus* (Forster) Weddell, DC Prodr. 16(1): 235, 19, 1869.**

Tau: in thicket on plateau back of Siufaga, alt. 45 m., Oct. 2, 1939, *Yuncker 9185*; beach at Fitiuta, Aug. 8, 1921, *Garber 585*; Faleasao, Jan. 8, 1922, *Garber 696*.

Ofu: in thicket along northwest shore, Nov. 28, 1939, *Yuncker 9547*; in thicket on ridge back of Ofu village, alt. 120 m., Nov. 25, 1939, *Yuncker 9472*;

in sandy soil along the northwest shore, alt. 5 m., Nov. 28, 1939, *Yuncker 9564, 9567*; Nuu Islet, July 4, 1925, *Garber 1110*. Abundant on all the islands, Guest. Native name, *soga*.

Genus **Maoutia** Weddell

Maoutia australis Weddell, Nouv. Arch. Mus. Hist. Nat. Paris 9: 480, 1856.

Ofu: in thicket on ridge above Ofu village, alt. 360 m., Nov. 27, 1939, *Yuncker 9530*. Native name, *ole tau'uta*.

FAMILY AMARANTHACEAE

Genus **Amaranthus** Linnaeus

Amaranthus viridis Linnaeus, Sp. Pl. ed. 2, 1405, 1763.

Tau: weed in taro patch near Siufaga, Oct. 3, 1939, *Yuncker 9197*; along trail behind Luma, Aug. 5, 1938, *W. Harris 9*; in sand near the beach, Papatea, July 15, 1938, *W. Harris 262*.

Olosega: Olosega-Olosegauta trail, June 17, 1925, *Garber 1069*. Native name, *autasi*.

Genus **Cyathula** Loureiro

Cyathula prostrata (Linnaeus) Blume, Bijdr. Fl. Jav., 549, 1825.

Tau: weed, common in sandy soil, Siufaga, Oct. 2, 1939, *Yuncker 9170*; Sani Ridge back of Fitiuta, alt. 150 m., Aug. 9, 1921, *Garber 592*; Fitiuta trail, alt. 100 m., Dec. 16, 1921, *Garber 669*; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris 215*. Native names, *manutoga, lau talatala*.

Genus **Achyranthes** Linnaeus

Achyranthes aspera Linnaeus, Sp. Pl., 204, 1753.

Tau: Fitiuta trail, alt. 125 m., Aug. 4, 1921, *Garber 563, 568*; south side of South Cove, Papatea, on tuff near beach, July 15, 1938, *W. Harris 322*. Native names, *lau tamatama, la'au talatala*.

Genus **Alternanthera** Forskål

Alternanthera sessilis (Linnaeus) R. Brown, ex Roemer and Schultes, Syst. 5: 554, 1819.

Tau: weed in sandy soil, Siufaga, Oct. 2, 1939, *Yuncker 9172*; in shallow water of taro patch near Siufaga, Oct. 6, 1939, *Yuncker 9275*; south side of South Cove, Papatea, on tuff at beach, July 15, 1938, *W. Harris 321*; along trail behind Luma, Aug. 5, 1938, *W. Harris 8*; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1939, *W. Harris 236*.

Ofu: Aloafao, June 6, 1925, *Garber 1010*. Native name, *vao sosolo*.

Genus **Gomphrena** Linnaeus

Gomphrena globosa Linnaeus, Sp. Pl., 224, 1753.

Tau: dooryard ornamental, Siufaga, Oct. 3, 1939, *Yuncker 9206*. Native name, *malila*.

FAMILY NYCTAGINACEAE

Genus *Boerhaavia* Linnaeus

Boerhaavia diffusa Linnaeus, Sp. Pl., 3, 1753.

Tau: in sandy soil, Siufaga, Oct. 6, 1939, *Yuncker* 9281; roadside, Fitiuta, Aug. 8, 1921, *Garber* 586; in sand along trail behind Luma, Aug. 1938, *W. Harris* 10; in sand near the beach, Papatea, July 15, 1938, *W. Harris* 76, 96.

Ofu: on old lava along northwestern shore, Nov. 28, 1939, *Yuncker* 9551.

Genus *Pisonia* Linnaeus

Pisonia grandis R. Brown, Prodr. Fl. Nov. Holl. 1: 422, 1810.

Ofu: on rocky talus slope at base of cliff along northwestern shore, Nov. 28, 1939, *Yuncker* 9556. Abundant on all the islands, Guest. Native name, *pu'a vai*.

Pisonia umbellifera (Forster) Seemann, Bonplandia 10: 154, 1862.

Tau: in moist ravine south of Siufaga, alt. 100 m., Sept. 25, 1939, *Yuncker* 9002; between one and two miles south of Siufaga, in dark ravine, alt. 100 m., Sept. 27, 1939, *Yuncker* 9064; plateau back of Faleasao, alt. 150 m., Aug. 4, 1921, *Garber* 574; Faleasao-Fitiuta trail, April 19, 1922, *Garber* 761.

FAMILY AIZOACEAE

Genus *Sesuvium* Linnaeus

Sesuvium Portulacastrum Linnaeus, Syst., 10th ed., 1058, 1759.

Ofu: on spray-swept lava rocks along the northwestern shore, Nov. 28, 1939, *Yuncker* 9558; rock crevices on spray-swept cliff, Ofu-Samoi trail, June 7, 1925, *Garber* 1024.

FAMILY PORTULACACEAE

Genus *Portulaca* Linnaeus

Portulaca lutea Solander, Forster f., De Pl. Esc., 72, 1786.

Ofu: on old lava along the shore, Nov. 28, 1939, *Yuncker* 9559 (this specimen is sterile but agrees well with this species in its vegetative characters); rocky cliff, Feaiia Point, June 7, 1925, *Garber* 1023. Native name, *tamole*.

Portulaca oleracea Linnaeus, Sp. Pl., 445, 1753.

Tau: weed in taro patch, near Luma, Oct. 6, 1939, *Yuncker* 9279; Papatea, in sand near the beach, July 15, 1938, *W. Harris* 79, 99. Native name, *tamole*.

Portulaca samoensis von Poellnitz, in Fedde, Repert. Nov. Spec. 33: 163, 1933.

Tau: on lava rocks near the shore along the Luma-Faleasao trail, Sept. 20, 1939, *Yuncker* 9122; Faasamene Spring near Faleasao, Jan. 8, 1922, *Garber* 697; Papatea, in sandy soil near the beach, July 15, 1938, *W. Harris* 77, 78, 80, 97, 98.

Ofu: in sandy soil near the beach along the southeastern shore, Nov. 26, 1939, Yuncker 9518; along the Ofu-Samoi trail on exposed wind-swept rocky cliffs, June 7, 1925, Garber 1025. Native name, *tamole*.

FAMILY ANNONACEAE

Genus **Cananga** Hooker f. and Thomson

Cananga odorata (Lamarck) Hooker f. and Thomson, Fl. Ind. 1: 130, 1855.

Tau: in thicket on plateau above Luma, alt. 45 m., Oct. 2, 1939, Yuncker 9178; on plateau back of Lumā, alt. 125 m., Garber 650. Abundant on all the islands, Guest. Native name, *moso'oi*.

FAMILY MYRISTICACEAE

Genus **Myristica** Linnaeus

Myristica inutilis Rich, ex A. Gray, U. S. Expl. Exped., Bot. 15(1): 33, 1854.

Tau: in thicket on plateau behind Siufaga, alt. 45 m., Oct. 2, 1939, Yuncker 9182; in forest on slope two miles south of Siufaga, alt. 150 m., Oct. 1, 1939, Yuncker 9161; Fitiuta trail back of Faleasao, Aug. 11, 1921, Garber 609.

Ofu: in forest on main ridge, Nov. 27, 1939, Yuncker 9541. Abundant on all the islands, Guest. Native name, *atone*.

FAMILY MONIMIACEAE

Genus **Hedycarya** Forster

Hedycarya denticulata (A. Gray) Perkins and Gilg, Pflanzenr. 49(IV.101): 19, 1901.

Ofu: on central ridge, alt. 250 m., Nov. 27, 1939, Yuncker 9523. Native name, *la'au fai vaitusi*.

FAMILY LAURACEAE

Genus **Cinnamomum** Linnaeus

Cinnamomum pallidum Gillespie, B. P. Bishop Mus., Bull. 91: 6, fig. 5, 1932.

Tau: a few trees at Fitiuta, Guest. Native name, *tigamoni*.

Genus **Litsea** Lamarck

?**Litsea magnifolia** Gillespie var. **samoensis** Christophersen, B. P. Bishop Mus., Bull. 128: 89, fig. 10, 1935.

Tau: abundant at Fitiuta, Guest. Native name, *papaono*.

FAMILY HERNANDIACEAE

Genus **Gyrocarpus** Jacquin

Gyrocarpus americanus Jacquin, Stirp. Am., 282, pl. 178, 1763.

Ofu: Nuu Islet, July 4, 1925, Garber 1108.

Genus **Hernandia** Plumier**Hernandia Moerenhoutiana** Guillemin, Sci. Nat., Ann. II, 7: 189, 1837.Tau: abundant, Guest. Native name, *pipi*.**Hernandia ovigera** Linnaeus, in Stickman Herb. Amb., 14, 1754.Tau: Amouli beach trail, Jan. 7, 1922, *Garber* 676.Ofu: in thicket along southeastern shore, Nov. 26, 1939, *Yuncker* 9511.Abundant on all islands, Guest. Native name, *pu'a*.

FAMILY CRUCIFERAE

Genus **Nasturtium** R. Brown**Nasturtium sarmentosum** (Solander) O. E. Schulz, Engl. Bot. Jahrb. 32: 595, 1902.Tau: weed in sandy soil, Siufaga, Oct. 2, 1939, *Yuncker* 9171; on beach, Faleasao-Fanga [Faga] trail, Apr. 19, 1922, *Garber* 762; south side of South Cove, Papatea, on tuff at beach, July 15, 1938, *W. Harris* 325; in sand near the beach, Papatea, July 15, 1938, *W. Harris* 81, 100.Olosega: taro patch, back of Olosega village, June 23, 1925, *Garber* 1089. Native name, *lau taliga*.

FAMILY CAPPARIDACEAE

Genus **Crataeva** Linnaeus**Crataeva religiosa** Forster f., Fl. Ins. Austr. Prodr., 35, 1786.Ofu: exposed coast near Mafafa, alt. 40 m., May 26, 1925, *Garber* 983.

FAMILY CUNONIACEAE

Genus **Weinmannia** Linnaeus**Weinmannia manuana** Christophersen, B. P. Bishop Mus., Bull. 154: 10, fig. 2, 1938.Tau: old taro clearing on trail to peak, alt. 530 m., Jan. 28, 1922, *Garber* 720.Olosega: top of Piumafua mountain, alt. 630 m., June 17, 1925, *Garber* 1027, 1031; Piumafua ridge, alt. 480 m., June 17, 1925, *Garber* 1066. Native name, *laga'ali*.

FAMILY ROSACEAE

Genus **Parinarium** Aublet**Parinarium glaberrimum** Hasskarl, Flora 27: 583, 1844.

Tau: abundant at Fitiuta, Guest.

Ofu: scarce, Guest. Native name, *ififi*.**Parinarium insularum** A. Gray, U. S. Expl. Exped., Bot. 15(1): 488, pl. 54, 1854.Scarce on Tau and Olosega, Guest. Native name, *sea*.

FAMILY LEGUMINOSAE

Genus **Leucaena** Bentham

Leucaena glauca (Linnaeus) Bentham, Hook. Jour. Bot. 4: 416, 1842.

Tau: in thicket near the sea, south of Siufaga, Sept. 29, 1939, Yuncker 9108. Native name, *lopa samoae*.

Genus **Adenanthera** Royen

Adenanthera pavonina Linnaeus, Sp. Pl., 384, 1753.

Abundant on all of the islands, Guest. Native name, *lopa*.

Genus **Caesalpinia** Linnaeus

Caesalpinia jayabo Maza, Soc. Esp. Hist. Nat., Anal. 19: 234, 1890.

Ofu: at Aofa, alt. 275 m., Sept. 3, 1925, Garber 1115. Native name, *anaoso*.

Caesalpinia pulcherrima (Linnaeus) Swartz, Observ. Bot., 166, 1791.

Ofu: cultivated ornamental, village of Ofu, June 1925, Garber 1087. Native name, *lau pa*.

Genus **Sophora** Linnaeus

Sophora tomentosa Linnaeus, Sp. Pl., 373, 1753.

Ofu: along southeastern shore near the beach, Nov. 26, 1939, Yuncker 9496; beach at Muliulu, May 26, 1925, Garber 985. Native name, *fua feti iti*.

Genus **Indigofera** Linnaeus

Indigofera suffruticosa Miller, Gard. Dict., 8th ed., no. 2, 1768.

Tau: along Luma-Faleasao trail, alt. 50 m., Sept. 30, 1939, Yuncker 9127; along trail near Fitiuta village, Sept. 26, 1939, Yuncker 9053; plateau back of Luma, alt. 60 m., Aug. 2, 1921, Garber 550; plateau back of Luma, alt. 105 m., May 28, 1937, W. Harris 271.

Genus **Tephrosia** Persoon

Tephrosia purpurea (Linnaeus) Persoon, Syn. Pl. 2: 329, 1807.

Tau: in sandy soil near the shore south of Siufaga, Oct. 4, 1939, Yuncker 9229.

Ofu: in sandy soil near the shore, Ofu village, Nov. 25, 1939, Yuncker 9490; Aloafao, June 4, 1925, Garber 988. Native name, 'avasa.

Genus **Desmodium** Desvaux

Desmodium heterocarpum (Linnaeus) DeCandolle, Prodr. 2: 337, 1825.

Tau: in thicket along trail south of Siufaga, Sept. 27, 1939, Yuncker 9083; Sani Ridge back of Fitiuta, alt. 150 m., Aug. 9, 1921, Garber 603; Amouli trail, alt. 175 m., Sept. 16, 1921, Garber 634; plateau back of Luma, alt. 100 m., Dec. 15, 1921, Garber 645a.

Ofu: in coconut grove near the village of Ofu, alt. 150 m., Nov. 25, 1939, Yuncker 9469; back of Ofu village, alt. 40 m., May 25, 1925, Garber 979.

Desmodium umbellatum (Linnaeus) DeCandolle, Prodr. 2: 325, 1825.

Tau: in thicket along Luma-Faleasao trail, alt. 45-50 m., Sept. 30, 1939, *Yuncker 9139*; on lava cliff near the sea, Luma-Faleasao trail, alt. 45-50 m., Sept. 30, 1939, *Yuncker 9121*; Luma-Faleasao trail, Jan. 8, 1922, *Garber 692*. Native names, *lala, lau lala*.

Genus *Uraria* Desvaux

Uraria lagopodioides (Linnaeus) Desvaux, Soc. Linn. Paris, Mém. 4: 309, 1826.

Tau: plateau back of Luma, alt. 100 m., Dec. 15, 1921, *Garber 645b*. Native name, *lala*.

Genus *Lourea* Necker

Lourea vespertilionis (Linnaeus f.) Desvaux, Jour. Bot. 1: 122, pl. 5, fig. 18, 1813.

Tau: cultivated ornamental, Luma, Oct. 3, 1939, *Yuncker 9207*.

Genus *Inocarpus* Forster

Inocarpus fagiferus (Parkinson) Fosberg, Washington Acad. Sci., Jour. 31: 95, 1941.

Tau: in open woods along Luma-Faleasao trail, alt. 45-50 m., Sept. 30, 1939, *Yuncker 9130*; Luma, Dec. 15, 1921, *Garber 661*. Abundant on all of the islands, Guest. Native name, *ifi*.

Genus *Abrus* Linnaeus

Abrus precatorius Linnaeus, Syst., ed. 12, 472, 1767.

Ofu: Nuu Islet, July 4, 1925, *Garber 1111*. Native name, *fuefue laulili'i*.

Genus *Erythrina* Linnaeus

Erythrina variegata Linnaeus var. *orientalis* (Linnaeus) Merrill, Interpret. Herb. Amb., 276, 1917.

Tau: in thicket near the shore two miles south of Siufaga, Oct. 1, 1939, *Yuncker 9143*; plateau back of Luma, alt. 100-125 m., Aug. 4, 1921, *Garber 581*.

Ofu: along the southeastern shore, Nov. 26, 1939, *Yuncker 9502*; at water's edge, Toanga [Toaga], June 23, 1925, *Garber 1094*. Moderate on Olosega, Guest. Native name, *gatae*.

Genus *Mucuna* Adanson

Mucuna gigantea (Willdenow) DeCandolle, Prodr. 2: 405, 1825.

Ofu: climbing in trees along the southeastern shore, Nov. 26, 1939, *Yuncker 9512*. Native name, *fuefue*.

Genus *Pueraria* DeCandolle

Pueraria Thunbergiana (Siebold and Zuccarini) Bentham, Linn. Soc. Bot., Jour. 9: 122, 1865.

Tau: plateau back of Luma, alt. 125 m., Dec. 15, 1921, *Garber 651*.

Genus ***Canavalia*** DeCandolle***Canavalia rosea*** (Swartz) DeCandolle, Prodr. 2: 404, 1825.Tau: sandy beach, Faga-Faleasao, Apr. 19, 1922, *Garber* 763.Ofu: plantation, June 5, 1925, *Garber* 1000; beach, Aloafao, July 5, 1925, *Garber* 1114. Native names, *va'a*, *fuefue fai va'a*.Genus ***Vigna*** Savi***Vigna marina*** (Burman) Merrill, Interpret. Herb. Amb., 285, 1917.Tau: in sandy soil near the beach south of Siufaga, Sept. 29, 1939, *Yuncker* 9101; sandy beach, Fangamalo [Fagamalo], Jan. 7, 1922, *Garber* 684; Faleasao-Fitiuta trail, alt. 20 m., Jan. 8, 1922, *Garber* 700.Ofu: beach, Aloafao, July 5, 1925, *Garber* 1113. Native names, *lau fuefue*, *fuefue*.

FAMILY OXALIDACEAE

Genus ***Oxalis*** Linnaeus***Oxalis corniculata*** Linnaeus var. *repens* (Thunberg) Zuccarini, Akad.

Muensch., Abh. 1: 230, 1829-30.

Tau: in sand near the beach, Papatea, July 15, 1938, *W. Harris* 82; on lava cliff behind Luma, alt. 45 m., July 24, 1938, *W. Harris* 34.Ofu: on rocks about the water reservoir near Ofu village, alt. 15 m., Nov. 25, 1939, *Yuncker* 9482.Olosega: plantation above Olosegauta, alt. 330 m., June 17, 1925, *Garber* 1082. Native name, *a'aa'a*.

FAMILY RUTACEAE

Genus ***Evodia*** Forster***Evodia hortensis*** Forster f. *simplicifolia* (Rechinger) K. Schumann, Engl.

Bot. Jahrb. 55: 232, 1918.

Ofu: cultivated ornamental, Ofu village, Nov. 25, 1939, *Yuncker* 9446. Abundant on all the islands, Guest. Native name, *usi*.***Evodia samoensis*** Christophersen, B. P. Bishop Mus., Bull. 154: 12, 1938.Tau: in thicket on plateau behind Siufaga, Oct. 2, 1939, *Yuncker* 9193; Amouli trail, alt. 200 m., Sept. 16, 1921, *Garber* 617.Ofu: in open forest on central ridge, alt. 180-200 m., Nov. 25, 1939, *Yuncker* 9448.Olosega: top of Piumafua mountain, June 17, 1925, *Garber* 1026. Native name, *so'opini*.Genus ***Micromelum*** Blume***Micromelum minutum*** (Forster f.) Seemann, Viti, Govt. Mission, 434, 1862.Tau: Fitiuta trail, Dec. 16, 1921, *Garber* 668.Ofu: Nuu Islet, June 4, 1925, *Garber* 990. On all the islands, Guest. Native name, *talafalu*.

Genus **Citrus** Linnaeus

Citrus aurantifolia (Christman) Swingle, Washington Acad. Sci., Jour. 3 : 465, 1913.

Ofu: Aloafao, alt. 5 m., June 6, 1925, *Garber* 1015. Found in all villages in the islands, Guest. Native name, *moli tipolo*.

Citrus Aurantium Linnaeus, Sp. Pl., 782, 1753.

A few seedling trees are found around every village, Guest. Native name, *moli aina*.

Citrus Hystrix DeCandolle, Cat. Hort. Monsp., 97, 1813.

Tau: Fitiuta trail, Aug. 4, 1921, *Garber* 569. Found in or around almost every village, Guest. Native name, *moli u'u*.

Citrus maxima (Burman) Merrill, Interpret. Herb. Amb., 296, 1917.

Tau: a few trees at Fitiuta, Guest. Native name, *moli si-suka*.

Citrus Medica Linnaeus, Sp. Pl., 782, 1753.

A few trees in almost every village, Guest. Native names, *moli patupatu*, *tipolo*.

FAMILY BURSERACEAE

Genus **Canarium** Linnaeus

Canarium samoense Engler, DC Monog. Phanerog. 4: 134, 1883.

On all of the islands, Guest. Native name, *ma'ali*.

Canarium species.

Tau: abundant at village of Tau, Guest. Native name, *a'amati'e*.

FAMILY MELIACEAE

Genus **Dysoxylum** Blume

Dysoxylum Richii (A. Gray) DeCandolle, Monog. Phanerog. 1: 511, 1878.

Tau: in open forest on plateau behind Siufaga, alt. 45 m., Oct. 2, 1939, *Yuncker* 9183. Native name, *maota*.

Genus **Aglaia** Loureiro

Aglaia samoensis A. Gray, U. S. Expl. Exped., Bot. 15(1) : 236, 1854.

Ofu: plantation at Toaga, June 22, 1925, *Garber* 1098. Abundant on all of the islands, Guest. Native name, *laga'ali*.

FAMILY POLYGALACEAE

Genus **Polygala** Linnaeus

Polygala paniculata Linnaeus, Amoen. Acad. 5: 402, 1759.

Tau: along trail on plateau behind Siufaga, Oct. 2, 1939, *Yuncker* 9175. Native names, *fua paepae*, *fua lili'i*.

FAMILY EUPHORBIACEAE

Genus *Glochidion* Forster

Glochidion ramiflorum Forster var. ***samoanum*** (Mueller-Argau) Pax, Reinecke, Engl. Bot. Jahrb. 25: 645, 1898.

Ofu: in forest on central ridge, alt. 200 m., Nov. 25, 1939, *Yuncker* 9463; in forest on ridge back of Ofu village, alt. 150 m., Nov. 25, 1939, *Yuncker* 9459; in forest on ridge above Ofu village, alt. 350 m., Nov. 27, 1939, *Yuncker* 9519.

Olosega: top of Piumafua mountain, alt. 625 m., June 17, 1925, *Garber* 1038. Abundant on all of the islands, Guest. Native name, *masame*.

Genus *Breynia* Forster

Breynia nivosa (Bull) Small, Torrey Bot. Club, Bull. 37: 516, 1910.

Tau: cultivated ornamental, Siufaga, Oct. 3, 1939, *Yuncker* 9195.

Genus *Phyllanthus* Linnaeus

Phyllanthus Niruri Linnaeus, Sp. Pl., 981, 1753.

Tau: weed in plantation, plateau back of Siufaga, alt. 45 m., Oct. 2, 1939, *Yuncker* 9188; in sand, Luma, July 19, 1938, *W. Harris* 24, 343; side of trail behind Luma, Aug. 5, 1938, *W. Harris* 15; in sand near the beach, Papatea, July 15, 1938, *W. Harris* 87, 104.

Ofu: weed near the shore, Nov. 28, 1939, *Yuncker* 9552. Native names, *lau lili'i, fua lili'i*.

Phyllanthus simplex Retzius, Obs. 5: 29, 1789.

Tau: among ferns on moist bank along Luma-Faleasao trail, alt. 50 m., Sept. 30, 1939, *Yuncker* 9133; Luma-Faleasao trail, alt. 40 m., Jan. 8, 1922, *Garber* 693.

Olosega: along the trail at Olosegauta, alt. 250 m., June 17, 1925, *Garber* 1074.

Genus *Securinega* A. L. deJussieu

Securinega samoana Croizat, new species.

Securinega (§ *Flueggea*) species, Christophersen, B. P. Bishop Mus., Bull. 128: 120, 1935.

Arbor parva, ligno durissimo insignis, innovationibus flexuosis. Foliis cum petiolis glaberrimis, more generis distichis 9-18 cm. longis, 3-7 cm. latis, tenuissime chartaceis, in sicco olivaceis subconcoloribus, oblanceolato-cuspidatis vel oblanceolato-ellipticis, margine integerrimis, venis ca. 6-7-jugis, petiolo canaliculato vix 5-8 mm. longo. Floribus in glomerulis axillaribus congestis ♂ aut ♀, pulvinulum vix 1-2 cm. longum efformantibus. Perianthio ♂ ca. 2 mm. longo, latoque, lobis ovato-rotundatis 5; staminibus 5 oppositilobis, cum glandulis magnitudine formaque variis alternantibus, filamento ca. 1 mm. longo, antheris ca. 0.25 mm. longis, pedicello ca. 3 mm. longo, gracili; pistillodio carnoso 2-partito. Periantio ♀ illi ♂ subsimili, ovario globuloso-truncato glaberrimo ca. 1-1.5 mm. magno, stylis 3 e basi liberis apice retuse bifidis, semine brunneo leviusculo ad 1.5 mm. longo.

Tau: Siufaga, Oct. 3, 1939, "Tree 6 m. tall, 20 cm. diam." *Yuncker* 9196.

Savaii: Tufutafoe-Falelima, "Edge of forest, alt. about 25 m., tree 5 m. high, female, fr. black", Sept. 1931, *Christophersen* 2765. Type in herbarium of New York Botanical Garden.

Upolu: Vailima "Slopes and rocky woods alt. 100 m. Tree 10-25 m. high, diam. 14-20 cm.", Feb. 1921, *A. J. Eames* 85; Falealupo, 1931, *Christophersen* 2794; Saluafata, "Poumouli, fruit blue black", 1929, *Christophersen* 943.

Without locality, *Y. Powell* 1880.

Christophersen doubts that this plant is native to Samoa and it must be admitted that its characters bring it close to *S. flexuosa* Mueller-Argau, endemic to the Philippine Islands. As a matter of fact, it would prove impossible to separate the Samoan plant specifically from that of the Philippines if the ranges were in common. A careful comparison of representative sets of *S. flexuosa* and *S. samoana* of both sexes and in every stage of growth shows, however, that (1) the Samoan plant has foliage which tends to be more broadly oblong and short-cuspidate than that of the Philippines; (2) the pedicel of the female flower near the young fruit stage reaches 6-8 mm. in *S. flexuosa*, while it does not exceed 4-5 mm. in *S. samoana*; (3) the style of *S. flexuosa* is more or less stipitate, instead of sessile, as in *S. samoana*; (4) the seed is slightly longer and manifestly more voluminous in *S. flexuosa* than it is in *S. samoana*. All these differences, taken severally, would be trifling, but together, with the different ranges, suggest that the description of a new species is advisable. The specific characters of *Securinega* are seldom strongly marked among species of the same affinity.

With the final addition of *S. samoana*, *Securinega* becomes one of the most widespread genera, its range including Oceania, Australasia, Asia proper, Africa, Europe, and America. Native name, *poumuli*.

Genus *Bischofia* Blume

Bischofia javanica Blume, *Bijdr. Fl. Jav.*, 1168, 1826.

Tau: plantation back of Luma, Sept. 13, 1921, *Garber* 613; plateau back of Luma, alt. 105 m., May 29, 1937, *W. Harris* 289. Abundant on all of the islands, Guest. Native name, 'o'a.

Genus *Aleurites* Forster

Aleurites moluccana (Linnaeus) Willdenow, *Sp. Pl.* 4: 590, 1805.

Tau: plateau back of Luma, alt. 100 m., Sept. 21, 1922, *Garber* 777.

Ofu: on ridge back of Ofu village, alt. 100 m., Nov. 27, 1939, *Yuncker* 9540.

Olosega: abundant, Guest. Native name, *lama*.

Genus *Macaranga* Thouars

Macaranga Harveyana Mueller-Argau, DC Prodr. 15(2): 998, 1866.

Tau: along the Fitiuta trail, alt. 50 m., Sept. 26, 1939, *Yuncker* 9037; Amouli trail, alt. 75 m., Sept. 16, 1921, *Garber* 614.

Ofu: in forest on main ridge, alt. 150 m., Nov. 25, 1939, *Yuncker 9467*; Feaiia Point, alt. 75 m., May 25, 1925, *Garber 978*. Native names, *pata*, *pata laitiiti*.

Macaranga stipulosa Mueller-Argau, DC Prodr. 15(2) : 1001, 1866.

Tau: Amouli trail, alt. 200 m., Sept. 16, 1921, *Garber 636*.

Olosega: Olosegauta-Piumafua Ridge, alt. 500 m., June 17, 1925, *Garber 1072*. Abundant on all of the islands, Guest. Native name, *pata*.

Genus **Acalypha** Linnaeus

Acalypha boehmerioides Miquel, Fl. Ind. Bat. Suppl., 459, 1861-62.

Tau: dooryard weed, Siufaga, Oct. 6, 1939, *Yuncker 9280*; along the trail behind Luma, Aug. 5, 1938, *W. Harris 7*; in sand near the beach, Papatea, July 15, 1938, *W. Harris 74*.

Ofu: Aloafao, June 6, 1925, *Garber 1014*. Native name, *ogoogo*.

Acalypha grandis Bentham, Hook. Lond. Jour. Bot. 2 : 232, 1843.

Olosega: at beach, Asanga [Asaga] Pass, June 23, 1925, *Garber 1093*. Native name, *lala*.

Acalypha hispida Burman f., Fl. Ind., 303, pl. 61, fig. 1, 1768.

Tau: cultivated ornamental, Siufaga, Oct. 2, 1939, *Yuncker 9168*.

Acalypha Wilkesiana Mueller-Argau, DC Prodr. 15(2) : 817, 1866.

Tau: cultivated ornamental, Siufaga, Oct. 3, 1939, *Yuncker 9223*.

Genus **Codiaeum** (Rumphius) Jussieu

Codiaeum variegatum (Linnaeus) Blume var. *pictum* (Loddiges) Mueller-Argau, DC Prodr. 15(2) : 1119, 1866.

Tau: on plateau back of Luma, alt. 105 m., Aug. 4, 1922, *Garber 769* (form *appendiculatum* Celakovsky). A number of different forms cultivated for ornament about dwellings on the islands of Tau and Ofu were noted but not collected. Native name, *motu*.

Genus **Manihot** Miller

Manihot esculenta Crantz, Inst. Rei Herb. 1 : 167, 1766.

Tau: about dwellings, Luma, Sept. 30, 1939, *Yuncker 9138*. Native names, *ufi maniota*, *manioka*, *tapioka*.

Genus **Homalanthus** Jussieu

Homalanthus acuminatus (Mueller-Argau) Pax, Reinecke, Engl. Bot. Jahrb. 25 : 648, 1898.

Abundant on all of the islands, Guest. Native name, *mamala*.

Homalanthus nutans (Forster) Pax, in Engler and Prantl, Nat. Pflanzenfam. 3(5) : 96, 1890.

Tau: in thicket along Fitiuta trail, alt. 50 m., Sept. 26, 1939, *Yuncker 9045*; Fitiuta trail back of Luma, alt. 60 m., Aug. 4, 1921, *Garber 553*.

Ofu: in thicket on ridge back of Ofu village, alt. 300 m., Nov. 27, 1939, *Yuncker* 9534.

Olosega: Olosegauta-Piumafua Ridge, alt. 475 m., June 17, 1925, *Garber* 1029. Native names, *mamala, fanuamamala*.

Genus *Euphorbia* Linnaeus

Euphorbia Atoto Forster f., Fl. Ins. Austr. Prodr., 207, 1786.

Tau: in sandy soil near the shore, Sept. 29, 1939, *Yuncker* 9112; in sandy soil, Luma, July 20, 1938, *W. Harris* 340; in sand near the beach, Papatea, July 15, 1938, *W. Harris* 84.

Ofu: coral beach, Muliulu, May 26, 1925, *Garber* 986. Native names, *pulutai, a'atasi*.

Euphorbia hirta Linnaeus, Sp. Pl., 454, 1753.

Tau: weed along Luma-Faleasao trail, Sept. 30, 1939, *Yuncker* 9120; Luma-Faleasao trail, alt. 40 m., Jan. 8, 1922, *Garber* 691; Luma, in sand near the beach, July 20, 1938, *W. Harris* 13, 14, 23, 341; Papatea, in sand near the beach, July 15, 1938, *W. Harris* 83, 85, 103; Faleulu, in plantation, alt. 180 m., July 27, 1938, *W. Harris* 237. Native name, *la'au fai moti*.

Euphorbia prostrata Aiton, Hort. Kew. 2: 136, 1789.

Tau: in sandy soil near the beach, Oct. 2, 1939, *Yuncker* 9116; Luma, in sand near the beach, July 20, 1938, *W. Harris* 342; Papatea, in sand near the beach, July 15, 1938, *W. Harris* 86. Native name, *lau lili'i*.

FAMILY ANACARDIACEAE

Genus *Buchanania* Roxburgh

Buchanania Merrillii Christophersen, B. P. Bishop Mus., Bull. 154: 14, 1938.

Tau: abundant in the mountains and at Fitiuta, Guest. Native name, *nasu*.

Genus *Mangifera* Linnaeus

Mangifera indica Linnaeus, Sp. Pl., 200, 1753.

Tau: Siufaga, Oct. 1, 1939, *Yuncker* 9158. Seedling trees found about almost every village on the islands, Guest. Native name, *mago*.

Genus *Spondias* Linnaeus

Spondias dulcis Parkinson, Jour. Voy. Endeavour, 39, 1773.

Tau: cultivated, Siufaga, Jan. 11, 1922, *Garber* 706. Common on all of the islands, Guest. Native name, *vi*.

Genus *Rhus* Linnaeus

Rhus taitensis Guillemin, Sci. Nat. Bot., Ann. II, 7: 361, 1837.

Tau: in forest on west slope of Mt. Vaoaimanu, alt. 450 m., Oct. 5, 1939, *Yuncker* 9245.

Ofu: in forest on central ridge, alt. 300 m., Nov. 27, 1939, *Yuncker* 9533. Abundant on all of the islands, Guest. This is a common forest tree. The above cited specimens agree well with those from Upolu and Savaii as reported by Christophersen, which he doubtfully refers to this species. Native name, *tavai*.

FAMILY ICACINACEAE

Genus *Chariessa* Miquel

Chariessa samoensis (A. Gray) Engler, in Engler and Prantl, Nat. Pflanzenfam. III, 5: 245, 1893.

Abundant on all of the islands, Guest. Native names, *ala'a*, *aloalo vao*.

FAMILY SAPINDACEAE

Genus *Pometia* Forster

Pometia pinnata Forster, Charact. Gen. Pl., 110, pl. 55, 1776.

Abundant on all of the islands, Guest. Native name, *tava*.

Genus *Alectryon* Gaertner

Alectryon samoensis Christophersen, B. P. Bishop Mus., Bull. 128: 130, fig. 16, 1935.

Moderate on all of the islands, Guest. Native name, *taputo'i*.

FAMILY RHAMNACEAE

Genus *Colubrina* L. C. Richard

Colubrina asiatica (Linnaeus) Brongniart, Sci. Nat., Ann. 10: 369, 1827.

Tau: in thicket along Luma-Faleasao trail, alt. 50 m., Sept. 30, 1939, *Yuncker* 9126; Luma-Faleasao trail, alt. 40 m., Jan. 8, 1922, *Garber* 687.

Ofu: Nuu Islet, June 4, 1925, *Garber* 993. Native name, *fisoa*.

Genus *Alphitonia* Reissek

Alphitonia zizyphoides (Sprengel) A. Gray, U. S. Expl. Exped., Bot. 15(1): 278, 1854.

Tau: Fitiuta trail, alt. 100 m., Aug. 4, 1921, *Garber* 559.

Ofu: Nuu Islet, alt. 75 m., July 4, 1925, *Garber* 1104. Abundant on all of the islands, Guest. Native name, *toi*.

FAMILY TILIACEAE

Genus *Trichospermum* Blume

Trichospermum Richii (A. Gray) Seemann, Bonplandia 9: 254, 1861.

Tau: in forest on slope south of Siufaga, alt. 300 m., Oct. 4, 1939, *Yuncker* 9231.

Genus *Grewia* Linnaeus

Grewia crenata (J. R. and G. Forster) Setchell, Dept. Marine Biol. Carnegie Inst. Washington 20: 72, 1924.

Abundant on all of the islands, Guest. Native name, *fau'ui*.

Genus *Triumfetta* Linnaeus

Triumfetta Bartramia Linnaeus, Syst., ed. 10, 1044, 1759.

Tau: Sani Ridge back of Fitiuta, alt. 150 m., Aug. 9, 1921, *Garber* 602.

Olosega: Olosegauta, alt. 250 m., June 17, 1925, *Garber* 1076. Native name, *mautofu vao*.

Triumfetta procumbens Forster f., Prodr., 35, 1786.

Tau: in sandy soil, coconut grove near the shore, Sept. 25, 1939, *Yuncker* 9022; beach at Fitiuta, Aug. 8, 1921, *Garber* 582; sandy beach, Faga, April 20, 1922, *Garber* 764.

Ofu: in sandy soil along southeastern shore, Nov. 26, 1939, *Yuncker* 9497. Native name, *fuatala*.

FAMILY MALVACEAE

Genus *Sida* Linnaeus

Sida rhombifolia Linnaeus, Sp. Pl., 684, 1753.

Tau: in sandy soil near the shore, Oct. 1, 1939, *Yuncker* 9144; plateau back of Luma, alt. 60 m., Aug. 2, 1921, *Garber* 548; along the trail behind Luma, Aug. 5, 1938, *W. Harris* 16; in sand along the beach, Papatea, July 15, 1938, *W. Harris* 88; in coconut, breadfruit and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 195.

Ofu: Tua, June 7, 1925, *Garber* 1020.

Olosega: Olosegauta-Piumafua Ridge, alt. 525 m., June 17, 1925, *Garber* 1075. Native name, *mautofu*.

Sida samoensis Rechinger, in Fedde, Report. Nov. Spec. 4: 228, 1907.

Tau: in sand near the beach, Oct. 2, 1939, *Yuncker* 9167.

Genus *Urena* Linnaeus

Urena lobata Linnaeus, Sp. Pl., 692, 1753.

Tau: in sandy soil near the shore two miles south of Siufaga, Oct. 1, 1939, *Yuncker* 9142; plateau back of Luma, alt. 60 m., Aug. 2, 1921, *Garber* 545.

Ofu: weed in plantation near the village of Ofu, Nov. 25, 1939, *Yuncker* 9492. Native name, *mautofu*.

Genus *Hibiscus* Linnaeus

Hibiscus Abelmoschus Linnaeus, Sp. Pl., 696, 1753.

Tau: in swampy area back of Siufaga, Oct. 7, 1939, *Yuncker* 9298; in clearing along trail to Fitiuta, Sept. 26, 1939, *Yuncker* 9035; Sani Ridge above

Fitiuta, alt. 150 m., Aug. 9, 1921, Garber 601; Luma-Faleasao trail, Oct. 1930, Judd.

Olosega: taro plantation back of Olosegauta, alt. 350 m., June 17, 1925, Garber 1085. Native names, *fau tagaloa*, *fua samasama*, *vavae samoan*.

Hibiscus Rosa-sinensis Linnaeus, Sp. Pl., 694, 1753.

Ofu: near the beach, village of Ofu, Nov. 28, 1939, Yuncker 9565. A number of hybrid varieties of *Hibiscus* are in cultivation as ornamentals in the various villages about the islands. Native name, *autē*.

Hibiscus Sabdariffa Linnaeus, Sp. Pl., 695, 1753.

Ofu: cultivated, Aloafao, June 4, 1925, Garber 989. Native name, *uaina*.

Hibiscus tiliaceus Linnaeus, Sp. Pl., 694, 1753.

Tau: near the shore south of Siufaga, Sept. 29, 1939, Yuncker 9094; at the beach, Fitiuta, Aug. 8, 1921, Garber 583.

Ofu: near the shore along the southeastern shore, Nov. 26, 1939, Yuncker 9495. Abundant on all of the islands, Guest. Native name, *fau*.

Genus *Thespesia* Solander

Thespesia populnea (Linnaeus) Solander, ex. Correa, Mus. Hist. Nat. Paris, Ann. 9: 290, 1807.

Tau: beach at Fagamalo, Jan. 7, 1922, Garber 683.

Ofu: near the shore on the northwestern shore, Nov. 28, 1939, Yuncker 9570; rocky coast near Tuafanua, May 26, 1925, Garber 982. Moderate to abundant on all of the islands, Guest. Native name, *milo*.

Genus *Gossypium* Linnaeus

Gossypium brasiliense Macfayden, Fl. Jam. 1: 72, 1837.

Tau: near dwelling, Siufaga, Oct. 3, 1939, Yuncker 9217.

Ofu: Nuu Islet, July 4, 1925, Garber 1105. Native name, *vavae samoan*.

FAMILY BOMBACACEAE

Genus *Ceiba* Medikus

Ceiba pentandra (Linnaeus) Gaertner, Fruct. et Semin. 2: 224, pl. 133, fig. 1, 1791.

Tau: in clearing on plateau back of Siufaga, alt. 45 m., Oct. 2, 1939, Yuncker 9189.

Ofu: cultivated, Aloafao, June 26, 1925, Garber 1056. In every village on the islands, Guest. Native names, *vavae*, *vavae papalagi*.

FAMILY STERCULIACEAE

Genus *Melochia* Linnaeus

Melochia odorata Linnaeus f., Suppl., 302, 1781.

Abundant on all of the islands, Guest. Native name, *ma'o*.

Genus **Theobroma** Linnaeus**Theobroma Cacao** Linnaeus, Sp. Pl., 782, 1753.

A few trees near the village of Tau on the island of Tau and also on Ofu, Guest. Native name, *koko* [*toto*].

Genus **Kleinhovia** Linnaeus**Kleinhovia hospita** Linnaeus, Sp. Pl., ed. 2, 1365, 1763.

Tau: Fitiuta trail, alt. 125 m., Aug. 4, 1921, *Garber* 565. Abundant on all of the islands, Guest. Native name, *fu'afu'a*.

Genus **Sterculia** Linnaeus

Sterculia fanaiho Setchell, Dept. Marine Biol. Carnegie Inst. Washington 20: 69, 1924.

Tau: plateau back of Luma, alt. 100 m., Sept. 21, 1921, *Garber* 775.

Ofu: in open woods along the southeastern shore, Nov. 26, 1939, *Yuncker* 9513; Nuu Islet, July 4, 1925, *Garber* 1112. Abundant on all of the islands, Guest. Native name, *fana'io*.

FAMILY GUTTIFERAE

Genus **Calophyllum** Linnaeus**Calophyllum inophyllum** Linnaeus, Sp. Pl., 513, 1753.

Tau: near the beach, Siufaga, Oct. 2, 1939, *Yuncker* 9194; in forest back of Luma, alt. 3 m., Jan. 11, 1922, *Garber* 704.

Ofu: in forest on the main ridge, alt. 250 m., Nov. 25, 1939, *Yuncker* 9468. Native name, *fetau*.

Calophyllum samoense Christophersen, B. P. Bishop Mus., Bull. 128: 147, fig. 20, 1935.

Abundant on all of the islands, Guest. Native name, *tamanu*.

FAMILY BIXACEAE

Genus **Bixa** Linnaeus**Bixa Orellana** Linnaeus, Sp. Pl., 512, 1753.

Tau: near dwelling, Siufaga, Oct. 3, 1939, *Yuncker* 9224. Moderate on all of the islands, Guest. Native name, *loa*.

FAMILY FLACOURTIACEAE

Genus **Flacourtia** L'Héritier**Flacourtia Rukam** Zollinger and Morren, Syst. Verz., 33, 1854.

Tau: in forest on mountain slope south of Siufaga, alt. 200 m., Sept. 27, 1939, *Yuncker* 9063; same locality, alt. 150 m., Sept. 27, 1939, *Yuncker* 9062; in forest two miles south of Siufaga, alt. 180 m., Oct. 1, 1939, *Yuncker* 9163;

plateau back of Luma, alt. 125 m., Dec. 15, 1921, Garber 646; plateau back of Luma, alt. 100 m., Sept. 21, 1922, Garber 776.

Ofu: in open forest along ridge trail, alt. 200 m., Nov. 27, 1939, Yuncker 9520; in open forest on ridge back of Ofu village, alt. 100 m., Nov. 25, 1939, Yuncker 9464. Abundant on all of the islands, Guest. Native name, *filimoto*.

FAMILY CARICACEAE

Genus *Carica* Linnaeus

Carica Papaya Linnaeus, Sp. Pl., 1036, 1753.

Occurs on all of the islands both cultivated and spontaneously. Native name, *esi*.

FAMILY BEGONIACEAE

Genus *Begonia* Linnaeus

Begonia species.

Tau: dooryard ornamental, Siufaga, Oct. 3, 1939, Yuncker 9216.

FAMILY THYMELAEACEAE

Genus *Phaleria* Jack

Phaleria acuminata (A. Gray) Gilg, in Engler and Prantl, Nat. Pflanzenfam. III, 6a: 225, 1894.

Abundant on all of the islands, Guest. Native name, *suni*.

Phaleria Burnettiana (Bentham) Knuth, Handb. Blütenbiol. III, 1: 522, 1904.

Tau: exposed cliff, Siulagi Point, alt. 75 m., Aug. 26, 1922, Garber 772. Abundant on all of the islands, Guest. Native name, *suni*.

FAMILY LYTHRACEAE

Genus *Lagerstroemia* Linnaeus

Lagerstroemia indica Linnaeus, Syst. ed. 10, 1076, 1759.

Tau: near dwelling, cultivated, Siufaga, Oct. 3, 1939, Yuncker 9201.

FAMILY LECYTHIDACEAE

Genus *Barringtonia* Forster

Barringtonia asiatica (Linnaeus) Kurz, Asiatic Soc. Beng., Jour. 45(2): 131, 1876.

Tau: abundant along the shore south of Siufaga, Sept. 29, 1939, Yuncker 9098; Fitiuta trail, alt. 125 m., Aug. 4, 1921, Garber 564. Abundant along the seacoast around all the islands, Guest. Native name, *futu*.

Barringtonia samoensis A. Gray, U. S. Expl. Exped., Bot. 15(1) : 508, 1854.

Tau: near the shore, Sept. 27, 1939, *Yuncker* 9082; edge of thicket on plateau behind Siufaga, alt. 45 m., Oct. 2, 1939, *Yuncker* 9173; along trail back of Luma, alt. 30 m., Aug. 8, 1921, *Garber* 543. Moderate to abundant on all of the islands, Guest. Native name, *falaga*.

FAMILY COMBRETACEAE

Genus *Terminalia* Linnaeus

Terminalia Catappa Linnaeus, Mant., 128, 1767.

Ofu: slope of mountain at Aaofa, alt. 15 m., May 26, 1925, *Garber* 980. Abundant on all of the islands, Guest. Native name, *talie*.

FAMILY MYRTACEAE

Genus *Psidium* Linnaeus

Psidium Guajava Linnaeus, Sp. Pl., 470, 1753.

Tau: in thicket near Siufaga, Oct. 7, 1939, *Yuncker* 9293. Native name, *kuava* [*tuava*].

Genus *Decaspermum* Forster

Decaspermum fruticosum Forster, Charact. Gen. Pl., 74, pl. 37, 1776.

Abundant on the island of Olosega, Guest. Native name, *nu'anu'a*.

Genus *Eugenia* Micheli

Eugenia corynocarpum A. Gray, U. S. Expl. Exped., Bot. 15(1) : 528, 1854, Atlas, pl. 65, 1857.

Tau: plateau back of Luma, alt. 150 m., Dec. 15, 1921, *Garber* 652, 658. Native name, *seasea*.

Eugenia inophylloides A. Gray, U. S. Expl. Exped., Bot. 15(1) : 521, 1854.

Tau: Amouli trail, alt. 200 m., Sept. 16, 1921, *Garber* 637. Abundant on all of the islands, Guest. Native names, *asi*, *asi vao*.

Eugenia malaccensis Linnaeus, Sp. Pl., 470, 1753.

Found in and around most every village, Guest. Native name, *nonu fafia*.

Eugenia neurocalyx A. Gray, U. S. Expl. Exped., Bot. 15(1) : 512, 1854, Atlas pl. 59, 1857.

On all of the islands, Guest. Native name, *fena*.

Eugenia Richii A. Gray, U. S. Expl. Exped., Bot. 15(1) : 510, 1854.

Olosega: Olosegauta-Piumafua ridge, alt. 500 m., June 17, 1925, *Garber* 1030; top of Piumafua mountain, alt. 655 m., June 17, 1925, *Garber* 1028.

FAMILY MELASTOMACEAE

Genus *Melastoma* Linnaeus

Melastoma denticulatum Labillardière, Sert. Austro-Caled. 2 : 65, pl. 64, 1825.

Tau: in thicket along trail near Fitiuta, alt. 50 m., Sept. 26, 1939, *Yuncker 9058*; edge of forest on slope south of Siufaga, alt. 240 m., Oct. 4, 1939, *Yuncker 9232*; Amouli trail, alt. 175 m., Sept. 16, 1921, *Garber 635*.

Olosega: Olosegauta-Piumafua ridge, alt. 500 m., June 17, 1925, *Garber 1037*. Native names, *fua lole*, *moegalo*, *olive vao*.

Genus *Astronidium* A. Gray

Astronidium Pickeringii (A. Gray) var. *samoense* (A. Gray) Christophersen, B. P. Bishop Mus., Bull. 154 : 32, 1938.

Tau: in forest on west slope of Mt. Vaoaimanu, alt. 360 m., Oct. 5, 1939, *Yuncker 9252*; on slopes of peak back of Siufaga, alt. 670 m., Jan. 28, 1922, *Garber 746*; in old taro clearing, below peak at back of Siufaga, alt. 530 m., Jan. 28, 1922, *Garber 721*.

Olosega: Olosegauta-Piumafua ridge, alt. 550 m., June 17, 1925, *Garber 1032*. Native name, *matoi*.

FAMILY ONAGRACEAE

Genus *Jussiaea* Linnaeus

Jussiaea erecta Linnaeus, Sp. Pl., 388, 1753.

Tau: in swampy area behind Siufaga, Oct. 7, 1939, *Yuncker 9294*.

Olosega: taro swamp back of Olosega village, June 23, 1925, *Garber 1091*. Native names, *la'avai*, *fua samasama*, *mautofu vao*.

FAMILY ARALIACEAE

Genus *Reynoldsia* A. Gray

***Reynoldsia* species.**

Tau: top of peak back of Tau village, alt. 600-700 m., Jan. 28, 1922, *Garber 752*. Abundant on all of the islands, Guest. Native name, *vi vao*.

Genus *Meryta* Forster

Meryta capitata Christophersen, B. P. Bishop Mus., Bull. 128 : 161, fig. 23, 1935.

Tau: in ravine on slope two miles south of Siufaga, alt. 100 m., Oct. 1, 1939, *Yuncker 9162*.

Ofu: in forest on ridge above Ofu village, alt. 300 m., Nov. 27, 1939, *Yuncker 9545*. Native name, *fagufagu*.

Meryta aff. macrophylla (Rich) Seemann, Bonplandia 10: 294, 1862.

Tau: Amouli trail, alt. 180 m., Sept. 16, 1921, Garber 622.

Olosega: Olosegauta-Piumafua Ridge, alt. 510 m., June 17, 1925, Garber 1079. Native name, *fagufagu*.

Genus **Polyscias** Forster

Polyscias Guilfoylei (Bull) L. H. Bailey, Rhodora 18: 153, 1916, var. (?).

Tau: ornamental shrub with coarsely toothed, rounded leaflets, Siufaga village, Oct. 3, 1939, Yuncker 9212. Native name, *tanitani*.

Polyscias Reinecke Harms, Engl. Bot. Jahrb. 25: 663, 1898.

Abundant at high elevations on Tau, Guest. Native name, *aifa*.

FAMILY UMBELLIFERAE

Genus **Centella** Linnaeus

Centella asiatica (Linnaeus) Urban, in Mart. Fl. Bras. 11(1): 287, 1879.

Tau: in sandy soil near the shore south of Siufaga, Sept. 29, 1939, Yuncker 9113; in sandy soil, Luma, July 20, 1938, W. Harris 344.

Ofu: sandy soil along the southeastern shore, Nov. 26, 1939, Yuncker 9504. Native name, *togo*.

FAMILY MYRSINACEAE

Genus **Maesa** Forskål

Maesa samoana Mez, Pflanzenr. 9(IV. 236): 53, 1902.

Tau: in thicket south of Siufaga, alt. 10 m., Sept. 29, 1939, Yuncker 9105; Luma-Faleasao trail, alt. 40 m., Jan. 8, 1922, Garber 686.

Ofu: beach at Vaoto, June 23, 1925, Garber 1097.

FAMILY PLUMBAGINACEAE

Genus **Plumbago** Linnaeus

Plumbago zeylanica Linnaeus, Sp. Pl., 151, 1753.

Olosega: along rocky shore, Lotufaga by Sili, May 29, 1925, Garber 987. Native name, *lau tafifi*.

FAMILY SAPOTACEAE

Genus **Planchonella** Pierre

Planchonella Garberi Christophersen, B. P. Bishop Mus., Bull. 128: 170, fig. 25, 1935.

Ofu: top of Nuu Islet, alt. 75 m., July 4, 1925, Garber 1101. Native name, *ala'a*.

Planchonella samoensis (Reinecke) H. J. Lam, Jard. Bot. Buit., Bull. III, 7: 218, fig. 10, 1925.

Scarce on the islands of Tau and Ofu, Guest. Native name, *mamalava*.

FAMILY EBENACEAE

Genus *Diospyros* Linnaeus

Diospyros ellipticifolia (Stokes) Bakhuizen, Gard. Bull. Straits Settlem. 7: 162, 1933.

Tau: top of bluff, north side, Siufaga Point, Aug. 26, 1922, Garber 773.

Ofu: top of Nuu Islet, alt. 75 m., July 4, 1925, Garber 1102. Native name, *anume*.

Diospyros samoensis A. Gray, Am. Acad., Proc. 5: 326, 1862.

Ofu: in thicket along southeastern shore, Nov. 26, 1939, Yuncker 9514, 9516; along beach at Toaga, June 23, 1925, Garber 1095; Nuu Islet, July 4, 1925, Garber 1103. Abundant on all of the islands, Guest. Native name, *au'aule*.

FAMILY OLEACEAE

Genus *Jasminum* Linnaeus

Jasminum didymum Forster f., Fl. Ins. Austr. Prodr., 3, 1786.

Ofu: Nuu Islet, July 4, 1925, Garber 1107. Native name, *fuefue*.

Jasminum pubescens Willdenow, Sp. Pl. 1: 37, 1792.

Tau: near dwelling in Siufaga, Oct. 3, 1939, Yuncker 9200.

Jasminum Sambac (Linnaeus) Aiton, Hort. Kew 1: 8, 1789.

Tau: in thicket near the sea two miles south of Siufaga (as escape?), Oct. 1, 1939, Yuncker 9141. Native name, *pua sosolo*.

FAMILY LOGANIACEAE

Genus *Geniostoma* Forster

Geniostoma rupestre Forster, Charact. Gen. Pl., 24, pl. 12, 1776.

Tau: in thicket on plateau behind Siufaga, alt. 45 m., Oct. 2, 1939, Yuncker 9192; in thicket, ravine south of Siufaga, alt. 50 m., Sept. 25, 1939, Yuncker 9005; peak back of Siufaga, alt. 650 m., Jan. 29, 1922, Garber 755.

Ofu: in thicket on slope behind Ofu village, alt. 45 m., Nov. 25, 1939, Yuncker 9477; in thicket on ridge back of Ofu village, alt. 200 m., Nov. 25, 1939, Yuncker 9453. Native name, *taitai ipu*.

Genus *Fagraea* Thunberg

Fagraea Berteriana A. Gray, Linn. Soc., Jour. 1: 98, 1857.

Tau: in open woods on plateau behind Siufaga, alt. 45 m., Oct. 2, 1939, Yuncker 9176; on west slope of Mt. Vaoaimanu, alt. 300 m., Oct. 5, 1939, Yuncker 9263; on plateau back of Luma, alt. 125 m., Aug. 4, 1921, Garber 580.

Ofu: in open forest on ridge above Ofu village, alt. 200 m., Nov. 27, 1939, Yuncker 9535. Moderate on all of the islands, Guest. Native name, *pua lulu*.

FAMILY APOCYNACEAE

Genus *Plumeria* Linnaeus*Plumeria acuminata* Aiton, Hort. Kew, ed. 2, 2: 70, 1811.Cultivated as an ornamental in most villages. Native name *pua fiti*.*Lochnera rosea* (Linnaeus) Reichenbach, Conspect., 134, 1828.Tau: Siufaga village, Oct. 3, 1939, *Yuncker* 9208.Genus *Alyxia* R. Brown*Alyxia bracteolosa* Rich, ex Gray, Am. Acad., Proc. 5: 332, 1862.Tau: climbing in trees in ravine on slope south of Siufaga, alt. 100 m., Sept. 25, 1939, *Yuncker* 9021; plateau back of Faleasao, alt. 200 m., Aug. 4, 1921, *Garber* 577.Ofu: climbing in trees on ridge back of Ofu village, alt. 300 m., Nov. 27, 1939, *Yuncker* 9536. Native names, *gau*, *lau maile*, *maile*.*?Alyxia stellata* (Forster) Roemer and Schultes, Syst. Veg. 4: 439, 1819.Tau: climbing in trees near Siufaga, Oct. 3, 1939, *Yuncker* 9199; near dwelling, Siufaga, Oct. 3, 1939, *Yuncker* 9227.Ofu: Feaia Point, alt. 90 m., May 25, 1925, *Garber* 977. Native names, *gau*, *gau lili'i*, *lava*, *lau maile*, *maile*.Genus *Cerbera* Linnaeus*Cerbera Manghas* Linnaeus, Sp. Pl., 208, 1753.Tau: in open forest along Luma-Falesao trail, alt. 45-50 m., Sept. 30, 1939, *Yuncker* 9134; plateau back of Luma, alt. 60 m., Aug. 2, 1921, *Garber* 546. Natiye name, *leva*.

FAMILY ASCLEPIADACEAE

Genus *Hoya* R. Brown*Hoya bicarinata* A. Gray, Am. Acad., Proc. 5: 335, 1862.Tau: climbing in thicket in ravine on slope south of Siufaga, alt. 150 m., Sept. 25, 1939, *Yuncker* 9012; Fitiuta trail, alt. 100-125 m., Aug. 4, 1921, *Garber* 556, 566; Amouli trail, alt. 100 m., Sept. 16, 1921, *Garber* 615.Ofu: climbing in trees in thicket on ridge back of Ofu village, alt. 250 m., Nov. 25, 1939, *Yuncker* 9449; climbing in trees along the southeastern shore, Nov. 26, 1939, *Yuncker* 9509. Native names, *fue se le la*, *olive vao*.*Hoya chlorantha* Rechinger, in Fedde, Repert. Nov. Sp. 5: 131, 1908.Ofu: climbing in thicket on ridge behind Ofu village, alt. 350 m., Nov. 27, 1939, *Yuncker* 9528.*Hoya samoensis* Seemann, Fl. Vit., 163, 1866.Ofu: top of Tumu mountain, alt. 450 m., Sept. 3, 1925, *Garber* 1116.

FAMILY CONVOLVULACEAE

Genus *Ipomoea* Linnaeus*Ipomoea campanulata* Linnaeus, Sp. Pl., 160, 1753.Ofu: Tua, in plantation, June 5, 1925, Garber 999. Native name, *tanamimi*.*Ipomoea gracilis* R. Brown, Prodr. Fl. Nov. Holl., 484, 1810.

Tau: climbing in thicket south of Siufaga, Sept. 25, 1939, Yuncker 9006; weed in banana plantation above Siufaga, alt. 50 m., Oct. 4, 1939, Yuncker 9234; Luma-Faleasao trail, alt. 50 m., Jan. 8, 1922, Garber 688; south side of South Cove, Papatea, on tuff at beach, July 15, 1938, W. Harris 327; on lava cliff behind Luma, alt. 45 m., July 24, 1938, W. Harris 35, 108.

Ofu: climbing in thicket near the northwestern shore, Nov. 28, 1939, Yuncker 9563; in thicket along the southeastern shore, Nov. 26, 1939, Yuncker 9508. Native names, *tanamimi vao*, *fue tanamimi*.*Ipomoea pes-caprae* (Linnaeus) Sweet, Hort. Suburb. Lond., 35, 1818.Tau: in sandy soil near the beach, south of Siufaga, Sept. 25, 1939, Yuncker 9024; sandy beach at Faga, April 20, 1922, Garber 765. Native names, *fuefue tai*, *fue moa*, *fuefue tutupatu*.*Ipomoea Quamoclit* Linnaeus, Sp. Pl., 159, 1753.

Tau: near dwelling, Siufaga, Oct. 3, 1939, Yuncker 9198; on lava rock soil, Fitiuta, alt. 30 m., Oct. 12, 1921, Garber 642.

Ipomoea Turpethum (Linnaeus) R. Brown, Prodr. Fl. Nov. Holl., 485, 1810.

Tau: in thicket, low marshy area near Siufaga, Oct. 7, 1939, Yuncker 9289; climbing in thicket near the shore, south of Siufaga, Sept. 29, 1939, Yuncker 9097; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, W. Harris 135.

Ofu: Toaga, climbing on tree at Muaava, alt. 5 m., June 22, 1925, Garber 1099. Native name, *fue vao*.

FAMILY BORAGINACEAE

Genus *Cordia* Linnaeus*Cordia subcordata* Lamarck, Tabl. Encycl. 1: 421, 1791.Moderate on the islands of Tau and Olosega. None on Ofu, Guest. Native name, *tauagave*.Genus *Messerschmidia* Hebenstreit*Messerschmidia argentea* (Linnaeus f.) Johnston, Arnold Arb., Jour. 16: 164, 1935.

Tau: rocky shore at Amouli, Sept. 16, 1921, Garber 639.

Ofu: in thicket near the shore along the southeastern trail, Nov. 26, 1939, Yuncker 9507. Abundant on all of the islands, Guest. Native name, *tausuni*.

FAMILY VERBENACEAE

Genus *Stachytarpheta* Vahl

Stachytarpheta urticaefolia (Salisbury) Sims, Curtis, Bot. Mag. 43: pl. 1848, as *S. urticifolia*, 1816.

Tau: in sandy soil near the beach south of Siufaga, Sept. 25, 1939, *Yuncker 9026**; plateau back of Luma, alt. 150 m., Dec. 15, 1921, *Garber 656*. Native names, *mautofu vao*, *mautofu tala*.

Genus *Tectona* Linnaeus f.

Tectona grandis Linnaeus f., Suppl., 151, 1781.

A few introduced specimens on Tau, Guest. Native name, *tiki*.

Genus *Premna* Linnaeus

Premna taitensis Schauer, DC Prodr. 11: 638, 1847.

Tau: in thicket on slope south of Siufaga, alt. 150 m., Sept. 29, 1939, *Yuncker 9117*; Fitiuta trail, alt. 150 m., Jan. 24, 1922, *Garber 708*.

Ofu: in thicket on slope back of Ofu village, alt. 100 m., Nov. 25, 1939, *Yuncker 9462*. Abundant on all of the islands, Guest. Native name, *aloalo*.

Genus *Vitex* Linnaeus

Vitex trifolia Linnaeus var. *bicolor* (Willdenow) Moldenke, Known Geogr. Distrib. Verb., 79, 1942.

Tau: in thicket near the shore, Sept. 29, 1939, *Yuncker 9104**; Luma, Sept. 6, 1921, *Garber 611*.

Ofu: in thicket near the shore, Nov. 28, 1939, *Yuncker 9566**; Aloafao, June 5, 1925, *Garber 995*. Moderate on all of the islands, Guest. Native name, *namulega*.

Genus *Faradaya* F. Mueller

Faradaya Powellii Seemann, Powell, Seem. Jour. Bot. 6: 342, 1868.

Tau: climbing in trees on the plateau behind Siufaga, Oct. 2, 1939, *Yuncker 9174*; Fitiuta trail above Luma, alt. 60 m., Aug. 4, 1921, *Garber 552*; Fitiuta trail back of Luma, alt. 100 m., Dec. 16, 1921, *Garber 671*; Faleulu valley, in light forest, alt. 210 m., July 27, 1938, *W. Harris 200*.

Ofu: climbing in trees on ridge back of Ofu village, alt. 250 m., Nov. 25, 1939, *Yuncker 9445*. Native name, *mamalupe*.

Genus *Clerodendrum* Linnaeus

Clerodendrum speciosissimum Van Geert ex Morren, Hort. Belg. 3: 322-323, pl. 68 [as "Clerodendron"], 1836.

Tau: in coconut plantation along Fitiuta trail, Sept. 26, 1939, *Yuncker 9059**; pig yard, Faleasao-Fitiuta trail, Jan. 8, 1922, *Garber 699**.

Ofu: in open weedy area along southeastern shore, Nov. 26, 1939, *Yuncker 9494*.

* Identified by Dr. H. N. Moldenke.

Clerodendrum inerme (Linnaeus) Gaertner, Fruct. et Semin. 1: 271, 1788.

Tau: in thicket near the shore, Sept. 29, 1939, *Yuncker* 9096; beach at Fitiuta, Aug. 8, 1921, *Garber* 584; Luma-Faleasao trail, alt. 30 m., Jan. 8, 1922, *Garber* 694.

Ofu: in thicket near the shore, Nov. 28, 1939, *Yuncker* 9562. Native name, *aloalo tai*.

Clerodendrum nutans Wallich, Num. List Dried Spec., etc., no. 1793, 1828.

Tau: near dwelling, Siufaga, Oct. 3, 1939, *Yuncker* 9228.

FAMILY LABIATAE

Genus **Leucas** R. Brown

Leucas flaccida R. Brown, Prodr. Fl. Nov. Holl., 505, 1810.

Tau: weed in sandy soil near the beach, Sept. 25, 1939, *Yuncker* 9028; on lava cliff, Luma, alt. 45 m., July 24, 1938, *W. Harris* 378; Sani Ridge back of Fitiuta, alt. 150 m., Aug. 9, 1921, *Garber* 594; on lava cliff behind Luma, alt. 45 m., July 24, 1938, *W. Harris* 36, 93.

Genus **Coleus** Loureiro

Coleus Blumei Bentham, Lab. Gen. et Sp., 56, 1832, var.?

Tau: along the beach trail south of Siufaga, Sept. 29, 1939, *Yuncker* 9099; coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 239. Native name, *la'au fai sei*.

Genus **Ocimum** Linnaeus

Ocimum sanctum Linnaeus, Mant., 85, 1767.

Tau: dooryard, Siufaga, Oct. 4, 1939, *Yuncker* 9213. Native name, *la'au sauga*.

FAMILY SOLANACEAE

Genus **Capsicum** Linnaeus

Capsicum annuum Linnaeus, Sp. Pl., 188, 1753.

Ofu: cultivated, Aloafao, alt. 7 to 8 m., June 6, 1925, *Garber* 1011. Native name, *polo papalagi*.

Capsicum frutescens Linnaeus, Sp. Pl., 189, 1753.

Tau: Siufaga, Oct. 3, 1939, *Yuncker* 9201, 9226; plateau back of Faleasao, alt. 125 m., Aug. 4, 1921, *Garber* 571; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 196.

Ofu: in sandy soil near the shore, Nov. 28, 1939, *Yuncker* 9568; cultivated, Aloafao, June 5, 1925, *Garber* 981, 994. Native names, *polo vao*, *polo papalagi*, *polo uli*.

Genus *Solanum* Linnaeus

Solanum nigrum Linnaeus, Sp. Pl., 186, 1753.

Tau: Sani Ridge back of Fitiuta, alt. 150 m., Aug. 9, 1921, *Garber* 597.

Ofu: Feaia Point, alt. 75 to 100 m., May 25, 1925, *Garber* 976. Native names, *magalogalo*, *magalo*, *polo vao*.

Genus *Cestrum* Linnaeus

Cestrum diurnum Linnaeus, Sp. Pl., 191, 1753.

Tau: cultivated, Siufaga, Oct. 6, 1939, *Yuncker* 9277.

Cestrum nocturnum Linnaeus, Sp. Pl., 191, 1753.

Tau: cultivated, Siufaga, Oct. 2, 1939, *Yuncker* 9191.

FAMILY SCROPHULARIACEAE

Genus *Russelia* Jacquin

Russelia equisetiformis Schlechtendal and Chamisso, Linnaea 6: 377, 1831.

Tau: cultivated, Siufaga, Oct. 3, 1939, *Yuncker* 9209.

Genus *Angelonia* Humboldt and Bonpland

Angelonia grandiflora C. Morren, Soc. Hort. Gand., Ann. 3: 93, pl. 119, 1847.

Tau: cultivated, Siufaga, Oct. 3, 1939, *Yuncker* 9225.

Genus *Lindernia* Allioni

Lindernia crustacea (Linnaeus) F. Mueller, Census, 97, 1882.

Tau: in wet soil along the trail two miles south of Siufaga, alt. 50 m., Oct. 1, 1939, *Yuncker* 9147; Sani Ridge back of Fitiuta, alt. 150 m., Aug. 9, 1921, *Garber* 600; wet rocks, Amouli trail, alt. 175 m., Sept. 16, 1921, *Garber* 623; south side of South Cove, Papatea, on tuff at beach, July 15, 1938, *W. Harris* 324. Native names, *lau tafifi*, *fua lanumoana*.

FAMILY GESNERIACEAE

Genus *Cyrtandra* Forster

?*Cyrtandra compressa* C. B. Clarke, DC Monogr. Phanerog. 5(1): 279, 1883.

Olosega: Olosegauta-Piumafua ridge, alt. 450 m., June 17, 1925, *Garber* 1040. Native name, *lau taputa*.

?*Cyrtandra Godeffroyi* Reinecke, Engl. Bot. Jahrb. 25: 677, 1898.

Olosega: Olosegauta-Piumafua ridge, alt. 450 m., June 17, 1925, *Garber* 1067. Native names, *fua taulou*, *atiu vao*.

Cyrtandra pulchella Rich, Gray, Am. Acad. Arts. Sci., Proc. 6: 38, 1862.

Tau: in forest on the west slope of Mt. Vaoaimanu, alt. 450 m., Oct. 5, 1939, *Yuncker* 9253; Amouli trail by a stream, alt. 180 m., Sept. 16, 1921, *Garber* 633.

Olosega: Olosegauta-Piumafua ridge, alt. 550 m., June 17, 1925, *Garber* 1059. Native names, *ma'a*, *puafafafa*, *pua lulu vao*.

Cyrtandra samoensis A. Gray, Am. Acad. Arts Sci., Proc. 6: 39, 1862.

Tau: in thicket, dark ravine, lower slope of mountain south of Siufaga, alt. 50 m., Sept. 25, 1939, Yuncker 9018; trail to plateau at back of Luma, alt. 50 m., Aug. 2, 1921, Garber 551.

Olosega: Olosegauta-Piumafua ridge, alt. 530 m., June 17, 1925, Garber 1034. Native name, *mafia*.

Genus *Gloxinia* L'Héritier

Gloxinia maculata L'Héritier, Stirp. Nov., 149, 1784-85.

Tau: cultivated, Siufaga, Oct. 3, 1939, Yuncker 9202.

FAMILY ACANTHACEAE

Genus *Thunbergia* Linnaeus

Thunbergia erecta (Bentham) T. Anderson, Linn. Soc. Bot., Jour. 7: 18, 1864.

Tau: cultivated, near dwelling, Siufaga, Oct. 3, 1939, Yuncker 9203.

Genus *Hemigraphis* Nees

Hemigraphis colorata (Blume) Hallier f., Acad. Nat. Cur., Nov. Act. 70: 199, pl. 9, fig. 1, 1897.

Tau: near dwellings, Siufaga, Oct. 3, 1939, Yuncker 9215.

Genus *Pseuderanthemum* Radlkofner

Pseuderanthemum species.

Tau: naturalized on slope south of Siufaga, alt. 240 m., Oct. 4, 1939, Yuncker 9241; Fitiuta, Aug. 9, 1921, Garber 606.

FAMILY RUBIACEAE

Genus *Hedyotis* Linnaeus

Hedyotis biflora (Linnaeus) Lamarck, Tabl. Encycl. 1: 272, 1791.

Tau: in sand near the beach south of Siufaga, Sept. 29, 1939, Yuncker 9119; cliff crevices near Faleasao, Jan. 8, 1922, Garber 698; south side of South Cove, Papatea, on tuff at beach, July 15, 1938, W. Harris 328.

Ofu: on old lava flow along northwest shore, Nov. 28, 1939, Yuncker 9555; at beach, Tua, June 7, 1925, Garber 1019.

Olosega: taro plantation, back of Olosegauta, alt. 350 m., June 17, 1925, Garber 1084.

Hedyotis foetida (Forster) J. E. Smith, in Rees Cyclop. 17(2), 1811.

Tau: cliff crevices, Faleasao-Fitiuta trail, alt. 15 m., Jan. 8, 1922, Garber 701.

Ofu: on broken lava near the shore, Nov. 27, 1939, Yuncker 9553; on exposed cliff by the sea, Ofu-Samoi, June 7, 1925, Garber 1022.

Genus **Neonauclea** Merrill

Neonauclea Forsteri (Seemann) Merrill, Washington Acad. Sci., Jour. 5: 540, 1915.

Tau: in thicket on plateau behind Siufaga, alt. 45 m., Oct. 2, 1939, *Yuncker* 9179; back of Luma, alt. 75 m., Dec. 17, 1921, *Garber* 672.

Ofu: along southeastern shore trail, Nov. 26, 1939, *Yuncker* 9506; Sina, Aloafao, alt. 15 m., June 5, 1925, *Garber* 997. Abundant on all of the islands, Guest. Native name, *ofoa*.

Genus **Mussaenda** Linnaeus

Mussaenda frondosa Linnaeus, Sp. Pl., 177, 1753.

Tau: in thicket along Fitiuta trail, Sept. 26, 1939, *Yuncker* 9043; Fitiuta trail, alt. 125 m., Aug. 4, 1921, *Garber* 562; Fitiuta trail back of Luma, alt. 175 m., Aug. 11, 1921, *Garber* 610.

Ofu: along southeastern shore trail, Nov. 26, 1939, *Yuncker* 9515. Native name, *aloalo vao*.

Genus **Tarennia** Gaertner

Tarennia sambucina (Forster f.) Durand, in Drake, Ill. Fl. Mar. Pac. 6: 190, 1890.

Tau: on plateau back of Siufaga, alt. 45 m., Oct. 2, 1939, *Yuncker* 9184; Amouli forest trail, alt. 150 m., Sept. 22, 1922, *Garber* 778.

Ofu: in forest on ridge back of Ofu village, alt. 240 m., Nov. 25, 1939, *Yuncker* 9451; in thicket on central ridge, alt. 300 m., Nov. 27, 1939, *Yuncker* 9542. Moderate to abundant on all of the islands, Guest. Native name, *ma'a-nunu*.

Genus **Canthium** Lamarck

?**Canthium Merrillii** (Setchell) Christophersen, B. P. Bishop Mus., Bull. 128: 201, 1935.

Moderate to abundant on all of the islands, Guest. Native names, *ola*, *ola sina*.

Genus **Guettarda** Linnaeus

Guettarda speciosa Linnaeus, Sp. Pl., 991, 1753.

Ofu: in thicket along southeastern shore trail, Nov. 26, 1939, *Yuncker* 9493; beach, Nuu Islet, June 4, 1925, *Garber* 991; beach, Nuu Islet, July 4, 1925, *Garber* 1109. Moderate on all of the islands, Guest. Native name, *puapua*.

Genus **Ixora** Linnaeus

Ixora coccinea Linnaeus, Sp. Pl., 110, 1753.

Tau: cultivated, Siufaga, Oct. 3, 1939, *Yuncker* 9222.

Ixora species.

Small to medium-sized tree with very hard wood. Moderate on all of the islands, Guest. Native name, *filofiloa*.

Genus **Psychotria** Linnaeus

Psychotria Garberiana Christophersen, B. P. Bishop Mus., Bull. 154 : 51, fig. 12, 1938.

Tau: below top of peak, alt. 680 m., Jan. 28, 1922, *Garber* 754.

Olosega: Olosegauta-Piumafua ridge, alt. 500 m., June 17, 1925, *Garber* 1062.

Psychotria insularum A. Gray, Am. Acad. Arts Sci., Proc. 4 : 45, 1860.

Tau: Fitiuta trail, in thicket, Sept. 26, 1939, *Yuncker* 9060; in thicket along trail south of Siufaga, alt. 20 m., Sept. 25, 1939, *Yuncker* 9008; Fitiuta trail, alt. 75 m., Aug. 4, 1921, *Garber* 555; along stream bed on plateau back of Faleasao, alt. 140 m., Aug. 4, 1921, *Garber* 573; old village site near Laufuti, alt. 45 m., May 4, 1922, *Garber* 768.

Ofu: in thicket on slope back of Ofu village, alt. 60 m., Nov. 25, 1939, *Yuncker* 9475; alt. 250 m., *Yuncker* 9444. Abundant on Olosega, Guest. Native name, *matalafi*.

Genus **Geophila** D. Don

Geophila herbacea (Jacquin) O. Kuntze, Rev. Gen., 300, 1891.

Tau: on moist soil along trail, two miles south of Siufaga, alt. 50 m., Oct. 1, 1939, *Yuncker* 9148; on moist soil near Siufaga, Oct. 7, 1939, *Yuncker* 9291; Fitiuta trail, alt. 125 m., Aug. 4, 1921, *Garber* 567; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 252. Native names, *olia'a'a*, *lafifi*.

Genus **Morinda** Linnaeus

Morinda citrifolia Linnaeus, Sp. Pl., 176, 1753.

Tau: in thicket along Fitiuta trail, Sept. 26, 1939, *Yuncker* 9032; plateau back of Luma, alt. 90 m., Aug. 4, 1922, *Garber* 771.

Ofu: in thicket on ridge near Ofu village, alt. 100 m., Nov. 25, 1939, *Yuncker* 9466. Abundant on all of the islands, Guest. Native names, *nonu*, *nonu vao*.

Genus **Sarcopygme** Setchell and Christophersen

Sarcopygme pacifica (Reinecke) Setchell and Christophersen, B. P. Bishop Mus., Occ. Papers 11(5) : 4, pl. 2, A, B, fig. 20, 1935.

Tau: in forest on west slope of Mt. Vaoaimanu, alt. 350 m., Oct. 5, 1939, *Yuncker* 9254; trail back of Luma, alt. 450 m., Jan. 28, 1922, *Garber* 760. Native names, *u'unu*, *'ulu vao*, *fagufagu*.

FAMILY CUCURBITACEAE

Genus **Melothria** Linnaeus

Melothria samoensis A. Gray, U. S. Expl. Exped., Bot. 15(1) : 641, 1854.

Tau: climbing in thicket near Siufaga, Oct. 7, 1939, *Yuncker* 9287; in thicket near the beach, Sept. 29, 1939, *Yuncker* 9095; south side of South Cove,

Papatea, lower edge of strand forest at beach, on tuff, July 15, 1938, *W. Harris* 329. Native name, *meleni vao*.

Genus *Luffa* Tournefort

Luffa cylindrica (Linnaeus) M. Roemer var. *insularum* (A. Gray) Cogniaux, DC Monog. Phanerog. 3 : 459, 1881.

Ofu: in waste area near the shore, Nov. 28, 1939, *Yuncker* 9560; along the beach at north end of Aloafao, Sept. 1925, *Garber* 1119. Native name, *meleni vao*.

Genus *Citrullus* Forskål

Citrullus vulgaris Schrader, Linnaea 12 : 412, 1838.

Ofu: in sandy soil near the beach, Nov. 28, 1939, *Yuncker* 9569. Native name, *meleni*.

Genus *Cucumis* Linnaeus

Cucumis Melo Linnaeus, Sp. Pl., 1011, 1753.

Tau: in low, waste area near Siufaga, Oct. 6, 1939, *Yuncker* 9284; at Faasamene Spring, Faleasao, alt. 4-5 m., Jan. 8, 1922, *Garber* 702. Native names, *atiu, meleni*.

FAMILY GOODENIACEAE

Genus *Scaevola* Linnaeus

Scaevola frutescens (Miller) Krause, Engler Pflanzenr. 54(IV.277) : 125, fig. 25, 1912.

Tau: forming a thicket along the strand south of Siufaga, Sept. 25, 1939, *Yuncker* 9025.

Ofu: in thicket near the shore, Nov. 26, 1939, *Yuncker* 9503. Abundant on all of the islands, Guest. Native name, *to'ito'i*.

FAMILY COMPOSITAE

Genus *Vernonia* Schreber

Vernonia cinerea (Linnaeus) Lessing, Linnaea 4 : 291, 1829.

Tau: in sandy soil near the shore south of Siufaga, Sept. 29, 1939, *Yuncker* 9116; in sand, Luma, July 19, 1938, *W. Harris* 26, 345; along the trail behind Luma, Aug. 5, 1938, *W. Harris* 17; in sand near the beach, Papatea, July 15, 1938, *W. Harris* 89, 105; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 235; on lava cliff, behind Luma, alt. 45 m., July 27, 1938, *W. Harris* 122.

Ofu: on volcanic rocks along the northwestern shore, Nov. 28, 1939, *Yuncker* 9554; along trail, Toaga, June 23, 1925, *Garber* 1096.

Genus *Adenostemma* Forster

Adenostemma Lavenia (Linnaeus) O. Kuntze, Rev. Gen. 1 : 304, 1891.

Tau: Sani Ridge back of Fitiuta, alt. 150 m., Aug. 9, 1921, *Garber* 593.

Genus *Ageratum* Linnaeus***Ageratum conyzoides*** Linnaeus, Sp. Pl., 839, 1753.

Tau: in open ground along Siufaga-Faleasao trail, alt. 50 m., Sept. 30, 1939, *Yuncker* 9124; south side of South Cove, Papatea, near the beach, July 15, 1938, *W. Harris* 330; Luma, face of lava cliff, July 24, 1938, *W. Harris* 379; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 194.

Ofu: along trail at beach, June 7, 1925, *Garber* 1021. Native name, *lau taioti*.

Genus *Zinnia* Linnaeus***Zinnia elegans*** Jacquin, Coll. 3 : 152, 1789.

Tau: cultivated, Luma, *Yuncker*; Luma, in sand near the beach, July 20, 1938, *W. Harris* 346.

Genus *Siegesbeckia* Linnaeus***Siegesbeckia orientalis*** Linnaeus, Sp. Pl., 900, 1753.

Tau: Sani Ridge, back of Fitiuta, alt. 150 m., Aug. 9, 1921, *Garber* 604.

Genus *Wedelia* Jacquin***Wedelia biflora*** (Linnaeus) DeCandolle, in Wight, Contrib. Bot. Ind., 18, 1834.

Tau: weed in coconut plantation, along trail to village of Fitiuta, Sept. 26, 1939, *Yuncker* 9054; plateau back of Luma, alt. 60 m., Aug. 2, 1921, *Garber* 547. Native name, *ateate*.

Genus *Synedrella* Gaertner***Synedrella nodiflora*** (Linnaeus) Gaertner, Fruct. et Semin. 2 : 456, pl. 171, fig. 7, 1791.

Tau: in sandy soil near the shore south of Siufaga, Sept. 29, 1939, *Yuncker* 9111; plateau back of Luma, alt. 60 m., Aug. 2, 1921, *Garber* 549; in sand, Luma, July 19, 1938, *W. Harris* 27; south side of South Cove, Papatea, on tuff near the shore, July 15, 1938, *W. Harris* 331; along trail behind Luma, Aug. 5, 1938, *W. Harris* 18; in sand near the beach, Papatea, July 15, 1938, *W. Harris* 106; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 241; on lava cliff behind Luma, alt. 45 m., July 27, 1938, *W. Harris* 37. Native names, *taioti*, *lau otioti*.

Genus *Bidens* Linnaeus***Bidens pilosa*** Linnaeus, Sp. Pl., 832, 1753.

Tau: weed in taro patch, near Siufaga, Oct. 6, 1939, *Yuncker* 9278; in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 198.

Genus **Erechtites** Rafinesque

Erechtites valerianaeifolia (Wolf) DeCandolle, Prodr. 6: 295, 1837.

Tau: in sandy soil near the beach, south of Siufaga, Sept. 25, 1939, *Yuncker* 9029; weed in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 197. Native name, *fua lele*.

Genus **Emilia** Cassini

Emilia sonchifolia (Linnaeus) DeCandolle, Prodr. 6: 302, 1837.

Tau: on old lava flow near the sea, along Luma-Faleasao trail, alt. 40-45 m., Sept. 30, 1939, *Yuncker* 9123; along trail south of Siufaga, Sept. 9, 1939, *Yuncker* 9107; weed in coconut, breadfruit, and banana plantation, Faleulu, alt. 180 m., July 27, 1938, *W. Harris* 242.

Ofu: in clearing on main ridge, alt. 250 m., Nov. 27, 1939, *Yuncker* 9529. Native name, *fua lele lili'i*.

INDEX

Native names are in italic.

<i>a'aa'a</i> 43	<i>Asparagus plumosus</i> 29	<i>Calymperes serratum</i> 8
<i>a'amati'e</i> 44	<i>Asplenium laserpitiiifolium</i> 18	<i>Calymperes strictifolium</i> 6
<i>a'atasi</i> 48	<i>Asplenium Nidus</i> 19	<i>Calymperes tahitense</i> 7
<i>Abrus precatorius</i> 42	<i>Astronium Pickeringii</i> var. <i>samoense</i> 55	<i>Calymperes Vreisii</i> 7
<i>Acalypha boehmerioides</i> 47	<i>Athyrium accedens</i> 18	<i>Cananga odorata</i> 39
<i>Acalypha grandis</i> 47	<i>atiu</i> 66	<i>Canarium samoense</i> 44
<i>Acalypha hispida</i> 47	<i>atiu vao</i> 62	<i>Canarium</i> sp. 44
<i>Acalypha Wilkesiana</i> 47	<i>atone</i> 39	<i>Canavalia rosea</i> 43
<i>Achyranthes aspera</i> 37	<i>au'aule</i> 57	<i>Canna humilis</i> 30
<i>Acroporium sigmatodontium</i> 11	<i>autasi</i> 37	<i>Canthium Merrillii</i> 64
<i>Acrostichum aureum</i> 20	<i>autie</i> 51	<i>Capsicum annuum</i> 61
<i>Adenanthera pavonina</i> 41	<i>'ava</i> 34	<i>Carica Papaya</i> 53
<i>Adenostemma Lavenia</i> 66	<i>'ava'ava</i> <i>atiu</i> 34	<i>Casuarina equisetifolia</i> 34
<i>afia</i> 56	<i>'ava inu</i> 34	<i>Ceiba pentandra</i> 51
<i>Ageratum conyzoides</i> 67	<i>'avapui</i> 30	<i>Cenchrus echinatus</i> 24
<i>Aglaia samoensis</i> 44	<i>'avasa</i> 34, 41	<i>Centella asiatica</i> 56
<i>ago</i> 30		<i>Centotheca latifolia</i> 23
<i>Agrostophyllum megalurum</i> 31	<i>Barringtonia asiatica</i> 53	<i>Cerbera Manghas</i> 58
<i>ala'a</i> 49, 56	<i>Barringtonia samoensis</i> 54	<i>Cestrum diurnum</i> 62
<i>Alectryon samoensis</i> 49	<i>Begonia</i> sp. 53	<i>Cestrum nocturnum</i> 62
<i>Aleurites moluccana</i> 46	<i>Bidens pilosa</i> 67	<i>Chariessa samoensis</i> 49
<i>aloalo</i> 60	<i>Bischofia javanica</i> 46	<i>Cheirostylis</i> sp. 31
<i>aloalo tai</i> 61	<i>Bixa orellana</i> 52	<i>Cinnamomum pallidum</i> 39
<i>aloalo vao</i> 49, 64	<i>Blechnum orientale</i> 18	<i>Citrullus vulgaris</i> 66
<i>Alphitonia zizyphoides</i> 49	<i>Boerhaavia diffusa</i> 38	<i>Citrus aurantifolia</i> 44
<i>Alpinia purpurata</i> 30	<i>Breynia nivosa</i> 45	<i>Citrus Aurantium</i> 44
<i>Alsophila extensa</i> 17	<i>Broussonetia papyrifera</i> 35	<i>Citrus Hystrix</i> 44
<i>Alternanthera sessilis</i> 37	<i>Buchanania Merrillii</i> 48	<i>Citrus maxima</i> 44
<i>Alyxia bracteolosa</i> 58	<i>Bulbophyllum Christopher-</i> <i>senii</i> 33	<i>Citrus Medica</i> 44
<i>Alyxia stellata</i> 58	<i>Bulbophyllum longiscapum</i> 33	<i>Clerodendrum inerme</i> 61
<i>Amaranthus viridis</i> 37	<i>Bulbophyllum</i> sp. 33	<i>Clerodendrum nutans</i> 61
<i>Amorphophallus campanu-</i> <i>latus</i> 28	<i>Caesalpinia jayabo</i> 41	<i>Clerodendrum speciosis-</i> <i>simum</i> 60
<i>anaoso</i> 41	<i>Caesalpinia pulcherrima</i> 41	<i>Cocos nucifera</i> 27
<i>Ananas comosus</i> 28	<i>Calanthe furcata</i> 32	<i>Codiaeum variegatum</i> var. <i>pictum</i> 47
<i>Aneilema vitiense</i> 28	<i>Calanthe triplicata</i> 32	<i>Codiaeum variegatum</i> var. <i>pictum f. appendiculatum</i> 47
<i>Angelonia grandiflora</i> 62	<i>Calanthe</i> sp. 32	<i>Coix lachryma-jobi</i> 25
<i>Angiopteris evecta</i> var. <i>Vaupelii</i> 15	<i>Callicostella papillata</i> 10	<i>Coleus Blumei</i> 61
<i>Antrophyum alatum</i> 21	<i>Calophyllum inophyllum</i> 52	<i>Colocasia esculenta</i> 28
<i>Antrophyum angustatum</i> 21	<i>Calophyllum samoense</i> 52	<i>Colubrina asiatica</i> 49
<i>anume</i> 57	<i>Calymperes Dozyanum</i> 7	<i>Commelina diffusa</i> 28
<i>aoa</i> 35	<i>Calymperes hypophylaceum</i> 6	<i>Cordia subcordata</i> 59
<i>Appendicula bracteosa</i> 31	<i>Calymperes moluccense</i> 7	<i>Cordyline terminalis</i> 28
<i>Arthropteris oblitterata</i> 20	<i>Calymperes porrectum</i> 7	<i>Crataeva religiosa</i> 40
<i>Artocarpus altilis</i> 35	<i>Calymperes samoanum</i> 6	<i>Crinum asiaticum</i> 29
<i>asi</i> 54		<i>Cucumis Melo</i> 66
<i>asi vao</i> 54		<i>Curcuma longa</i> 30

- | | | | | |
|---|--|--|---|---|
| <i>Cyathula prostrata</i> 37
<i>Cyclophorus lanceolatus</i> 20
<i>Cynodon Dactylon</i> 23
<i>Cyperus brevifolius</i> 25
<i>Cyperus compressus</i> 25
<i>Cyperus cyperoides</i> 25
<i>Cyperus cyperoides</i> subsp.
<i>cyperinus</i> 25
<i>Cyperus javanicus</i> 26
<i>Cyperus Kyllingia</i> 26
<i>Cyperus Seemannianus</i> var.
<i>altus</i> 26
<i>Cyperus Seemannianus</i> var.
<i>monostachys</i> 26
<i>Cyperus Seemannianus</i> var.
<i>perdensus</i> 26
<i>Cyrtandra compressa</i> 62
<i>Cyrtandra Godeffroyi</i> 62
<i>Cyrtandra pulchella</i> 62
<i>Cyrtandra samoensis</i> 63 | <i>Erechtites valerianaefolia</i> 68
<i>Erythrina variegata</i>
var. <i>orientalis</i> 42
<i>Erythrodess</i> sp. 30
<i>esi</i> 53
<i>Eugenia corynocarpum</i> 54
<i>Eugenia inophylloides</i> 54
<i>Eugenia malaccensis</i> 54
<i>Eugenia neurocalyx</i> 54
<i>Eugenia Richii</i> 54
<i>Euphorbia Atoto</i> 48
<i>Euphorbia hirta</i> 48
<i>Euphorbia prostrata</i> 48
<i>Eupritchardia pacifica</i> 27
<i>Evodia hortensis</i> f.
<i>simplicifolia</i> 43
<i>Evodia samoensis</i> 43
<i>Exodiptyon dentatum</i> 6
<i>Exodiptyon scabrum</i> 5

<i>fagamanu</i> 30
<i>Fagraea Berteriana</i> 57
<i>fagufagu</i> 55, 56, 65
<i>fa'i</i> 29
<i>fala 'aina</i> 28
<i>falaga</i> 76
<i>fali funo</i> 22
<i>fana'io</i> 52
<i>fanuamamala</i> 48
<i>Faradaya Powellii</i> 60
<i>fau</i> 13, 51
<i>fau tagaloa</i> 51
<i>fau'ui</i> 50
<i>fena</i> 54
<i>fetau</i> 52
<i>Ficus ciliata</i> 35
<i>Ficus Godeffroyi</i> 35
<i>Ficus obliqua</i> 35
<i>Ficus prolixa</i> 36
<i>Ficus scabra</i> 36
<i>Ficus tinctoria</i> 36
<i>filimoto</i> 53
<i>filofiloa</i> 64
<i>Fimbristylis annua</i> 26
<i>Fimbristylis cymosa</i> var.
<i>pycnocephala</i> 26
<i>Fimbristylis diphylla</i> 26
<i>fiso</i> 25
<i>fiso vao</i> 21
<i>fisoa</i> 49
<i>Fissidens mangarevensis</i> 5
<i>Flacourzia Rukam</i> 52
<i>Fleurya interrupta</i> 36
<i>Floribundaria aeruginosa</i> 9
<i>Freycinetia Hombronii</i> 21
<i>Freycinetia Reineckei</i> 22 | <i>fu'afu'a</i> 52
<i>fua fe'i it'i</i> 36, 41
<i>fua lanumoana</i> 62
<i>fua lele</i> 68
<i>fua lele lili'i</i> 68
<i>fua lili'i</i> 44, 45
<i>fua lole</i> 36, 55
<i>fua paepae</i> 44
<i>fua samasama</i> 51, 55
<i>fua taulou</i> 62
<i>fuatala</i> 50
<i>fuefue</i> 42, 43, 57
<i>fuefue fai vao</i> 43
<i>fuefue laulili'i</i> 42
<i>fuefue tai</i> 59
<i>fuefue tutupatu</i> 59
<i>fuemanogi</i> 34
<i>fue moa</i> 59
<i>fue se le la</i> 58
<i>fue tanamimi</i> 59
<i>fue vao</i> 59
<i>futu</i> 13, 53 | <i>Garovaglia Powellii</i> 9
<i>gatae</i> 42
<i>gau</i> 58
<i>gau lili'i</i> 58
<i>Geniostoma rupestre</i> 57
<i>Geophila herbacea</i> 65
<i>Glochidion ramiflorum</i>
var. <i>samoanum</i> 45
<i>Glomera Reinekeana</i> 31
<i>Gloxinia maculata</i> 63
<i>Gomphrena globosa</i> 37
<i>Gossypium brasiliense</i> 51
<i>Grewia crenata</i> 50
<i>Guettarda speciosa</i> 64
<i>Gyrocarpus americanus</i> 39 | <i>Habenaria tradescantifolia</i>
30
<i>Hedycarya denticulata</i> 39
<i>Hedyotis biflora</i> 63
<i>Hedyotis foetida</i> 63
<i>Hemigraphis colorata</i> 63
<i>Hernandia Moerenhoutiana</i>
40
<i>Hernandia ovigera</i> 40
<i>Hibiscus Abelmoschus</i> 50
<i>Hibiscus Rosa-sinensis</i> 51
<i>Hibiscus Sabdariffa</i> 51
<i>Hibiscus tiliaceus</i> 51
<i>Himantocladium implanum</i>
10
<i>Homalanthus acuminatus</i> 47 |
|---|--|--|---|---|

- Homalanthus nutans 47
 Hoya bicarinata 58
 Hoya chlorantha 58
 Hoya samoensis 58
 Humata heterophylla 20
 Hymenophyllum formosum 17
 Hymenophyllum polyanthus 17
 Hymenophyllum praeter-visum 17
 Hyophila involuta 8

 'ie'ie 22
 'ie toga 22
 ifi 42
 ififi 40
 Indigofera suffruticosa 41
 Inocarpus fagiferus 42
 Ipomoea campanulata 59
 Ipomoea gracilis 59
 Ipomoea pes-caprae 59
 Ipomoea Quamoclit 59
 Ipomoea Turpethum 59
 Ixora coccinea 64
 Ixora sp. 64

 Jasminum didymum 57
 Jasminum pubescens 57
 Jasminum Sambac 57
 Jussiaea erecta 55

 Kleinhovia hospita 52
 koko [toto] 52
 kuava [tuava] 54

 la'au fai pale 14, 18
 la'au fai moti 48
 la'au fai sei 61
 la'au fai vaitusi 39
 la'au souga 61
 la'au sosolo i le ma'a 35
 la'au talatala 37
 la'au vao 14
 la'avae 55
 lafifi 65
 lag'aali 40, 44
 lagasese 20
 Lagerstroemia indica 53
 lala 42, 47
 lama 46
 lau fala 22
 lau fala vao 27
 lau fuefue 43
 lau 'ie'ie 22

 lau lala 42
 lau lili'i 29, 45, 48
 lau magamaga 18
 lau maile 58
 lau manamana 19, 28
 lau 'ofe'ofe 23
 lau otioti 67
 lau pa 41
 lau paogo 22
 lau selesele 27
 lau tafifi 56, 62
 lau taioti 67
 lau talatala 14, 37
 lau taliga 40
 lau talatalo 21, 29
 lau tamatama 37
 lau taputu 20, 62
 laugapapa 15
 laumafafia 32
 launapapa 19
 lava 58
 Lepturus repens 23
 Leucaena glauca 41
 Leucas flaccida 61
 Leucophanes candidum 5
 Leucophanes pungens 5
 levo 58
 Lindernia crustacea 62
 Lindsaya decomposita 19
 Liparis sp. 31
 Litsia magnifolia var. samoensis 39
 loa 52
 Lochnera rosea 58
 ole vao 36
 Lomagramma cordipinna 19
 lopa 41
 lopa samoae 41
 Lourea vespertilionis 42
 Luffa cylindrica var. insularum 66
 Lycopodium carinatum 14
 Lycopodium Phlegmaria 14
 Lycopodium squarrosum 14

 ma'a 62
 ma'ali 44
 ma'anunu 64
 Macaranga Harveyana 46
 Macaranga stipulosa 47
 Macromitrium semipellucidum 9
 Maesa samoana 56
 mafia 63
 magalo 62
 magalogalo 62

 magele 35
 mago 48
 maile 58
 Malaxis resupinata 31
 malila 37
 mamala 47, 48
 mamalava 56
 mamalupe 60
 Mangifera indica 48
 Manihot esculenta 47
 manioka 47
 manutoga 37
 ma'o 51
 maota 44
 Maoutia australis 37
 Mapania longifolia 27
 Mapania macrocephala 27
 Marattia Smithii 15
 masame 45
 masoa 29
 matafa 65
 mati 36
 matoi 55
 mautofu 50
 mautofu tala 60
 mautofu vao 50, 55, 60
 mau'u'u toga 28
 Melastoma denticulatum 55
 meleni 66
 meleni vao 66
 Melochia odorata 51
 Melothria samoensis 65
 Meryta capitata 55
 Meryta macrophylla 56
 Messerschmidia argentea 59
 Microlepia sp. 19
 Micromelum minutum 43
 Microstylis sp. 31
 milo 51
 Miscanthus japonicus 25
 Mnionalia semiimbata 8
 moegalo 55
 moliaina 44
 moli patupatu 44
 moli si-suka 44
 moli tipolo 44
 moli u'u 44
 Morinda citrifolia 65
 moso'oi 39
 motu 47
 Mucuna gigantea 42
 Musa paradisiaca 29
 Mussaenda frondosa 64
 mutia 26
 Myristica inutilis 39

<i>namulega</i> 60	<i>Phreatia samoensis</i> 33	<i>pua sosolo</i> 57
<i>Nasturtium sarmentosum</i> 40	<i>Phreatia upoluensis</i> 33	<i>pu'a voi</i> 38
<i>nasu</i> 48	<i>Phyllanthus Niruri</i> 45	<i>Pueraria Thunbergiana</i> 42
<i>Neonauclea Forsteri</i> 64	<i>Phyllanthus simplex</i> 45	<i>pututai</i> 48
<i>Nephrolepis biserrata</i> 19	<i>pia</i> 29	
<i>Nephrolepis hirsutula</i> 19	<i>pilita</i> 29	<i>Reynoldsia</i> sp. 55
<i>niu</i> 27	<i>Pinnatella Kuhliana</i> 10	<i>Rhus taitensis</i> 48
<i>niu piu</i> 27	<i>Piper Graeffei</i> 34	<i>Rhynchosstegium vitianum</i> 11
<i>niu vao</i> 32	<i>Piper Graeffei</i> var. <i>cordatum</i> 34	<i>Russelia equisetiformis</i> 62
<i>nonu</i> 65	<i>Piper Macgillivrayi</i> var. <i>subrotundifolium</i> 34	<i>Rynchospora corymbosa</i> 27
<i>nonu fiafa</i> 54	<i>Piper methysticum</i> 34	
<i>nonu vao</i> 65	<i>pipi</i> 40	<i>sa'ato</i> 20
<i>nu'anu'a</i> 54	<i>Pipturus argenteus</i> 36	<i>Saccharum officinarum</i> 25
	<i>Pisonia grandis</i> 38	<i>sagisagi</i> 25
'o'a 46	<i>Pisonia umbellifera</i> 38	<i>Sarcopygme pacifica</i> 65
<i>Ocimum sanctum</i> 61	<i>Planchonella Garberi</i> 56	<i>Scaevola frutescens</i> 66
<i>ofa</i> 64	<i>Planchonella samoensis</i> 56	<i>Schizaea dichotoma</i> 15
'ofe 22	<i>Platylepis heteromorpha</i> 30	<i>Schizostachyum glaucifolium</i> 22
'ofe samoan 22	<i>Plumbago zeylanica</i> 56	<i>Scleria lithosperma</i> 27
<i>ogoogo</i> 36, 47	<i>Plumeria acuminata</i> 58	<i>Scleria scrobiculata</i> 27
<i>ola</i> 64	<i>polo papalagi</i> 61	<i>sea</i> 40
<i>ola sina</i> 64	<i>polo uli</i> 61	<i>seasea</i> 54
<i>ole tou'uta</i> 37	<i>polo vao</i> 61, 62	<i>Securinega flexuosa</i> 46
<i>olia'a</i> 65	<i>Polygala paniculata</i> 44	<i>Securinega samoana</i> 45-46
<i>oli oli</i> 15, 17, 20	<i>Polypodium blechnoides</i> 20	<i>sefa</i> 23
<i>olive vao</i> 55, 58	<i>Polypodium Scolopendria</i> 21	<i>Selaginella Reinecke</i> 14
<i>Ophioglossum pendulum</i> 15	<i>Polypodium viresi</i> 21	<i>Selaginella Whitmei</i> 15
<i>Oplismenus compositus</i> 24	<i>Polyscias Guilfoylei</i> var. 56	<i>selesele</i> 26
<i>Oplismenus undulatifolius</i> var. <i>imbecillus</i> 24	<i>Polyscias Reinecke</i> 56	<i>Sesuvium Portulacastrum</i> 38
<i>Oxalis corniculata</i> var. <i>repens</i> 43	<i>Pometia pinnata</i> 49	<i>siapo</i> 35
	<i>popo vao</i> 27	<i>Sida rhombifolia</i> 50
<i>Pandanus calostigma</i> f. <i>samoana</i> 22	<i>Portulaca lutea</i> 38	<i>Sida samoensis</i> 50
<i>Pandanus tauensis</i> 22	<i>Portulaca oleracea</i> 38	<i>Siegesbeckia orientalis</i> 67
<i>Pandanus tectorius</i> var. <i>laevis</i> 22	<i>Portulaca samoensis</i> 38	<i>soga</i> 37
<i>Panicum ambiguum</i> 24	<i>poumuli</i> 46	<i>soi</i> 29
<i>Panicum pilipes</i> 24	<i>Premna taitensis</i> 60	<i>Solanum nigrum</i> 62
<i>papaono</i> 39	<i>Procris pedunculata</i> 36	<i>so'opini</i> 43
<i>Parinarium glaberrimum</i> 40	<i>Prosaptia contigua</i> 21	<i>Sophora tomentosa</i> 41
<i>Parinarium insularum</i> 40	<i>Pseudereria ramosa</i> 33	<i>Spathoglottis plicata</i> 32
<i>Paspalum conjugatum</i> 24	<i>Pseudereria</i> sp. 33	<i>Spiridens aristifolius</i> 9
<i>Paspalum vaginatum</i> 24	<i>Pseuderanthemum</i> sp. 63	<i>Spondias dulcis</i> 48
<i>pata</i> 47	<i>Psidium Guava</i> 54	<i>Stachytarpheta urticaefolia</i> 60
<i>pata laitiuti</i> 47	<i>Psilotum complanatum</i> 14	<i>Stenotaphrum subulatum</i> 24
<i>Pelekium velatum</i> 10	<i>Psychotria Garberiana</i> 65	<i>Sterculia fanaiho</i> 52
<i>Pennisetum macrostachyum</i> 24	<i>Psychotria insularum</i> 65	<i>sun</i> 53
<i>Peperomia tutuilana</i> 35	<i>Pteris comans</i> 19	<i>Synedrella nodiflora</i> 67
<i>Phaius Tankervilliae</i> 31	<i>Pteris ensiformis</i> 19	
<i>Phaleria acuminata</i> 53	<i>Pteris tripartita</i> 20	<i>ta'ata'a</i> 23, 25
<i>Phaleria Burnettiana</i> 53	<i>pua fasafa</i> 62	<i>Tacca pinnatifida</i> 29
<i>Phreatia Reinecke</i> 33	<i>pua fiti</i> 58	<i>Taeniophyllum</i> sp. 34
	<i>pua lulu</i> 57	<i>tafa tolu</i> 27
	<i>pua lulu vao</i> 62	<i>taioti</i> 67
	<i>pu'a</i> 40	<i>taitai ipu</i> 57
	<i>puapua</i> 64	

<i>talafalu</i> 43	<i>toi</i> 49	<i>utuutu</i> 26
<i>talie</i> 54	<i>to'ito'i</i> 66	<i>u'unu</i> 65
<i>talo</i> 28	<i>tolo</i> 25	
<i>tamanu</i> 52	<i>togo</i> 56	
<i>tamole</i> 38, 39	<i>toto</i> [koko] 52	
<i>tanamimi</i> 59	<i>Trema orientalis</i> var. <i>viridis</i>	
<i>tanamimi vao</i> 59	35	
<i>tanitani</i> 56	<i>Trichomanes apiifolium</i> 16	
<i>tapioka</i> 47	<i>Trichomanes assimile</i> 16	
<i>taputo'i</i> 49	<i>Trichomanes bipunctatum</i> 16	
<i>Tarennia sambucina</i> 64	<i>Trichomanes Boryanum</i> 16	
<i>tauagave</i> 59	<i>Trichomanes dentatum</i> 16	
<i>tausuni</i> 59	<i>Trichomanes Endlicherianum</i>	
<i>tava</i> 49	16	
<i>tavai</i> 49	<i>Trichomanes humile</i> 16	
<i>Taxithelium samoanum</i> 12	<i>Trichomanes intermedium</i> 17	
<i>Taxithelium tenuisetum</i> 12	<i>Trichomanes pallidum</i> 17	
<i>Tectaria Stearnsii</i> 18	<i>Trichomanes saxifragoides</i>	
<i>Tectona grandis</i> 60	17	
<i>Tephrosia purpurea</i> 41	<i>Trichospermum Richii</i> 49	
<i>Terminalia Catappa</i> 54	<i>Triumfetta Bartramia</i> 50	
<i>teuila</i> 30	<i>Triumfetta procumbens</i> 50	
<i>teve</i> 28	<i>tuava</i> [kuava] 54	
<i>Theobroma Cacao</i> 52	<i>tuse</i> 25	
<i>Thespesia populnea</i> 51	<i>u</i> 25	
<i>Thrixspermum</i> sp. 33	<i>u'a</i> 35	
<i>Thuarea involuta</i> 25	<i>uaina</i> 51	
<i>Thuidium cymbifolium</i> 11	<i>ufi maniota</i> 47	
<i>Thuidium samoanum</i> 10	<i>'ulu</i> 35	
<i>Thunbergia erecta</i> 63	<i>'ulu vao</i> 65	
<i>ti</i> 29		
<i>tigamoni</i> 39	<i>Uraria lagopodioides</i> 42	<i>Zephyranthes rosea</i> 29
<i>tiki</i> 60	<i>Urena lobata</i> 50	<i>Zingiber Zerumbet</i> 30
<i>tipolo</i> 44	<i>usi</i> 43	<i>Zinnia elegans</i> 67
<i>toa</i> 34		