## Erratum and further notes on *Erinna newcombi* (Mollusca: Lymnaeidae) in Hanakoa, Kaua'i, Hawai'i

KENNETH R. WOOD<sup>1</sup> (National Tropical Botanical Garden, 3530 Papalina Road, Kalāheo, Kauaʻi, Hawaiʻi 96741, USA; email: kwood@ntbg.org)



FIGURE 1. Adult Erinna newcombi with young (Hanakoa, Kaua'i, Hawai'i).

The front cover photograph and Figure 1 on page 53 of Volume No. 96 (*in* Records of the Hawaii Biological Survey for 2006—Part 2: Notes) represents the Hawaiian endemic *Lymnaea aulacospira* Ancey (Lymnaeidae) and not *Erinna newcombi* Adams & Adams (R.H. Cowie pers. comm. 2008; Hubendick 1952; Boynton & Wood 2007). The correct image representing the Hanakoa, Kaua'i population of *E. newcombi* is included below (see Figure 1). Further research indicates that there are three separate sub-populations of *E. newcombi* located on the walls of fast-flowing perennial natural spring waterfalls in Hanakoa Valley. Sites range between 250 and 370 m and each site covers approximately 20 m<sup>2</sup> with densities of 5–7 snails per square meter.

## **Literature Cited**

Boynton, D. & Wood, K.R. 2007. *Erinna newcombi* Adams & Adams (Mollusca: Lymnaeidae): a rediscovered population in Hanakoa, Kaua'i, Hawai'i. *Bishop Museum Occasional Papers* **96**: 52–54.

Hubendick, B. 1952. Hawaiian Lymnaeidae. Bishop Museum Occasional Papers 20: 307–328.

Research Associate, Hawaii Biological Survey, Bishop Museum, 1525 Bernice Street, Honolulu, Hawai'i 96817, USA.